

- A comparison on Beck's cognitive therapy and jogging as treatments for depression [dissertation]. Missoula: University of Montana, 1981.
- European Heart Failure Training Group. Experience from controlled trials of physical training in chronic heart failure. Protocol and patient factors in effectiveness in the improvement in exercise tolerance. *Eur Heart J* 1998; 19: 466–475.
- Pulmonary rehabilitation. BTSS statement. *Thorax* 2001a; 56: 827–834. Working Group Report. Recommendations for exercise training in chronic heart failure patients. *Eur Heart J* 2001b; 22: 125–135.
- Abramson JL, Weintraub WS, Vaccarino V. Association between pulse pressure and C-reactive protein among apparently healthy US adults. *Hypertension* 2002; 39: 197–202.
- Adamopoulos S, Coats AJ, Brunotte F, Arnolda L, Meyer T, Thompson CH, Dunn JF, Stratton J, Kemp GJ, Rada GK. Physical training improves skeletal muscle metabolism in patients with chronic heart failure. *J Am Coll Cardiol* 1993; 21: 1101–1106.
- Adamopoulos S, Parissis J, Karatzas D, Kroupis C, Georgiadis M, Karavolias G, Paraskevaidis J, Koniavitou K, Coats AJ, Kremastinos DT. Physical training modulates proinflammatory cytokines and the soluble Fas/soluble Fas ligand system in patients with chronic heart failure. *J Am Coll Cardiol* 2002; 39: 653–663.
- Adamopoulos S, Parissis J, Kroupis C, Georgiadis M, Karatzas D, Karavolias G, Koniavitou K, Coats AJ, Kremastinos DT. Physical training reduces peripheral markers of inflammation in patients with chronic heart failure. *Eur Heart J* 2001; 22: 791–797.
- Agren B, Olin C, Castenfors J, Nilsson-Ehle P. Improvements of the lipoprotein profile after coronary bypass surgery: additional effects of an exercise training program. *Eur Heart J* 1989; 10: 451–458.
- Agurs-Collins TD, Kumanyika SK, TenHave TR, Adams-Campbell LL. A randomized controlled trial of weight reduction and exercise for diabetes management in older African-American subjects. *Diabetes Care* 1997; 20: 1503–1511.
- Ahmaidi SB, Varray AL, Savy-Pacaux AM, Prefaut CG. Cardiorespiratory fitness evaluation by the shuttle test in asthmatic subjects during aerobic training. *Chest* 1993; 103: 1135–1141.
- Akinpelu AO. Responses of the African hypertensive to exercise training: preliminary observations. *J Hum Hypertens* 1990; 4: 74–76.
- Alam S, Stolinski M, Pentecost C, Boroujerdi MA, Jones RH, Sonksen PH, Umpleby AM. The effect of a six-month exercise program on very low-density lipoprotein apolipoprotein B secretion in type 2 diabetes. *J Clin Endocrinol Metab* 2004; 89: 688–694.
- Albright A, Franz M, Hornsby G, Kriska A, Marrero D, Ullrich I, Verity LS. American College of Sports Medicine position stand. Exercise and type 2 diabetes. *Med Sci Sports Exerc* 2000; 32: 1345–1360.
- Albright CL, King AC, Taylor CB, Haskell WL. Effect of a six-month aerobic exercise training program on cardiovascular responsiveness in healthy middle-aged adults. *J Psychosom Res* 1992; 36: 25–36.
- Allenberg K, Johansen K, Saltin B. Skeletal muscle adaptations to physical training in type II (non-insulin-dependent) diabetes mellitus. *Acta Med Scand* 1988; 223: 365–373.
- American College of Sports Medicine Position stand. Physical activity, physical fitness, and hypertension. *Med Sci Sports Exerc* 1993; 25: i–x.
- American Diabetes Association Clinical practice recommendations. *Diabetes Care* 2002; Jan: S1–S147.
- American Thoracic Society Standards for the diagnosis and care of patients with COPD. *Am J Respir Crit Care Med* 1995; 152: S77–S120.
- American Thoracic Society Pulmonary rehabilitation – 1999. Official statement of the American Thoracic Society, November 1998. *Am J Respir Crit Care Med* 1999; 159: 1666–1682.
- Andersen GS, Christiansen P, Madsen S, Schmidt G. Value of regular supervised physical training after acute myocardial infarction]. *Ugeskr Laeger* 1981; 143: 2952–2955.
- Andersen RE, Wadden TA, Bartlett SJ, Zemel B, Verde TJ, Franckowiak SC. Effects of lifestyle activity vs structured aerobic exercise in obese women: a randomized trial. *JAMA* 1999; 281: 335–340.
- Anderson JW, Konz EC, Frederich RC, Wood CL. Long-term weight-loss maintenance: a meta-analysis of US studies. *Am J Clin Nutr* 2001; 74: 579–584.
- Anderssen S, Holme I, Urdal P, Hjermandt I. Diet and exercise intervention have favourable effects on blood pressure in mild hypertensives: the Oslo Diet and Exercise Study (ODES). *Blood Press* 1995; 4: 343–349.
- Anker SD, Chua TP, Ponikowski P, Harrington D, Swan JW, Poole-Wilson PA, Coats AJ. Hormonal changes and catabolic/anabolic imbalance in chronic heart failure and their importance for cardiac cachexia. *Circulation* 1997; 96: 526–534.
- Anshel MH. Effect of chronic aerobic exercise and progressive relaxation on motor performance and affect following acute stress. *Behav Med* 1996; 21: 186–196.
- Armstrong N, Simons-Morton BG. Physical activity and blood lipids in adolescents. *Pediatr Exerc* 1994; 6: 631–640.
- Arroll B, Beaglehole R. Salt restriction and physical activity in treated hypertensives. *N Z Med J* 1995; 108: 266–268.
- Asikainen TM, Miilunpalo S, Kukkonen-Harjula K, Nenonen A, Pasanen M, Rinne M, Uusi-Rasi K, Oja P, Vuori I. Walking trials in postmenopausal women: effect of low doses of exercise and exercise fractionation on coronary risk factors. *Scand J Med Sci Sports* 2003; 13: 284–292.



- Babyak M, Blumenthal JA, Herman S, Khatri P, Doraiswamy M, Moore K, Craighead WE, Baldeewicz TT, Krishnan KR. Exercise treatment for major depression: maintenance of therapeutic benefit at 10 months. *Psychosom Med* 2000; 62: 633–638.
- Bagnall AM, Whiting P, Richardson R, Sowden AJ. Interventions for the treatment and management of chronic fatigue syndrome/myalgic encephalomyelitis. *Qual Saf Health Care* 2002; 11: 284–288.
- Baker KR, Nelson ME, Felson DT, Layne JE, Sarno R, Roubeno R. The efficacy of home based progressive strength training in older adults with knee osteoarthritis: a randomized controlled trial. *J Rheumatol* 2001; 28: 1655–1665.
- Ballantyne FC, Clarke RS, Simpson HS, Ballantyne D. The effect of moderate physical exercise on the plasma lipoprotein subfractions of male survivors of myocardial infarction. *Circulation* 1982; 65: 913–918.
- Barefoot JC, Heitmann BL, Helms MJ, Williams RB, Surwit RS, Siegler IC. Symptoms of depression and changes in body weight from adolescence to mid-life. *Int J Obes Relat Metab Disord* 1998; 22: 688–694.
- Baslund B, Lyngberg K, Andersen V, Halkjaer Kristensen J, Hansen M, Klokke M, Pedersen BK. Effect of 8 wk of bicycle training on the immune system of patients with rheumatoid arthritis. *J Appl Physiol* 1993; 75: 1691–1695.
- Bauld GS, Hoaman LA, Zullo TG, Sciuaba FC. Exercise maintenance following pulmonary rehabilitation: effect of distractive stimuli. *Chest* 2002; 122: 948–954.
- Bautch JC, Clayton MK, Chu Q, Johnson KA. Synovial fluid chondroitin sulphate epitopes 3B3 and 7D4, and glycosaminoglycan in human knee osteoarthritis after exercise. *AnnRheum Dis* 2000; 59: 887–891.
- Beals CA, Lampman RM, Banwell BF, Braunstein EM, Albers JW, Castor CW. Measurement of exercise tolerance in patients with rheumatoid arthritis and osteoarthritis. *J Rheumatol* 1985; 12: 458–461.
- Bearne LM, Scott DL, Hurley MV. Exercise can reverse quadriceps sensorimotor dysfunction that is associated with rheumatoid arthritis without exacerbating disease activity. *Rheumatology (Oxford)* 2002; 41: 157–166.
- Beck-Nielsen H, Henriksen JE, Hermansen K, Madsen LD, Olivarius NF, Mandrup-Poulsen TR, Pedersen OB, Richelsen B, Schmitz OE. Type 2 diabetes and the metabolic syndrome – diagnosis and treatment. 2000; 6: 1–36. Copenhagen, Lægeföreningens forlag.
- Behnke M, Taube C, Kirsten D, Lehnigk B, Jorres RA, Magnussen H. Home-based exercise is capable of preserving hospital-based improvements in severe chronic obstructive pulmonary disease. *Respir Med* 2000; 94: 1184–1191.
- Belardinelli R, Georgiou D, Cianci G, Purcaro A. Randomized, controlled trial of long-term moderate exercise training in chronic heart failure: effects on functional capacity, quality of life, and clinical outcome. *Circulation* 1999; 99: 1173–1182.
- Belardinelli R, Georgiou D, Ginzton L, Cianci G, Purcaro A. Effects of moderate exercise training on thallium uptake and contractile response to low-dose dobutamine of dysfunctional myocardium in patients with ischemic cardiomyopathy. *Circulation* 1998; 97: 553–561.
- Belardinelli R, Georgiou D, Scocco V, Barstow TJ, Purcaro A. Low intensity exercise training in patients with chronic heart failure. *J Am Coll Cardiol* 1995; 26: 975–982.
- Belardinelli R, Paolini I, Cianci G, Piva R, Georgiou D, Purcaro A. Exercise training intervention after coronary angioplasty: the ETICA trial. *J Am Coll Cardiol* 2001; 37: 1891–1900.
- Bell JA. Comparison of a multi-disciplinary home based cardiac rehabilitation programme with comprehensive conventional rehabilitation in post-myocardial infarction patients. London: University of London, 1998.
- Bengtsson K. Rehabilitation after myocardial infarction. *Scand J Rehab Med* 1983; 15: 1–9.
- Bennell KL, Hinman RS, Metcalf BR, Buchbinder R, McConnell J, McColl G, Green S, Crossley KM. Efficacy of physiotherapy management of knee joint osteoarthritis: a randomised, double blind, placebo controlled trial. *Ann Rheum Dis* 2005; 64: 906–912.
- Bennett RM, Clark SR, Goldberg L, Nelson D, Bonafede RP, Porter J, Specht D. Aerobic fitness in patients with fibrositis. A controlled study of respiratory gas exchange and ¹³³xenon clearance from exercising muscle. *Arthritis Rheum* 1989; 32: 454–460.
- Bertie J, King A, Reed N, Marshall AJ, Ricketts C. Benefits and weaknesses of a cardiac rehabilitation programme. *J R Coll Physicians London* 1992; 26: 147–151.
- Bethell HJ, Mullee MA. A controlled trial of community based coronary rehabilitation. *Br Heart J* 1990; 64: 370–375.
- Bild DE, Sholinsky P, Smith DE, Lewis CE, Hardin JM, Burke GL. Correlates and predictors of weight loss in young adults: the CARDIA study. *Int J Obes Relat Metab Disord* 1996; 20: 47–55.
- Binder EF, Birge SJ, Kohrt WM. Effects of endurance exercise and hormone replacement therapy on serum lipids in older women. *J Am Geriatr Soc* 1996; 44: 231–236.
- Blood Pressure Lowering Treatment Trialists' Collaboration. Effects of ACE inhibitors, calcium antagonists, and other blood-pressure-lowering drugs: results of prospectively designed overviews of randomised trials. *Lancet* 2000; 355: 1955–1964.
- Blumenthal JA, Babyak MA, Moore KA, Craighead WE, Herman S, Khatri P, Waugh R, Napolitano MA, Forman LM, Appelbaum M, Doraiswamy PM, Krishnan KR. Effects of exercise training on older patients with major depression. *Arch Intern Med* 1999; 159: 2349–2356.
- Blumenthal JA, Sherwood A, Gullette EC, Babyak M, Waugh R, Georgiades A, Craighead LW, Tweedy D, Feinglos M, Appelbaum M, Hayano J, Hinderliter A. Exercise and weight loss reduce blood pressure in men and women with mild hypertension: effects on cardiovascular, metabolic, and hemodynamic functioning. *Arch Intern Med* 2000; 160: 1947–1958.
- Blumenthal JA, Siegel WC, Appelbaum M. Failure of exercise to reduce blood pressure in patients with mild hyperten-



- sion. Results of a randomized controlled trial. *JAMA* 1991; 266: 2098–2104.
- Bogardus C, Ravussin E, Robbins DC, Wolfe RR, Horton ES, Sims EA. Effects of physical training and diet therapy on carbohydrate metabolism inpatients with glucose intolerance and non-insulin-dependent diabetes mellitus. *Diabetes* 1984; 33: 311–318.
- Bonaiuto D, Shea B, Iovine R, Negrini S, Robinson V, Kemper HC, Wells G, Tugwell P, Cranney A. Exercise for preventing and treating osteoporosis in postmenopausal women. *Cochrane Database Syst Rev* 2002 CD000333.
- Booker HA. Exercise training and breathing control in patients with chronic airflow limitation. *Physiotherapy* 1984; 70: 258–260.
- Borjesson M, Robertson E, Weidenhielm L, Mattsson E, Olsson E. Physiotherapy in knee osteoarthritis: effect on pain and walking. *Physiother Res Int* 1996; 1: 89–97.
- Bosscher RJ. Running and mixed physical exercises with depressed psychiatric patients. *Int J Sport Psychol* 1993; 24: 170–184.
- Boule NG, Haddad E, Kenny GP, Wells GA, Sigal RJ. Effects of exercise on glycemic control and body mass in type 2 diabetes mellitus: a meta-analysis of controlled clinical trials. *JAMA* 2001; 286: 1218–1227.
- Boule NG, Kenny GP, Haddad E, Wells GA, Sigal RJ. Meta-analysis of the effect of structured exercise training on cardiorespiratory fitness in Type 2 diabetes mellitus. *Diabetologia* 2003; 46: 1071–1081.
- Bradham WS, Moe G, Wendt KA, Scott AA, Konig A, Romanova M, Naik G, Spinale FG. TNF-alpha and myocardial matrix metalloproteinases in heart failure: relationship to LV remodeling. *Am J Physiol Heart Circ Physiol* 2002; 282: H1288–H1295.
- Braith RW, Pollock ML, Lowenthal DT, Graves JE, Limacher MC. Moderate- and high-intensity exercise lowers blood pressure in normotensive subjects 60 to 79 years of age. *Am J Cardiol* 1994; 73: 1124–1128.
- Brandon LJ, Gaasch DA, Boyette LW, Lloyd AM. Effects of long-term resistive training on mobility and strength in older adults with diabetes. *J Gerontol A Biol Sci Med Sci* 2003; 58: 740–745.
- Brandsma JW, Robeer BG, van den HS, Smit B, Wittens CH, Oostendorp RA. The effect of exercises on walking distance of patients with intermittent claudication: a study of randomized clinical trials. *Phys Ther* 1998; 78: 278–286.
- Bravo G, Gauthier P, Roy PM, Payette H, Gaulin P, Harvey M, Peloquin L, Dubois MF. Impact of a 12-month exercise program on the physical and psychological health of osteopenic women. *J Am Geriatr Soc* 1996; 44: 756–762.
- Bremer BA, Moore CT, Bourbon BM, Hess DR, Bremer KL. Perceptions of control, physical exercise, and psychological adjustment to breast cancer in South African women. *Ann Behav Med* 1997; 19: 51–60.
- Brennan FM, Maini RN, Feldmann M. TNF alpha – a pivotal role in rheumatoid arthritis? *Br J Rheumatol* 1992; 31: 293–298.
- Brooks GA, Fahey TD, White TP. Exercise physiology: human bioenergetics and its applications. Mountain View, CA: Mayfield Publishing Company, 1995.
- Brosse AL, Sheets ES, Lett HS, Blumenthal JA. Exercise and the treatment of clinical depression in adults: recent findings and future directions. *Sports Med* 2002; 32: 741–760.
- Brown DR, Pate RR, Pratt M, Wheeler F, Buchner D, Ainsworth B, Macera C. Physical activity and public health: training courses for researchers and practitioners. *Public Health Rep* 2001; 116: 197–202.
- Buckelew SP, Conway R, Parker J, Deuser WE, Read J, Witty TE, Hewett JE, Minor M, Johnson JC, Van Male L, McIntosh MJ, Nigh M, Kay DR. Biofeedback/relaxation training and exercise interventions for fibromyalgia: a prospective trial. *Arthritis Care Res* 1998; 11: 196–209.
- Burckhardt CS, Clark SR, Padrick KP. Use of the modified Balke treadmill protocol for determining the aerobic capacity of women with fibromyalgia. *Arthritis Care Res* 1989; 2: 165–167.
- Burt VL, Whelton P, Roccella EJ, Brown C, Cutler JA, Higgins M, Horan MJ, Labarthe D. Prevalence of hypertension in the US adult population. Results from the Third National Health and Nutrition Examination Survey, 1988–1991. *Hypertension* 1995; 25: 305–313.
- Busch A, Schachter CL, Peloso PM, Bombardier C. Exercise for treating fibromyalgia syndrome. *Cochrane Database Syst Rev* 2002 CD003786.
- Busch AJ, McClements JD. Effects of a supervised home exercise program on patients with severe chronic obstructive pulmonary disease. *Phys Ther* 1988; 68: 469–474.
- Calver A, Collier J, Vallance P. Inhibition and stimulation of nitric oxide synthesis in the human forearm arterial bed of patients with insulin-dependent diabetes. *J Clin Invest* 1992; 90: 2548–2554.
- Cambach W, Chadwick-Straver RV, Wagenaar RC, van Keimpema AR, Kemper HC. The effects of a community-based pulmonary rehabilitation programme on exercise tolerance and quality of life: a randomized controlled trial. *Eur Respir J* 1997; 10: 104–113.
- Campbell K, Waters E, O'Meara S, Summerbell C. Interventions for preventing obesity in childhood. A systematic review. *Obes Rev* 2001; 2: 149–157.
- Campbell K, Waters E, O'Meara S, Kelly S, Summerbell C. Interventions for preventing obesity in children. *Cochrane Database Syst Rev* 2002 CD001871.
- Carlsen KH. Therapeutic strategies for allergic airways diseases. *Paediatr Respir Rev* 2004; 5: 45–51.
- Carlsen KH, Carlsen KC. Exercise-induced asthma. *Paediatr Respir Rev* 2002; 3: 154.
- Carlsson R. Serum cholesterol, lifestyle, working capacity and quality of life in patients with coronary artery disease. Experiences from a hospital-based secondary prevention programme. *Scand Card J* 1998; 50: 1–20.
- Carlsson R, Lindberg G, Westin L, Israelsson B. Influence of coronary nursing management follow up on lifestyle after acute myocardial infarction. *Heart* 1997; 77: 256–259.

- Carson P, Phillips R, Lloyd M, Tucker H, Neophytou M, Buch NJ, Gelson A, Lawton A, Simpson T. Exercise after myocardial infarction: a controlled trial. *J R Coll Physicians Lond* 1982; 16: 147-151.
- Carter ND, Khan KM, McKay HA, Petit MA, Waterman C, Heinonen A, Janssen PA, Donaldson MG, Mallinson A, Riddell L, Kruse K, Prior JC, Flicker L. Community-based exercise program reduces risk factors for falls in 65- to 75-year-old women with osteoporosis: randomized controlled trial. *Can Med Assoc J* 2002; 167: 997-1004.
- Carter ND, Khan KM, Petit MA, Heinonen A, Waterman C, Donaldson MG, Janssen PA, Mallinson A, Riddell L, Kruse K, Prior JC, Flicker L, McKay HA. Results of a 10 week community based strength and balance training programme to reduce fall risk factors: a randomised controlled trial in 65-75 year old women with osteoporosis. *Br J Sports Med* 2001; 35: 348-351.
- Cauza E, Hanusch-Enserer U, Strasser B, Ludvik B, Metz-Schimmerl S, Pacini G, Wagner O, Georg P, Prager R, Kostner K, Dunky A, Haber P. The relative benefits of endurance and strength training on the metabolic factors and muscle function of people with type 2 diabetes mellitus. *Arch Phys Med Rehabil* 2005; 86: 1527-1533.
- Chamberlain MA, Care G, Harfield B. Physiotherapy in osteoarthritis of the knees. A controlled trial of hospital versus home exercises. *Int Rehabil Med* 1982; 4: 101-106.
- Chavannes N, Vollenberg JJ, van Schayck CP, Wouters EF. Effects of physical activity in mild to moderate COPD: a systematic review. *Br J Gen Pract* 2002; 52: 574-578.
- Chong PF, Golledge J, Greenhalgh RM, Davies AH. Exercise therapy or angioplasty? A summation analysis. *Eur J Vasc Endovasc Surg* 2000; 20: 4-12.
- Chow R, Harrison JE, Notarius J. Effect of two randomised exercise programmes on bone mass of healthy postmenopausal women. *Br Med J (Clin Res Ed)* 1987; 295: 1441-1444.
- Cider A, Tygesson H, Hedberg M, Seligman L, Wennerblom B, Sunnerhagen KS. Peripheral muscle training in patients with clinical signs of heart failure. *Scand J Rehabil Med* 1997; 29: 121-127.
- Ciucci G, Paltriccia R, Lombardini R, Lupattelli G, Pasqualini L, Mannarino E. Treating peripheral arterial occlusive disease: pentoxifylline vs exercise. *Int Angiol* 1994; 13: 33-39.
- Clark CJ, Cochrane LM. Assessment of work performance in asthma for determination of cardiorespiratory fitness and training capacity. *Thorax* 1988; 43: 745-749.
- Clark CJ, Cochrane L, Mackay E. Low intensity peripheral muscle conditioning improves exercise tolerance and breathlessness in COPD. *Eur Respir J* 1996; 9: 2590-2596.
- Clark CJ, Cochrane LM, Mackay E, Paton B. Skeletal muscle strength and endurance in patients with mild COPD and the effects of weight training. *Eur Respir J* 2000; 15: 92-97.
- Clark SR. Prescribing exercise for fibromyalgia patients. *Arthritis Care Res* 1994; 7: 221-225.
- Clark SR, Burckhardt CS, O'Rielly C, Bennett RM. Fitness characteristics and perceived exertion in women with fibromyalgia. *J Musculoskeletal Pain* 1993; 1(3/4): 191-197.
- Clarkson P, Celermajer DS, Donald AE, Sampson M, Sorensen KE, Adams M, Yue DK, Betteridge DJ, Deanfield JE. Impaired vascular reactivity in insulin-dependent diabetes mellitus is related to disease duration and low density lipoprotein cholesterol levels. *J Am Coll Cardiol* 1996; 28: 573-579.
- Cleroux J, Feldman RD, Petrella RJ. Recommendations on physical exercise training. *Can Med Assoc J* 1999; 160.
- Coakley EH, Rimm EB, Colditz G, Kawachi I, Willett W. Predictors of weight change in men: results from the Health Professionals Follow-up Study. *Int J Obes Relat Metab Disord* 1998; 22: 89-96.
- Coats AJ, Adamopoulos S, Meyer TE, Conway J, Sleight P. Effects of physical training in chronic heart failure. *Lancet* 1990; 335: 63-66.
- Coats AJ, Adamopoulos S, Radaelli A, McCance A, Meyer TE, Bernardi L, Solda PL, Davey P, Ormerod O, Forfar C. Controlled trial of physical training in chronic heart failure. Exercise performance, hemodynamics, ventilation, and autonomic function. *Circulation* 1992; 85: 2119-2131.
- Cochrane LM, Clark CJ. Benefits and problems of a physical training programme for asthmatic patients. *Thorax* 1990; 45: 345-351.
- Cockcroft A, Berry G, Brown EB, Exall C. Psychological changes during a controlled trial of rehabilitation in chronic respiratory disability. *Thorax* 1982; 37: 413-416.
- Cockcroft AE, Saunders MJ, Berry G. Randomised controlled trial of rehabilitation in chronic respiratory disability. *Thorax* 1981; 36: 200-203.
- Coggan AR, Spina RJ, Kohrt WM, Holloszy JO. Effect of prolonged exercise on muscle citrate concentration before and after endurance training in men. *Am J Physiol* 1993; 264: E215-E220.
- Cohen-Solal A, Chabernaud JM, Gourgon R. Comparison of oxygen uptake during bicycle exercise in patients with chronic heart failure and in normal subjects. *J Am Coll Cardiol* 1990; 16: 80-85.
- Colberg S. The diabetic athlete. Prescription for exercise and sports. *Champaign, IL: Human Kinetics*, 2001.
- Collins R, MacMahon S. Blood pressure, antihypertensive drug treatment and the risks of stroke and of coronary heart disease. *Br Med Bull* 1994; 50: 272-298.
- Collins R, Peto R, MacMahon S, Hebert P, Fiebach NH, Eberlein KA, Godwin J, Qizilbash N, Taylor JO, Hennekens CH. Blood pressure, stroke, and coronary heart disease. Part 2, Short-term reductions in blood pressure: overview of randomised drug trials in their epidemiological context. *Lancet* 1990; 335: 827-838.
- Conn EH, Williams RS, Wallace AG. Exercise responses before and after physical conditioning in patients with severely depressed left ventricular function. *Am J Cardiol* 1982; 49: 296-300.



- Cononie CC, Graves JE, Pollock ML, Phillips MI, Summers C, Hagberg JM. Effect of exercise training on blood pressure in 70- to 79-yr-old men and women. *Med Sci Sports Exerc* 1991; 23: 505–511.
- Conraads VM, Beckers P, Bosmans J, De Clerck LS, Stevens WJ, Vrints CJ, Brutsaert DL. Combined endurance/ resistance training reduces plasma TNF-alpha receptor levels in patients with chronic heart failure and coronary artery disease. *Eur Heart J* 2002; 23: 1854–1860.
- Cook NR, Cohen J, Hebert PR, Taylor JO, Hennekens CH. Implications of small reductions in diastolic blood pressure for primary prevention. *Arch Intern Med* 1995; 155: 701–709.
- Cooper AR, Moore LA, McKenna J, Riddoch CJ. What is the magnitude of blood pressure response to a programme of moderate intensity exercise? Randomised controlled trial among sedentary adults with unmedicated hypertension. *Br J Gen Pract* 2000; 50: 958–962.
- Cornelissen VA, Fagard RH. Effects of endurance training on blood pressure, blood pressure-regulating mechanisms, and cardiovascular risk factors. *Hypertension* 2005; 46: 667–675.
- Corra U, Giannuzzi P, Adamopoulos S, Bjornstad H, Bjarnason-Wehrens B, Cohen-Solal A, Dugmore D, Fioretti P, Gaita D, Hambrecht R, Hellermans I, McGee H, Mendes M, Perk J, Saner H, Vanhees L. Executive summary of the working group on cardiac rehabilitation and exercise physiology of the European Society of Cardiology (ESC): core components of cardiac rehabilitation in chronic heart failure. *Eur J Cardiovasc Prev Rehabil* 2005; 12: 321–325.
- Corvera-Tindel T, Doering LV, Woo MA, Khan S, Dracup K. Effects of a home walking exercise program on functional status and symptoms in heart failure. *Am Heart J* 2004; 147: 339–346.
- Courneya KS, Friedenreich CM. Relationship between exercise during cancer treatment and current quality of life in survivors of breast cancer. *J Psychosocial Oncol* 1997a; 5: 120–127.
- Courneya KS, Friedenreich CM. Relationship between exercise pattern across the cancer experience and current quality of life in colorectal cancer survivors. *J Altern Complement Med* 1997b; 3: 215–226.
- Courneya KS, Friedenreich CM. Physical exercise and quality of life following cancer diagnosis: a literature review. *Ann Behav Med* 1999; 21: 171–179.
- Courneya KS, Friedenreich CM, Quinney HA, Fields AL, Jones LW, Fairey AS. A randomized trial of exercise and quality of life in colorectal cancer survivors. *Eur J Cancer Care (Engl)* 2003; 12: 347–357.
- Courneya KS, Mackey JR, Jones LW. Coping with cancer experience: can physical exercise help? *Physician Sportsmed* 2000; 28: 49–73.
- Coutinho M, Gerstein HC, Wang Y, Yusuf S. The relationship between glucose and incident cardiovascular events. A meta-regression analysis of published data from 20 studies of 95, 783 individuals followed for 12.4 years. *Diabetes Care* 1999; 22: 233–240.
- Cox KL, Puddey IB, Morton AR, Burke V, Beilin LJ, McAleer M. Exercise and weight control in sedentary overweight men: effects on clinic and ambulatory blood pressure. *J Hypertens* 1996; 14: 779–790.
- Crawford DA, Jefferies RW, French SA. Television viewing, physical inactivity and obesity. *Int J Obes Relat Metab Disord* 1999; 23: 437–440.
- Creasy TS, McMillan PJ, Fletcher EW, Collin J, Morris PJ. Is percutaneous transluminal angioplasty better than exercise for claudication? Preliminary results from a prospective randomised trial. *Eur J Vasc Surg* 1990; 4: 135–140.
- Crouse SF, O'Brien BC, Grandjean PW, Lowe RC, Rohack JJ, Green JS, Tolson H. Training intensity, blood lipids, and apolipoproteins in men with high cholesterol. *J Appl Physiol* 1997; 82: 270–277.
- Cummings SR, Nevitt MC, Browner WS, Stone K, Fox KM, Ensrud KE, Cauley J, Black D, Vogt TM. Risk factors for hip fracture in white women. Study of Osteoporotic fractures research group. *N Engl J Med* 1995; 332: 767–773.
- Dahllof AG, Bjornorp P, Holm J, Schersten T. Metabolic activity of skeletal muscle in patients with peripheral arterial insufficiency. *Eur J Clin Invest* 1974; 4: 9–15.
- Dansk Lungemedicinsk Selskab and Dansk Selskab for Almen Medicin (1998). Kronisk obstruktiv lungesygdom. *Ugeskr. Laeger*.
- Dansk Selskab for Adipositasforskning and Dansk Kirurgisk Selskab (2001). Er der indikation for kirurgisk behandling af ekstrem overvægt i Danmark? *Ugeskr. Laeger*. Davey P, Meyer T, Coats A, Adamopoulos S, Casadei B, Conway J, Sleight P. Ventilation in chronic heart failure: effects of physical training. *Br Heart J* 1992; 68: 473–477.
- Day L, Fildes B, Gordon I, Fitzharris M, Flamer H, Lord S. Randomised factorial trial of falls prevention among older people living in their own homes. *BMJ* 2002; 325: 128.
- de Geus EJ, Klufft C, de Bart AC, van Doornen LJ. Effects of exercise training on plasminogen activator inhibitor activity. *Med Sci Sports Exerc* 1992; 24: 1210–1219.
- de Jong Z, Munneke M, Lems WF, Zwinderman AH, Kroon HM, Pauwels EK, Jansen A, Ronday KH, Dijkmans BA, Breedveld FC, Vliet Vlieland TP, Hazes JM. Slowing of bone loss in patients with rheumatoid arthritis by long-term high-intensity exercise: results of a randomized, controlled trial. *Arthritis Rheum* 2004; 50: 1066–1076.
- de Jong Z, Munneke M, Zwinderman AH, Kroon HM, Jansen A, Ronday KH, van Schaardenburg D, Dijkmans BA, Van Den Ende CH, Breedveld FC, Vliet Vlieland TP, Hazes JM. Is a long-term high-intensity exercise program effective and safe in patients with rheumatoid arthritis? Results of a randomized controlled trial. *Arthritis Rheum* 2003; 48: 2415–2424.
- De Plaen JF, Detry JM. Hemodynamic effects of physical training in established arterial hypertension. *Acta Cardiol*

- 1980; 35: 179–188.
- de Vries SO, Visser K, de Vries JA, Wong JB, Donaldson MC, Hunink MG. Intermittent claudication: cost- effectiveness of revascularization versus exercise therapy. *Radiology* 2002; 222: 25–36.
- Dela F, Handberg A, Mikines KJ, Vinten J, Galbo H. GLUT 4 and insulin receptor binding and kinase activity in trained human muscle. *J Physiol* 1993; 469(615–24): 615–624.
- Dela F, Larsen JJ, Mikines KJ, Ploug T, Petersen LN, Galbo H. Insulin- stimulated muscle glucose clearance in patients with NIDDM. Effects of one- legged physical training. *Diabetes* 1995; 44: 1010–1020.
- Dela F, Ploug T, Handberg A, Petersen LN, Larsen JJ, Mikines KJ, Galbo H. Physical training increases muscle GLUT4 protein and mRNA in patients with NIDDM. *Diabetes* 1994; 43: 862– 865.
- Dela F, von Linstow ME, Mikines KJ, Galbo H. Physical training may enhance beta-cell function in type 2 diabetes. *Am J Physiol Endocrinol Metab* 2004; 287: E1024–E1031.
- Delagardelle C, Feiereisen P, Krecke R, Essamri B, Beissel J. Objective effects of a 6 months' endurance and strength training program in outpatients with congestive heart failure. *Med Sci Sports Exerc* 1999; 31: 1102–1107.
- Demopoulos L, Bijou R, Fergus I, Jones M, Strom J, LeJemtel TH. Exercise training in patients with severe congestive heart failure: enhancing peak aerobic capacity while minimizing the increase in ventricular wall stress. *J Am Coll Cardiol* 1997; 29: 597–603.
- DePue JD, Clark MM, Ruggiero L, Medeiros ML, Pera V. Jr Maintenance of weight loss: a needs assessment. *Obes Res* 1995; 3: 241–248.
- Derby CA, Mohr BA, Goldstein I, Feldman HA, Johannes CB, McKinlay JB. Modifiable risk factors and erectile dysfunction: can lifestyle changes modify risk? *Urology* 2000; 56: 302– 306.
- Derman WE, Coleman KL, Noakes TD. Effects of exercise training in patients with cancer who have undergone chemotherapy. *Med Sci Sports Exerc* 1999; 31: 368.
- Deyle GD, Henderson NE, Matekel RL, Ryder MG, Garber MB, Allison SC. Effectiveness of manual physical therapy and exercise in osteoarthritis of the knee. A randomized, controlled trial. *Ann Intern Med* 2000; 132: 173–181.
- Di GX, Teng WP, Zhang J, Fu PY. Exercise therapy of non-insulin dependent diabetes mellitus a report of 10 year studies. The efficacy of exercise therapy. *Chin Med J (Engl)* 1993; 106: 757–759.
- Dietz JH. *Rehabilitation oncology*. New York: Wiley, 1981.
- Dimeo F, Rumberger BG, Keul J. Aerobic exercise as therapy for cancer fatigue. *Med Sci Sports Exerc* 1998; 30: 475–478.
- Dimeo FC. Effects of exercise on cancer- related fatigue. *Cancer* 2001; 92: 1689– 1693.
- Dimeo FC, Stieglitz RD, Novelli-Fischer U, Fetscher S, Keul J. Effects of physical activity on the fatigue and psychological status of cancer patients during chemotherapy. *Cancer* 1999; 85: 2273–2277.
- Dimeo FC, Tilmann MH, Bertz H, Kanz L, Mertelsmann R, Keul J. Aerobic exercise in the rehabilitation of cancer patients after high dose chemotherapy and autologous peripheral stem cell transplantation. *Cancer* 1997; 79: 1717–1722.
- Dinnes J, Kleijnen J, Leitner M, Thompson D. Cardiac rehabilitation. *Qual Health Care* 1999; 8: 65–71.
- Donner CF, Howard P. Pulmonary rehabilitation in chronic obstructive pulmonary disease (COPD) with recommendations for its use. Report of the European Respiratory Society Rehabilitation and Chronic Care Scientific Group (S.E.P.C.R. Rehabilitation Working Group). *Eur Respir J* 1992; 5: 266–275.
- Doyle EJ, Ossip-Klein DJ, Bowman ED, Osborn KM, McDougall-Wilson IB, Neimeyer RA. Running versus weight lifting in the treatment of depression. *J Consult Clin Psychol* 1987; 55: 748–754.
- Dubach P, Myers J, Dziekan G, Goebbels U, Reinhart W, Muller P, Buser P, Stulz P, Vogt P, Ratti R. Effect of high intensity exercise training on central hemodynamic responses to exercise in men with reduced left ventricular function. *J Am Coll Cardiol* 1997; 29: 1591–1598.
- Duey WJ, O'Brien WL, Crutchfield AB, Brown LA, Williford HN, Sharf- Olson M. Effects of exercise training on aerobic fitness in African-American females. *Ethn Dis* 1998; 8: 306–311.
- Dugmore LD, Tipson RJ, Phillips MH, Flint EJ, Stentiford NH, Bone MF, Littler WA. Changes in cardiorespiratory fitness, psychological wellbeing, quality of life, and vocational status following a 12 month cardiac exercise rehabilitation programme. *Heart* 1999; 81: 359–366.
- Duncan BB, Schmidt MI. Chronic activation of the innate immune system may underlie the metabolic syndrome. *Sao Paulo Med J* 2001; 119: 122–127.
- Duncan JJ, Farr JE, Upton SJ, Hagan RD, Oglesby ME, Blair SN. The effects of aerobic exercise on plasma catecholamines and blood pressure in patients with mild essential hypertension. *JAMA* 1985; 254: 2609– 2613.
- Duncan JJ, Gordon NF, Scott CB. Women walking for health and fitness. How much is enough? *JAMA* 1991; 266: 3295–3299.
- Dunn AL, Trivedi MH, O'Neal HA. Physical activity dose-response effects on outcomes of depression and anxiety. *Med Sci Sports Exerc* 2001; 33: S587– S597.
- Dunn AL, Trivedi MH, Kampert JB, Clark CG, Chambliss HO. The DOSE study: a clinical trial to examine efficacy and dose response of exercise as treatment for depression. *Control Clin Trials* 2002; 23: 584–603.
- Dunn AL, Trivedi MH, Kampert JB, Clark CG, Chambliss HO. Exercise treatment for depression: efficacy and dose response. *Am J Prev Med* 2005; 28: 1–8.
- Dunstan DW, Mori TA, Puddey IB, Beilin LJ, Burke V, Morton AR, Stanton KG. The independent and combined effects of aerobic exercise and dietary fish intake on serum lipids and glycemic control in NIDDM. A randomized controlled study. *Diabetes Care* 1997; 20: 913–921.
- Dunstan DW, Puddey IB, Beilin LJ, Burke V, Morton AR, Stanton KG. Effects of a short-term circuit weight training



- program on glycaemic control in NIDDM. *Diabetes Res Clin Pract* 1998; 40: 53–61.
- Durstine JL, Haskell WL. Effects of exercise training on plasma lipids and lipoproteins. *Exerc Sport Sci Rev* 1994; 22(477–521): 477–521.
- Ebeling P, Bourey R, Koranyi L, Tuominen JA, Groop LC, Henriksson J, Mueckler M, Sovijarvi A, Koivisto VA. Mechanism of enhanced insulin sensitivity in athletes. Increased blood flow, muscle glucose transport protein (GLUT-4) concentration, and glycogen synthase activity. *J Clin Invest* 1993; 92: 1623–1631.
- Ebrahim S, Smith GD. Lowering blood pressure: a systematic review of sustained effects of non-pharmacological interventions. *J Public Health Med* 1998; 20: 441–448.
- Ebrahim S, Thompson PW, Baskaran V, Evans K. Randomized placebo-controlled trial of brisk walking in the prevention of postmenopausal osteoporosis. *Age Ageing* 1997; 26: 253–260.
- Eid AA, Ionescu AA, Nixon LS, Lewis-Jenkins V, Matthews SB, Grieths TL, Shale DJ. Inflammatory response and body composition in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2001; 164: 1414–1418.
- Eklom B, Lovgren O, Alderin M, Fridstrom M, Satterstrom G. Physical performance in patients with rheumatoid arthritis. *Scand J Rheumatol* 1974; 3: 121–125.
- Ekdahl C, Broman G. Muscle strength, endurance, and aerobic capacity in rheumatoid arthritis: a comparative study with healthy subjects. *Ann Rheum Dis* 1992; 51: 35–40.
- Elliott TG, Cockcroft JR, Groop PH, Viberti GC, Ritter JM. Inhibition of nitric oxide synthesis in forearm vasculature of insulin-dependent diabetic patients: blunted vasoconstriction in patients with microalbuminuria. *Clin Sci (Lond)* 1993; 85: 687–693.
- Emtner M, Finne M, Stalenheim G. A 3-year follow-up of asthmatic patients participating in a 10-week rehabilitation program with emphasis on physical training. *Arch Phys Med Rehabil* 1998; 79: 539–544.
- Emtner M, Herala M, Stalenheim G. High-intensity physical training in adults with asthma. A 10-week rehabilitation program. *Chest* 1996; 109: 323–330.
- Engblom E, Hamalainen H, Lind J, Mattlar CE, Ollila S, Kallio V, Inberg M, Knuts LR. Quality of life during rehabilitation after coronary artery bypass surgery. *Qual Life Res* 1992a; 1: 167–175.
- Engblom E, Hietanen EK, Hamalainen H, Kallio V, Inberg M, Knuts LR. Exercise habits and physical performance during comprehensive rehabilitation after coronary artery bypass surgery. *Eur Heart J* 1992b; 13: 1053–1059.
- Engblom E, Korpilahi K, Hamalainen H, Puukka P, Ronnema T. Effects of five years of cardiac rehabilitation after coronary artery bypass grafting on coronary risk factors. *Am J Cardiol* 1996; 78: 1428–1431.
- Engblom E, Korpilahi K, Hamalainen H, Ronnema T, Puukka P. Quality of life and return to work 5 years after coronary artery bypass surgery. Long-term results of cardiac rehabilitation. *J Cardiopulm Rehabil* 1997; 17: 29–36.
- Engblom E, Ronnema T, Hamalainen H, Kallio V, Vanttinen E, Knuts LR. Coronary heart disease risk factors before and after bypass surgery: results of a controlled trial on multifactorial rehabilitation. *Eur Heart J* 1992c; 13: 232–237.
- Epstein D. Aerobic activity versus group cognitive therapy: an evaluative study of contrasting interventions for the alleviation of clinical depression [dissertation]. Reno: University of Nevada, 1986.
- Epstein LH, Goldfield GS. Physical activity in the treatment of childhood overweight and obesity: current evidence and research issues. *Med Sci Sports Exerc* 1999; 31: S553–S559.
- Erdman RA, Duivenvoorden HJ, Verhage F, Kazmier M, Hugenholtz PG. Predictability of beneficial effects in cardiac rehabilitation: a randomized clinical trial of psychosocial variables. *J Cardiopulm Rehabil* 1986; 6: 206–213.
- Eriksson J, Tuominen J, Valle T, Sundberg S, Sovijarvi A, Lindholm H, Tuomilehto J, Koivisto V. Aerobic endurance exercise or circuit-type resistance training for individuals with impaired glucose tolerance? *Horm Metab Res* 1998; 30: 37–41.
- Eriksson KF, Lindgarde F. No excess 12-year mortality in men with impaired glucose tolerance who participated in the Malmo Preventive Trial with diet and exercise. *Diabetologia* 1998; 41: 1010–1016.
- Ernst E. Exercise for female osteoporosis. A systematic review of randomised clinical trials. *Sports Med* 1998; 25: 359–368.
- Ernst EE, Matrai A. Intermittent claudication, exercise, and blood rheology. *Circulation* 1987; 76: 1110–1114.
- Esler M, Rumanir M, Kaye D, Lambert G. The sympathetic neurobiology of essential hypertension: disparate influences of obesity, stress, and noradrenaline transporter dysfunction? *Am J Hypertens* 2001; 14: 139S–146S.
- Espallargues M, Sampietro-Colom L, Estrada MD, Sola M, del Rio L, Setoain J, Granados A. Identifying bone-mass-related risk factors for fracture to guide bone densitometry measurements: a systematic review of the literature. *Osteoporos Int* 2001; 12: 811–822.
- Esposito K, Giugliano F, Di Palo C, Giugliano G, Marfella R, D'Andrea F, D'Armiento M, Giugliano D. Effect of lifestyle changes on erectile dysfunction in obese men: a randomized controlled trial. *JAMA* 2004; 291: 2978–2984.
- Ettinger WH Jr, Burns R, Messier SP, Applegate W, Rejeski WJ, Morgan T, Shumaker S, Berry MJ, O'Toole M, Monu J, Craven T. A randomized trial comparing aerobic exercise and resistance exercise with a health education program in older adults with knee osteoarthritis. The Fitness Arthritis and Seniors Trial (FAST). *JAMA* 1997; 277: 25–31.
- European Respiratory Society. Optimal assessment and management of COPD. *Eur Respir J* 1995; 8: 1398–1420.
- Ewbank PP, Darga LL, Lucas CP. Physical activity as a predictor of weight maintenance in previously obese subjects. *Obes Res* 1995; 3: 257–263.
- Fagard RH. Physical activity in the prevention and treatment of hypertension in the obese. *Med Sci Sports Exerc* 1999; 31: S624–S630.
- Fagard RH. Exercise characteristics and the blood pressure response to dynamic physical training. *Med Sci Sports*

- Exerc 2001: 33: S484-S492.
- Fairfield KM., Fletcher RH. Vitamins for chronic disease prevention in adults: scientific review. *JAMA* 2002: 287: 3116-3126.
- Falkenbach A. Physical exercise, nutrition and sunshine exposure for the prevention of osteoporosis. *Forsch Komplementarmed Klass Naturheilkd* 2001: 8: 196-204.
- Farmer ME, Harris T, Madans JH, Wallace RB, Cornoni-Huntley J, White LR. Anthropometric indicators and hip fracture. The NHANES I epidemiologic follow-up study. *J Am Geriatr Soc* 1989: 37: 9-16.
- Febbraio MA, Pedersen BK. Muscle-derived interleukin-6: mechanisms for activation and possible biological roles. *FASEB J* 2002: 16: 1335-1347.
- Feinleib M, Rosenberg HM, Collins JG, Delozier JE, Pokras R, Chevarley FM. Trends in COPD morbidity and mortality in the United States. *Am Rev Respir Dis* 1989: 140: S9-18.
- Felson DT, Naimark A, Anderson J, Kazis L, Castelli W, Meenan RF. The prevalence of knee osteoarthritis in the elderly. The Framingham Osteoarthritis Study. *Arthritis Rheum* 1987: 30: 914-918.
- Ferrier KE, Waddell TK, Gatzka CD, Cameron JD, Dart AM, Kingwell BA. Aerobic exercise training does not modify large-artery compliance in isolated systolic hypertension. *Hypertension* 2001: 38: 222-226.
- Fitch KD, Blitvich JD, Morton AR. The effect of running training on exercise-induced asthma. *Ann Allergy* 1986: 57: 90-94.
- Fletcher BJ, Dunbar SB, Felner JM, Jensen BE, Almon L, Cotsonis G, Fletcher GF. Exercise testing and training in physically disabled men with clinical evidence of coronary artery disease. *Am J Cardiol* 1994: 73: 170-174.
- Flynn TJ, Walsh MF. Thirty-month evaluation of a popular very-low-calorie diet program. *Arch Fam Med* 1993: 2: 1042-1048.
- Fogelholm M, Kukkonen-Harjula K. Does physical activity prevent weight gain - a systematic review. *Obes Rev* 2000: 1: 95-111.
- Fogelholm M, Kukkonen-Harjula K, Nenonen A, Pasanen M. Effects of walking training on weight maintenance after a very-low-energy diet in premenopausal obese women: a randomized controlled trial. *Arch Intern Med* 2000: 160: 2177-2184.
- Fortmann SP, Haskell WL, Wood PD. Effects of weight loss on clinic and ambulatory blood pressure in normotensive men. *Am J Cardiol* 1988: 62: 89-93.
- Foxx ML, Keteyian SJ. Fox's physiological basis for exercise physiology. New York: McGraw-Hill Co, 1998.
- Fremont J, Wilcoxon Craighead L. Aerobic exercise and cognitive therapy in the treatment of dysphoric moods. *Cognitive Ther Res* 1987: 11: 241-251.
- Fridlund B, Hogstedt B, Lidell E, Larsson PA. Recovery after myocardial infarction. Effects of a caring rehabilitation programme. *Scand J Caring Sci* 1991: 5: 23-32.
- Fuchsjaeger-Mayrl G, Pleiner J, Wiesinger GF, Sieder AE, Quittan M, Nuhr MJ, Francesconi C, Seit HP, Francesconi M, Schmetterer L, Wolzt M. Exercise training improves vascular endothelial function in patients with type 1 diabetes. *Diabetes Care* 2002: 25: 1795-1801.
- Fujii S, Okuno Y, Okada K, Tanaka S, Seki J, Wada M, Iseki T. Effects of physical training on glucose tolerance and insulin response in diabetics. *Osaka City Med J* 1982: 28: 1-8.
- Fulcher KY, White PD. Randomised controlled trial of graded exercise in patients with the chronic fatigue syndrome. *BMJ* 1997: 314: 1647-1652.
- Fulcher KY, White PD. Strength and physiological response to exercise in patients with chronic fatigue syndrome. *J Neurol Neurosurg Psychiatry* 2000: 69: 302-307.
- Gaede P, Vedel P, Larsen N, Jensen GV, Parving HH, Pedersen O. Multifactorial intervention and cardiovascular disease in patients with type 2 diabetes. *N Engl J Med* 2003: 348: 383-393.
- Ga□ney FA, Grimby G, Danneskiold-Samsoe B, Halskov O. Adaptation to peripheral muscle training. *Scand J Rehabil Med* 1981: 13: 11-16.
- Galipeau DM, Yao L, McNeill JH. Relationship among hyperinsulinemia, insulin resistance, and hypertension is dependent on sex. *Am J Physiol Heart Circ Physiol* 2002: 283: H562-H567.
- Gardner AW, Poehlman ET. Exercise rehabilitation programs for the treatment of claudication pain. A meta-analysis. *JAMA* 1995: 274: 975-980.
- Gardner AW, Katzell LI, Sorkin JD, Goldberg AP. Effects of long-term exercise rehabilitation on claudication distances in patients with peripheral arterial disease: a randomized controlled trial. *J Cardiopulm Rehabil* 2002: 22: 192-198.
- Garfinkel SK, Kesten S, Chapman KR, Rebeck AS. Physiologic and nonphysiologic determinants of aerobic fitness in mild to moderate asthma. *Am Rev Respir Dis* 1992: 145: 741-745.
- Geddes EL, Reid WD, Crowe J, O'Brien K, Brooks D. Inspiratory muscle training in adults with chronic obstructive pulmonary disease: a systematic review. *Respir Med* 2005.
- Gettman LR, Pollock ML, Durstine JL, Ward A, Ayres J, Linnerud AC. Physiological responses of men to 1, 3, and 5 day per week training programs. *Res Q* 1976: 47: 638-646.
- Giannuzzi P, Mezzani A, Saner H, Bjornstad H, Fioretti P, Mendes M, Cohen-Solal A, Dugmore L, Hambrecht R, Hellemans I, McGee H, Perk J, Vanhees L, Veress G. Physical activity for primary and secondary prevention. Position paper of the working group on cardiac rehabilitation and exercise physiology of the European society of cardiology. *Eur J Cardiovasc Prev Rehabil* 2003: 10: 319-327.
- Gielen S, Adams V, Mobius-Winkler S, Linke A, Erbs S, Yu J, Kempf W, Schubert A, Schuler G, Hambrecht R. Anti-inflammatory effects of exercise training in the skeletal muscle of patients with chronic heart failure. *J Am Coll Cardiol* 2003: 42: 861-868.
- Gillespie LD, Gillespie WJ, Robertson MC, Lamb SE, Cumming RG, Rowe BH. Interventions for preventing falls in



- elderly people. *Cochrane Database Syst Rev* 2001 CD000340.
- Girodo M, Ekstrand KA, Metivier GJ. Deep diaphragmatic breathing: rehabilitation exercises for the asthmatic patient. *Arch Phys Med Rehabil* 1992; 73: 717-720.
- Gokce N, Vita JA, Bader DS, Sherman DL, Hunter LM, Holbrook M, O'Malley C, Keaney JF Jr., Balady GJ. Effect of exercise on upper and lower extremity endothelial function in patients with coronary artery disease. *Am J Cardiol* 2002; 90: 124-127.
- Goldbourt U, Yaari S, Medalie JH. Factors predictive of long-term coronary heart disease mortality among 10, 059 male Israeli civil servants and municipal employees. A 23-year mortality follow-up in the Israeli Ischemic Heart Disease Study. *Cardiology* 1993; 82: 100-121.
- Goldstein RS, Gort EH, Stubbing D, Avendano MA, Guyatt GH. Randomised controlled trial of respiratory rehabilitation. *Lancet* 1994; 344: 1394-1397.
- Gordon A., Tyni-Lenne R, Persson H, Kaijser L, Hultman E, Sylven C. Markedly improved skeletal muscle function with local muscle training in patients with chronic heart failure. *Clin Cardiol* 1996; 19: 568-574.
- Gordon NF, Scott CB, Levine BD. Comparison of single versus multiple lifestyle interventions: Are the antihypertensive effects of exercise training and diet-induced weight loss additive? *Am J Cardiol* 1997; 79: 763-767.
- Gowans SE, deHueck A, Voss S, Richardson M. A randomized, controlled trial of exercise and education for individuals with fibromyalgia. *Arthritis Care Res* 1999; 12: 120-128.
- Green RH, Singh SJ, Williams J, Morgan MD. A randomised controlled trial of four weeks versus seven weeks of pulmonary rehabilitation in chronic obstructive pulmonary disease. *Thorax* 2001; 56: 143-145.
- Greist JH, Klein MH, Eischens RR, Faris J, Gurman AS, Morgan WP. Running as treatment for depression. *Comp Psychiatry* 1979; 20: 41-54.
- Grieths TL, Burr ML, Campbell IA, Lewis-Jenkins V, Mullins J, Shiels K, Turner-Lawlor PJ, Payne N, Newcombe RG, Ionescu AA, Thomas J, Tunbridge J. Results at 1 year of outpatient multidisciplinary pulmonary rehabilitation: a randomised controlled trial. *Lancet* 2000; 355: 362-368.
- Grieths TL, Phillips CJ, Davies S, Burr ML, Campbell IA. Cost effectiveness of an outpatient multidisciplinary pulmonary rehabilitation programme. *Thorax* 2001; 56: 779-784.
- Grodstein F, Levine R, Troy L, Spencer T, Colditz GA, Stampfer MJ. Three-year follow-up of participants in a commercial weight loss program. Can you keep it off? *Arch Intern Med* 1996; 156: 1302-1306.
- Grosbois JM, Lamblin C, Lemaire B, Chekroud H, Dernis JM, Douay B, Fortin F. Long-term benefits of exercise maintenance after outpatient rehabilitation program in patients with chronic obstructive pulmonary disease. *J Cardiopulm Rehabil* 1999; 19: 216-225.
- Grove KA, Londeree BR. Bone density in postmenopausal women: high impact vs low impact exercise. *Med Sci Sports Exerc* 1992; 24: 1190-1194.
- Gueyffer F, Boutitie F, Boissel JP, Pocock S, Coope J, Cutler J, Ekbom T, Fagard R, Friedman L, Perry M, Prineas R, Schron E. Effect of antihypertensive drug treatment on cardiovascular outcomes in women and men. A meta-analysis of individual patient data from randomized, controlled trials. The INDANA Investigators. *Ann Intern Med* 1997; 126: 761-767.
- Gunnarsson R, Wallberg-Henriksson H, Rossner S, Wahren J. Serum lipid and lipoprotein levels in female type I diabetics: relationships to aerobic capacity and glycaemic control. *Diabetes Metab* 1987; 13: 417-421.
- Guo SS, Zeller C, Chumlea WC, Siervogel RM. Aging, body composition, and lifestyle: the Fels Longitudinal Study. *Am J Clin Nutr* 1999; 70: 405-411.
- Gustafsson T, Bodin K, Sylven C, Gordon A, Tyni-Lenne R, Jansson E. Increased expression of VEGF following exercise training in patients with heart failure. *Eur J Clin Invest* 2001; 31: 362-366.
- Guyatt GH, Kirshner B, Jaeschke R. A methodologic framework for health status measures: clarity or oversimplification? *J Clin Epidemiol* 1992; 45: 1353-1355.
- Guyatt GH, Pugsley SO, Sullivan MJ, Thompson PJ, Berman L, Jones NL, Fallen EL, Taylor DW. Effect of encouragement on walking test performance. *Thorax* 1984; 39: 818-822.
- Guyatt GH, Thompson PJ, Berman LB, Sullivan MJ, Townsend M, Jones NL, Pugsley SO. How should we measure function in patients with chronic heart and lung disease? *J Chronic Dis* 1985; 38: 517-524.
- Haapanen N, Miilunpalo S, Vuori I, Oja P, Pasanen M. Association of leisure time physical activity with the risk of coronary heart disease, hypertension and diabetes in middle-aged men and women. *Int J Epidemiol* 1997; 26: 739-747.
- Hagberg JM, Montain SJ, Martin WH III, Ehsani AA. Effect of exercise training in 60- to 69-year-old persons with essential hypertension. *Am J Cardiol* 1989; 64: 348-353.
- Hakkinen A, Hakkinen K, Hannonen P, Alen M. Strength training induced adaptations in neuromuscular function of premenopausal women with fibromyalgia: comparison with healthy women. *Ann Rheum Dis* 2001a; 60: 21-26.
- Hakkinen A, Hannonen P, Hakkinen K. Muscle strength in healthy people and in patients suffering from recent-onset inflammatory arthritis. *Br J Rheumatol* 1995; 34: 355-360.
- Hakkinen A, Sokka T, Kotaniemi A, Hannonen P. A randomized two-year study of the effects of dynamic strength training on muscle strength, disease activity, functional capacity, and bone mineral density in early rheumatoid arthritis. *Arthritis Rheum* 2001b; 44: 515-522.
- Hakkinen A, Sokka T, Kotaniemi A, Kautiainen H, Jappinen I, Laitinen L, Hannonen P. Dynamic strength training in patients with early rheumatoid arthritis increases muscle strength but not bone mineral density. *J Rheumatol* 1999; 26: 1257-1263.

- Hakkinen K, Pakarinen A, Hannonen P, Hakkinen A, Airaksinen O, Valkeinen H, Alen M. Effects of strength training on muscle strength, cross-sectional area, maximal electromyographic activity, and serum hormones in premenopausal women with fibromyalgia. *J Rheumatol* 2002; 29: 1287–1295.
- Halle M, Berg A, Garwers U, Baumstark MW, Knisel W, Grathwohl D, Konig D, Keul J. Influence of 4 weeks' intervention by exercise and diet on low-density lipoprotein subfractions in obese men with type 2 diabetes. *Metabolism* 1999; 48: 641–644.
- Hambrecht R, Gielen S, Linke A, Fiehn E, Yu J, Walther C, Schoene N, Schuler G. Effects of exercise training on left ventricular function and peripheral resistance in patients with chronic heart failure: a randomized trial. *JAMA* 2000; 283: 3095–3101.
- Hambrecht R, Niebauer J, Fiehn E, Kalberer B, O'ner B, Hauer K, Riede U, Schlierf G, Kubler W, Schuler G. Physical training in patients with stable chronic heart failure: effects on cardiorespiratory fitness and ultrastructural abnormalities of leg muscles. *J Am Coll Cardiol* 1995; 25: 1239–1249.
- Hamdorf PA, Penhall RK. Walking with its training effects on the fitness and activity patterns of 79–91 year old females. *Aust N Z J Med* 1999; 29: 22–28.
- Hamdorf PA, Withers RT, Penhall RK, Haslam MV. Physical training effects on the fitness and habitual activity patterns of elderly women. *Arch Phys Med Rehabil* 1992; 73: 603–608.
- Hansen TM, Hansen G, Langgaard AM, Rasmussen JO. Longterm physical training in rheumatoid arthritis. A randomized trial with different training programs and blinded observers. *Scand J Rheumatol* 1993; 22: 107–112.
- Harkcom TM, Lampman RM, Banwell BF, Castor CW. Therapeutic value of graded aerobic exercise training in rheumatoid arthritis. *Arthritis Rheum* 1985; 28: 32–39.
- Harrington D, Anker SD, Chua TP, Webb-Peploe KM, Ponikowski PP, Poole-Wilson PA, Coats AJ. Skeletal muscle function and its relation to exercise tolerance in chronic heart failure. *J Am Coll Cardiol* 1997; 30: 1758–1764.
- Hartman CA, Manos TM, Winter C, Hartman DM, Li B, Smith JC. Effects of Tai Chi training on function and quality of life indicators in older adults with osteoarthritis. *J Am Geriatr Soc* 2000; 48: 1553–1559.
- Hartman WM, Stroud M, Sweet DM, Saxton J. Long-term maintenance of weight loss following supplemented fasting. *Int J Eat Disord* 1993; 14: 87–93.
- Haskell WL, Alderman EL, Fair JM, Maron DJ, Mackey SF, Superko HR, Williams PT, Johnstone IM, Champagne MA, Krauss RM. Effects of intensive multiple risk factor reduction on coronary atherosclerosis and clinical cardiac events in men and women with coronary artery disease. The Stanford Coronary Risk Intervention Project (SCRIP). *Circulation* 1994; 89: 975–990.
- Hass CJ, Garzarella L, de Hoyos DV, Connaughton DP, Pollock ML. Concurrent improvements in cardiorespiratory and muscle fitness in response to total body recumbent stepping in humans. *Eur J Appl Physiol* 2001; 85: 157–163.
- Hatori M, Hasegawa A, Adachi H, Shinozaki A, Hayashi R, Okano H, Mizunuma H, Murata K. The effects of walking at the anaerobic threshold level on vertebral bone loss in postmenopausal women. *Calcif Tissue Int* 1993; 52: 411–414.
- Haus G, Hoerr SL, Mavis B, Robison J. Key modifiable factors in weight maintenance: fat intake, exercise, and weight cycling. *J Am Diet Assoc* 1994; 94: 409–413.
- Havlik RJ, Phillips CL, Brock DB, Lohman K, Haskell W, Snell P, O'Toole M, Ribisl P, Vaitkevicius P, Spurgeon HA, Lakatta EG, Pullen P. Walking may be related to less vascular stiffness in the Activity Counseling Trial (ACT). *Am Heart J* 2005; 150: 270–275.
- Hawkins P, Johnson LC, Nikolettou D, Hamnegard CH, Sherwood R, Polkey MI, Moxham J. Proportional assist ventilation as an aid to exercise training in severe chronic obstructive pulmonary disease. *Thorax* 2002; 57: 853–859.
- Heitmann BL, Kaprio J, Harris JR, Rissanen A, Korkeila M, Koskenvuo M. Are genetic determinants of weight gain modified by leisure-time physical activity? A prospective study of Finnish twins. *Am J Clin Nutr* 1997; 66: 672–678.
- Heldal M, Sire S, Dale J. Randomised training after myocardial infarction: short and long-term effects of exercise training after myocardial infarction in patients on beta-blocker treatment. A randomized, controlled study. *Scand Cardiovasc J* 2000; 34: 59–64.
- Helge EW. High prevalence of eating disorders among elite athletes. Increased risk of amenorrhea and premenopausal osteoporosis. *Ugeskr Laeger* 2001; 163: 3473–3475.
- Hellenius ML, de Faire U, Berglund B, Hamsten A, Krakau I. Diet and exercise are equally effective in reducing risk for cardiovascular disease. Results of a randomized controlled study in men with slightly to moderately raised cardiovascular risk factors. *Atherosclerosis* 1993; 103: 81–91.
- Heller RF, Knapp JC, Valenti LA, Dobson AJ. Secondary prevention after acute myocardial infarction. *Am J Cardiol* 1993; 72: 759–762.
- Hiatt WR, Regensteiner JG, Hargarten ME, Wolfel EE, Brass EP. Benefit of exercise conditioning for patients with peripheral arterial disease. *Circulation* 1990; 81: 602–609.
- Hiatt WR, Wolfel EE, Meier RH, Regensteiner JG. Superiority of treadmill walking exercise versus strength training for patients with peripheral arterial disease. Implications for the mechanism of the training response. *Circulation* 1994; 90: 1866–1874.
- Higashi Y, Sasaki S, Kurisu S, Yoshimizu A, Sasaki N, Matsuura H, Kajiyama G, Oshima T. Regular aerobic exercise augments endothelium-dependent vascular relaxation in normotensive as well as hypertensive subjects: role of endothelium-derived nitric oxide. *Circulation* 1999a; 100: 1194–1202.
- Higashi Y, Sasaki S, Sasaki N, Nakagawa K, Ueda T, Yoshimizu A, Kurisu S, Matsuura H, Kajiyama G, Oshima T.



- Daily aerobic exercise improves reactive hyperemia in patients with essential hypertension. *Hypertension* 1999b; 33: 591–597.
- Higgins HC, Hayes RL, McKenna KT. Rehabilitation outcomes following percutaneous coronary interventions (PCI). *Patient Educ Couns* 2001; 43: 219–230.
- Hinkleman LL, Nieman DC. The effects of a walking program on body composition and serum lipids and lipoproteins in overweight women. *J Sports Med Phys Fitness* 1993; 33: 49–58.
- Hochberg MC, Altman RD, Brandt KD, Clark BM, Dieppe PA, Gribbin MR, Moskowitz RW, Schnitzer TJ. Guidelines for the medical management of osteoarthritis. Part I. Osteoarthritis of the hip. *American College of Rheumatology. Arthritis Rheum* 1995a; 38: 1535–1540.
- Hochberg MC, Altman RD, Brandt KD, Clark BM, Dieppe PA, Gribbin MR, Moskowitz RW, Schnitzer TJ. Guidelines for the medical management of osteoarthritis. Part II. Osteoarthritis of the knee. *American College of Rheumatology. Arthritis Rheum* 1995b; 38: 1541–1546.
- Hoiberg A, Berard S, Watten RH, Caine C. Correlates of weight loss in treatment and at follow-up. *Int J Obes* 1984; 8: 457–465.
- Hoidrup S. Risk factors for hip fracture. Copenhagen: Kommunehospitalet, Institute of Preventive Medicine, 1997: 1–120.
- Holden JH, Darga LL, Olson SM, Stettner DC, Ardito EA, Lucas CP. Long-term follow-up of patients attending a combination very-low calorie diet and behaviour therapy weight loss programme. *Int J Obes Relat Metab Disord* 1992; 16: 605–613.
- Holm J, Dahllöf AG, Björntorp P, Schersten T. Enzyme studies in muscles of patients with intermittent claudication. Effect of training. *Scand J Clin Lab Invest Suppl* 1973; 128: 201–205.
- Holmback AM, Sawe U, Fagher B. Training after myocardial infarction: lack of long-term effects on physical capacity and psychological variables. *Arch Phys Med Rehabil* 1994; 75: 551–554.
- Holmes GP, Kaplan JE, Gantz NM, Komaroff AL, Schonberger LB, Straus SE, Jones JF, Dubois RE, Cunningham-Rundles C, Pahwa S. Chronic fatigue syndrome: a working case definition. *Ann Intern Med* 1988; 108: 387–389.
- Holmes MD, Chen WY, Feskanich D, Kroenke CH, Colditz GA. Physical activity and survival after breast cancer diagnosis. *JAMA* 2005; 293: 2479–2486.
- Holtén MK, Zacho M, Gaster M, Juel C, Wojtaszewski JF, Dela F. Strength training increases insulin-mediated glucose uptake, GLUT4 content, and insulin signaling in skeletal muscle in patients with type 2 diabetes. *Diabetes* 2004; 53: 294–305.
- Honkola A, Forsen T, Eriksson J. Resistance training improves the metabolic profile in individuals with type 2 diabetes. *Acta Diabetol* 1997; 34: 245–248.
- Hopman-Rock M, Westhög MH. The effects of a health educational and exercise program for older adults with osteoarthritis for the hip or knee. *J Rheumatol* 2000; 27: 1947–1954.
- Hornsby WG, Boggess KA, Lyons TJ, Barnwell WH, Lazarchick J, Colwell JA. Hemostatic alterations with exercise conditioning in NIDDM. *Diabetes Care* 1990; 13: 87–92.
- Houde SC, Melillo KD. Cardiovascular health and physical activity in older adults: an integrative review of research methodology and results. *J Adv Nurs* 2002; 38: 219–234.
- Houmar J, Tanner CJ, Slentz CA, Duscha BD, McCartney JS, Kraus WE. Effect of the volume and intensity of exercise training on insulin sensitivity. *J Appl Physiol* 2004; 96: 101–106.
- Hsieh LF, Didenko B, Schumacher HR Jr, Torg JS. Isokinetic and isometric testing of knee musculature in patients with rheumatoid arthritis with mild knee involvement. *Arch Phys Med Rehabil* 1987; 68: 294–297.
- Ivy JL, Zderic TW, Fogt DL. Prevention and treatment of non-insulin-dependent diabetes mellitus. *Exerc Sport Sci Rev* 1999; 27(1–35): 1–35.
- Iwamoto J, Takeda T, Ichimura S. Effect of exercise training and detraining on bone mineral density in postmenopausal women with osteoporosis. *J Orthop Sci* 2001; 6: 128–132.
- Jaeschke R, Singer J, Guyatt GH. Measurement of health status. Ascertaining the minimal clinically important difference. *Control Clin Trials* 1989; 10: 407–415.
- Jan MH, Lai JS. The effects of physiotherapy on osteoarthritic knees of females. *J Formos Med Assoc* 1991; 90: 1008–1013.
- Jennings G, Nelson L, Nestel P, Esler M, Korner P, Burton D, Bazelmans J. The effects of changes in physical activity on major cardiovascular risk factors, hemodynamics, sympathetic function, and glucose utilization in man: a controlled study of four levels of activity. *Circulation* 1986; 73: 30–40.
- Jessup JV, Lowenthal DT, Pollock ML, Turner T. The effects of endurance exercise training on ambulatory blood pressure in normotensive older adults. *Geriatr Nephrol Urol* 1998; 8: 103–109.
- Jette M, Heller R, Landry F, Blumchen G. Randomized 4-week exercise program in patients with impaired left ventricular function. *Circulation* 1991; 84: 1561–1567.
- Jewitt DE, Reid D, Thomas M, Mercer CJ, Valori C, Shillingford JP. Free noradrenaline and adrenaline excretion in relation to the development of cardiac arrhythmias and heart failure in patients with acute myocardial infarction. *Lancet* 1969; 1: 635–641.
- Johnson PH, Cowley AJ, Kinnear WJ. A randomized controlled trial of inspiratory muscle training in stable chronic heart failure. *Eur Heart J* 1998; 19: 1249–1253.
- Johnstone MT, Creager SJ, Scales KM, Cusco JA, Lee BK, Creager MA. Impaired endothelium-dependent vasodilation in patients with insulin-dependent diabetes mellitus. *Circulation* 1993; 88: 2510–2516.
- Jolliffe JA, Rees K, Taylor RS, Thompson D, Oldridge N, Ebrahim S. Exercise-based rehabilitation for coronary heart disease. *Cochrane Database Syst Rev* 2000 4-CD001800.

- Jonason T, Ringqvist I. Prediction of the effect of training on the walking tolerance in patients with intermittent claudication. *Scand J Rehabil Med* 1987; 19: 47–50.
- Jones DR., Speier J, Canine K, Owen R, Stull GA. Cardiorespiratory responses to aerobic training by patients with postpoliomyelitis sequelae. *JAMA* 1989; 261: 3255–3258.
- Jones DT, Thomson RJ, Sears MR. Physical exercise and resistive breathing training in severe chronic airways obstruction – are they effective? *Eur J Respir Dis* 1985; 67: 159–166.
- Jones KD, Burckhardt CS, Clark SR, Bennett RM, Potempa KM. A randomized controlled trial of muscle strengthening versus flexibility training in fibromyalgia. *J Rheumatol* 2002; 29: 1041–1048.
- Jones NL. In *Clinical exercise testing*. Philadelphia, PA, USA: W.B. Saunders, 1988: 123–134.
- Joslin EP, Root EF, White P. The treatment of diabetes mellitus. Philadelphia: Lea & Febiger, 1959. Jurca R, Lamonte MJ, Church TS, Earnest CP, Fitzgerald SJ, Barlow CE, Jordan AN, Kampert JB, Blair SN. Associations of muscle strength and fitness with metabolic syndrome in men. *Med Sci Sports Exerc* 2004; 36: 1301–1307.
- Kallio V, Hamalainen H, Hakkila J, Luurila OJ. Reduction in sudden deaths by a multifactorial intervention programme after acute myocardial infarction. *Lancet* 1979; 2: 1091–1094.
- Kanis JA. The incidence of hip fracture in Europe. *Osteoporos Int* 1993; 3(Suppl. 1:10–5): 10–15.
- Kannel WB, McGee DL. Diabetes and cardiovascular disease. The Framingham study. *JAMA* 1979; 241: 2035–2038.
- Kaplan RM, Hartwell SL, Wilson DK, Wallace JP. Effects of diet and exercise interventions on control and quality of life in non-insulin-dependent diabetes mellitus. *J Gen Intern Med* 1987; 2: 220–228.
- Kavanagh T, Myers MG, Baigrie RS, Mertens DJ, Sawyer P, Shephard RJ. Quality of life and cardiorespiratory function in chronic heart failure: effects of 12 months' aerobic training. *Heart* 1996; 76: 42–49.
- Kayman S, Bruvold W, Stern JS. Maintenance and relapse after weight loss in women: behavioral aspects. *Am J Clin Nutr* 1990; 52: 800–807.
- Keats MR, Courneya KS, Danielsen S, Whitsett SF. Leisure-time physical activity and psychosocial well-being in adolescents after cancer diagnosis. *J Pediatr Oncol Nurs* 1999; 16: 180–188.
- Keefe FJ, Caldwell DS, Baucom D, Salley A, Robinson E, Timmons K, Beaupre P, Weisberg J, Helms M. Spouse-assisted coping skills training in the management of knee pain in osteoarthritis: long-term followup results. *Arthritis Care Res* 1999; 12: 101–111.
- Kellemen MH, Eron MB, Valenti SA, Stewart KJ. Exercise training combined with antihypertensive drug therapy. Effects on lipids, blood pressure, and left ventricular mass. *JAMA* 1990; 263: 2766–2771.
- Kelley G. Aerobic exercise and lumbar spine bone mineral density in postmenopausal women: a meta-analysis. *J Am Geriatr Soc* 1998a; 46: 143–152.
- Kelley GA. Aerobic exercise and bone density at the hip in postmenopausal women: a meta-analysis. *Prev Med* 1998b; 27: 798–807.
- Kelley GA. Exercise and regional bone mineral density in postmenopausal women: a meta-analytic review of randomized trials. *Am J Phys Med Rehabil* 1998c; 77: 76–87.
- Kelley GA. Aerobic exercise and resting blood pressure among women: a meta-analysis. *Prev Med* 1999; 28: 264–275.
- Kelley GA, Kelley KS. Aerobic exercise and resting blood pressure in women: a meta-analytic review of controlled clinical trials. *J Womens Health Gend Based Med* 1999; 8: 787–803.
- Kelley GA, Kelley KS. Progressive resistance exercise and resting blood pressure: a meta-analysis of randomized controlled trials. *Hypertension* 2000; 35: 838–843.
- Kelley GA, Kelley KS. Efficacy of resistance exercise on lumbar spine and femoral neck bone mineral density in premenopausal women: a meta-analysis of individual patient data. *J Womens Health (Larchmt)* 2004; 13: 293–300.
- Kelley GA, Sharpe KK. Aerobic exercise and resting blood pressure in older adults: a meta-analytic review of randomized controlled trials. *J Gerontol A Biol Sci Med Sci* 2001; 56: M298–M303.
- Kelley GA, Kelley KA, Tran ZV. Aerobic exercise and resting blood pressure: a meta-analytic review of randomized, controlled trials. *Prev Cardiol* 2001a; 4: 73–80.
- Kelley GA, Kelley KS, Tran ZV. Resistance training and bone mineral density in women: a meta-analysis of controlled trials. *Am J Phys Med Rehabil* 2001b; 80: 65–77.
- Kelley GA, Kelley KS, Tran ZV. Walking and resting blood pressure in adults: a meta-analysis. *Prev Med* 2001c; 33: 120–127.
- Kemper HC, Twisk JW, van Mechelen W, Post GB, Roos JC, Lips P. A fifteen-year longitudinal study in young adults on the relation of physical activity and fitness with the development of the bone mass: the Amsterdam Growth and Health Longitudinal Study. *Bone* 2000; 27: 847–853.
- Kentala E. Physical fitness and feasibility of physical rehabilitation after myocardial infarction in men of working age. *Ann Clin Res Suppl* 1972; 9: 1–84.
- Kerr D, Morton A, Dick I, Prince R. Exercise effects on bone mass in postmenopausal women are site-specific and load-dependent. *J Bone Miner Res* 1996; 11: 218–225.
- Ketelhut RG, Franz IW, Scholze J. Regular exercise as an effective approach in antihypertensive therapy. *Med Sci Sports Exerc* 2004; 36: 4–8.
- Keteyian SJ, Levine AB, Brawner CA, Kataoka T, Rogers FJ, Schairer JR, Stein PD, Levine TB, Goldstein S. Exercise training in patients with heart failure. A randomized, controlled trial. *Ann Intern Med* 1996; 124: 1051–1057.
- Kiesewetter H, Blume J, Jung F, Gerhards M, Leipnitz G. Training by walking and drug therapy of peripheral arterial occlusive disease. *Vasa Suppl* 1987; 20: 384–387.



- Kiilavuori K, Sovijarvi A, Naveri H, Ikonen T, Leinonen H. Effect of physical training on exercise capacity and gas exchange in patients with chronic heart failure. *Chest* 1996; 110: 985–991.
- Kiilavuori K, Toivonen L, Naveri H, Leinonen H. Reversal of autonomic derangements by physical training in chronic heart failure assessed by heart rate variability. *Eur Heart J* 1995; 16: 490–495.
- King AC, Frey-Hewitt B, Dreon DM, Wood PD. Diet vs exercise in weight maintenance. The effects of minimal intervention strategies on long-term outcomes in men. *Arch Intern Med* 1989; 149: 2741–2746.
- King AC, Haskell WL, Taylor CB, Kraemer HC, DeBusk RF. Group- vs home-based exercise training in healthy older men and women. A community-based clinical trial. *JAMA* 1991; 266: 1535–1542.
- King SJ, Wessel J, Bhambhani Y, Sholter D, Maksymowych W. The effects of exercise and education, individually or combined, in women with fibromyalgia. *J Rheumatol* 2002; 29: 2620–2627.
- Kingwell BA, Jennings GL. Effects of walking and other exercise programs upon blood pressure in normal subjects. *Med J Aust* 1993; 158: 234–238.
- Kirk A, Mutrie N, MacIntyre P, Fisher M. Increasing physical activity in people with type 2 diabetes. *Diabetes Care* 2003; 26: 1186–1192.
- Klein MH, Greist JH, Gurman RA, Neimeyer RA, Lesser DP, Bushnell NJ, et al. A comparative outcome study of group psychotherapy vs. exercise treatments for depression. *Int J Ment Health* 1985; 13: 148–177.
- Knopp RH. Drug treatment of lipid disorders. *N Engl J Med* 1999; 341: 498–511.
- Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 2002; 346: 393–403.
- Koch M, Douard H, Broustet JP. The benefit of graded physical exercise in chronic heart failure. *Chest* 1992; 101: 231S–235S.
- Kohl HW, Gordon NF, Villegas JA, Blair SN. Cardiorespiratory fitness, glycemic status, and mortality risk in men. *Diabetes Care* 1992; 15: 184–192.
- Kohrt WM, Ehsani AA, Birge SJ, Jr. Effects of exercise involving predominantly either joint-reaction or ground-reaction forces on bone mineral density in older women. *J Bone Miner Res* 1997; 12: 1253–1261.
- Koivisto UM, Martinez-Valdez H, Bilan PJ, Burdett E, Ramlal T, Klip A. Differential regulation of the GLUT-1 and GLUT-4 glucose transport systems by glucose and insulin in L6 muscle cells in culture. *J Biol Chem* 1991; 266: 2615–2621.
- Koivisto VA, Felig P. Effects of leg exercise on insulin absorption in diabetic patients. *N Engl J Med* 1978; 298: 79–83.
- Kokkinos PF, Narayan P, Collieran JA, Pittaras A, Notargiacomo A, Reda D, Papademetriou V. Effects of regular exercise on blood pressure and left ventricular hypertrophy in African-American men with severe hypertension. *New Engl J Med* 1995; 333: 1462–1467.
- Komatireddy GR, Leitch RW, Cella K, Browning G, Minor M. Efficacy of low load resistive muscle training in patients with rheumatoid arthritis functional class II and III. *J Rheumatol* 1997; 24: 1531–1539.
- Kongsgaard M, Backer V, Jorgensen K, Kjaer M, Beyer N. Heavy resistance training increases muscle size, strength and physical function in elderly male COPD-patients – a pilot study. *Respir Med* 2004; 98: 1000–1007.
- Koukouvou G, Kouidi E, Iacovides A, Konstantinidou E, Kaprinis G, Deligiannis A. Quality of life, psychological and physiological changes following exercise training in patients with chronic heart failure. *J Rehabil Med* 2004; 36: 36–41.
- Kovar PA, Allegrante JP, MacKenzie CR, Peterson MG, Gutin B, Charlson ME. Supervised fitness walking in patients with osteoarthritis of the knee. A randomized, controlled trial. *Ann Intern Med* 1992; 116: 529–534.
- Krachler M, Lindschinger M, Eber B, Watzinger N, Wallner S. Trace elements in coronary heart disease: impact of intensified lifestyle modification. *Biol Trace Elem Res* 1997; 60: 175–185.
- Kraemer WJ, Keuning M, Ratamess NA, Volek JS, McCormick M, Bush JA, Nindl BC, Gordon SE, Mazzetti SA, Newton RU, Gomez AL, Wickham RB, Rubin MR, Hakkinen K. Resistance training combined with bench-step aerobics enhances women's health profile. *Med Sci Sports Exerc* 2001; 33: 259–269.
- Kraemer WJ, Volek JS, Clark KL, Gordon SE, Incledon T, Puhl SM, Triplett-McBride NT, McBride JM, Putukian M, Sebastianelli WJ. Physiological adaptations to a weight-loss dietary regimen and exercise programs in women. *J Appl Physiol* 1997; 83: 270–279.
- Kraemer WJ, Volek JS, Clark KL, Gordon SE, Puhl SM, Koziris LP, McBride JM, Triplett-McBride NT, Putukian M, Newton RU, Hakkinen K, Bush JA, Sebastianelli WJ. Influence of exercise training on physiological and performance changes with weight loss in men. *Med Sci Sports Exerc* 1999; 31: 1320–1329.
- Kraus WE, Houshard JA, Duscha BD, Knetzger KJ, Wharton MB, McCartney JS, Bales CW, Henes S, Samsa GP, Otvos JD, Kulkarni KR, Slentz CA. Effects of the amount and intensity of exercise on plasma lipoproteins. *N Engl J Med* 2002; 347: 1483–1492.
- Krolewski AS. Magnitude and determinants of coronary artery disease in juvenile-onset, insulin-dependent diabetes mellitus. *Am J Cardiol* 1987; 59: 750–755.
- Krolner B, Toft B. Vertebral bone loss: an unheeded side effect of therapeutic bed rest. *Clin Sci (Lond)* 1983; 64: 537–540.
- Krotkiewski M, Lonroth P, Mandroukas K, Wroblewski Z, Rebuffe-Scrive M, Holm G, Smith U, Bjorntorp P. The effects of physical training on insulin secretion and effectiveness and on glucose metabolism in obesity and type 2 (non-insulin-dependent) diabetes mellitus. *Diabetologia* 1985; 28: 881–890.
- Krummel DA, Koeman DM, Bronner Y, Davis J, Greenlund K, Tessaro I, Upson D, Wilbur J. Cardiovascular health interventions in women: what works? *J Womens Health Gend Based Med* 2001; 10: 117–136.
- Kukkonen K, Rauramaa R, Voutilainen E, Lansimies E. Physical training of middle-aged men with borderline hypertension. *Ann Clin Res* 1982; 14(Suppl. 34):139–145.



- Laaksonen DE, Atalay M, Niskanen LK, Mustonen J, Sen CK, Lakka TA, Uusitupa MI. Aerobic exercise and the lipid profile in type 1 diabetic men: a randomized controlled trial. *Med Sci Sports Exerc* 2000; 32: 1541–1548.
- Lacasse Y, Brosseau L, Milne S, Martin S, Wong E, Guyatt GH, Goldstein RS. Pulmonary rehabilitation for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* 2002 3- CD003793.
- Lacasse Y, Wong E, Guyatt GH, King D, Cook DJ, Goldstein RS. Meta-analysis of respiratory rehabilitation in chronic obstructive pulmonary disease. *Lancet* 1996; 348: 1115–1119.
- Lake FR, Henderson K, Briant T, Openshaw J, Musk AW. Upper-limb and lower-limb exercise training in patients with chronic airflow obstruction. *Chest* 1990; 97: 1077–1082.
- Langbein WE, Collins EG, Orebaugh C, Maloney C, Williams KJ, Littooy FN, Edwards LC. Increasing exercise tolerance of persons limited by claudication pain using polestriding. *J Vasc Surg* 2002; 35: 887–893.
- Lange P, Vestbo J. (2 A.D.). Obstruktive lungesygdomme. In *Medicinsk kompendium*.
- Lansimies E, Hietanen E, Huttunen JK, Hanninen O, Kukkonen K, Rauramaa R, Voutilainen E. Metabolic and hemodynamic effects of physical training in middle-aged men – a controlled trial. In: Komi PV, Nelson RC, Morehouse CA, eds. *Exercise and sport biology*. Champaign, IL: Human Kinetics, 1979: 199–206.
- Larsen OA, Lassen NA. Effect of daily muscular exercise in patients with intermittent claudication. *Lancet* 1966; 2: 1093–1096.
- Lau EM, Woo J, Leung PC, Swaminathan R, Leung D. The effects of calcium supplementation and exercise on bone density in elderly Chinese women. *Osteoporos Int* 1992; 2: 168–173.
- Law MR, Wald NJ, Meade TW. Strategies for prevention of osteoporosis and hip fracture. *BMJ* 1991; 303: 453–459.
- Lawlor DA, Hopker SW. The effectiveness of exercise as an intervention in the management of depression: systematic review and meta-regression analysis of randomised controlled trials. *BMJ* 2001; 322: 763–767.
- Lee AP, Ice R, Blessey R, Sanmarco ME. Long-term effects of physical training on coronary patients with impaired ventricular function. *Circulation* 1979; 60: 1519–1526.
- Lee S, Kuk JL, Davidson LE, Hudson R, Kilpatrick K, Graham TE, Ross R. Exercise without weight loss is an effective strategy for obesity reduction in obese individuals with and without Type 2 diabetes. *J Appl Physiol* 2005; 99: 1220–1225.
- Leermakers EA, Perri MG, Shigaki CL, Fuller PR. Effects of exercise-focused versus weight-focused maintenance programs on the management of obesity. *Addict Behav* 1999; 24: 219–227.
- Lehmann R, Engler H, Honegger R, Riesen W, Spinass GA. Alterations of lipolytic enzymes and high-density lipoprotein subfractions induced by physical activity in type 2 diabetes mellitus. *Eur J Clin Invest* 2001; 31: 37–44.
- Lehmann R, Kaplan V, Bingisser R, Bloch KE, Spinass GA. Impact of physical activity on cardiovascular risk factors in IDDM. *Diabetes Care* 1997; 20: 1603–1611.
- Lehmann R, Vokac A, Niedermann K, Agosti K, Spinass GA. Loss of abdominal fat and improvement of the cardiovascular risk profile by regular moderate exercise training in patients with NIDDM. *Diabetologia* 1995; 38: 1313–1319.
- LeMaitre JP, Harris S, Fox KA, Denvir M. Change in circulating cytokines after 2 forms of exercise training in chronic stable heart failure. *Am Heart J* 2004; 147: 100–105.
- LeMura LM, Maziekas MT. Factors that alter body fat, body mass, and fat-free mass in pediatric obesity. *Med Sci Sports Exerc* 2002; 34: 487–496.
- Leng GC, Fowler B, Ernst E. Exercise for intermittent claudication. *Cochrane Database Syst Rev* 2000 CD000990.
- Leon AS. Effects of exercise conditioning on physiologic precursors of CHD. *J Cardiopulm Rehabil* 1991; 11: 46–57.
- Leon AS. Exercise in the prevention and management of diabetes mellitus and blood lipid disorders. In: Shephard RJ, Miller HSJ, eds. *Exercise and the heart in health and disease*. New York: Marcel Dekker, 1999: 355–420.
- Leon AS, Sanchez OA. Response of blood lipids to exercise training alone or combined with dietary intervention. *Med Sci Sports Exerc* 2001; 33: S502–S515.
- Leon AS, Casal D, Jacobs D Jr. Effects of 2, 000 kcal per week of walking and stair climbing on physical fitness and risk factors for coronary heart disease. *J Cardiopulm Rehabil* 1996; 16: 183–192.
- Letac B, Cribier A, Desplanches JF. A study of left ventricular function in coronary patients before and after physical training. *Circulation* 1977; 56: 375–378.
- Levy WC, Cerqueira MD, Abrass IB, Schwartz RS, Stratton JR. Endurance exercise training augments diastolic filling at rest and during exercise in healthy young and older men. *Circulation* 1993; 88: 116–126.
- Lewin B, Robertson IH, Cay EL, Irving JB, Campbell M. Effects of self-help post-myocardial-infarction rehabilitation on psychological adjustment and use of health services. *Lancet* 1992; 339: 1036–1040.
- Lewington S, Clarke R, Qizilbash N, Peto R, Collins R. Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet* 2002; 360: 1903–1913.
- Li YP, Reid MB. Effect of tumor necrosis factor-alpha on skeletal muscle metabolism. *Curr Opin Rheumatol* 2001; 13: 483–487.
- Lindgarde F, Eriksson KF, Lithell H, Saltin B. Coupling between dietary changes, reduced body weight, muscle fibre size and improved glucose tolerance in middle-aged men with impaired glucose tolerance. *Acta Med Scand* 1982; 212: 99–106.
- Lindheim SR, Notelovitz M, Feldman EB, Larsen S, Khan FY, Lobo RA. The independent effects of exercise and estrogen on lipids and lipoproteins in postmenopausal women. *Obstet Gynecol* 1994; 83: 167–172.
- Lindstrom J, Eriksson JG, Valle TT, Aunola S, Cepaitis Z, Hakumaki M, Hamalainen H, Ilanne-Parikka P, Keinanen-Kiukkaanniemi S, Laakso M, Louheranta A, Mannelin M, Martikkala V, Moltchanov V, Rastas M, Salminen V, Sundvall J, Uusitupa M, Tuomilehto J. Prevention of diabetes mellitus in subjects with impaired glucose tolerance in the Finnish diabetes prevention study: results from a randomized clinical trial. *J Am Soc Nephrol* 2003a; 14:



S108– S113.

- Lindstrom J, Louheranta A, Mannelin M, Rastas M, Salminen V, Eriksson J, Uusitupa M, Tuomilehto J. The Finnish Diabetes Prevention Study (DPS): lifestyle intervention and 3-year results on diet and physical activity. *Diabetes Care* 2003b; 26: 3230–3236.
- Lisspers J, Sundin O, Hofman-Bang C, Nordlander R, Nygren A, Ryden L, Ohman A. Behavioral effects of a comprehensive, multifactorial program for lifestyle change after percutaneous transluminal coronary angioplasty: a prospective, randomized controlled study. *J Psychosom Res* 1999; 46: 143– 154.
- Lloyd-Williams F, Mair FS, Leitner M. Exercise training and heart failure: a systematic review of current evidence. *Br J Gen Pract* 2002; 52: 47–55.
- Loge JH, Abrahamsen AF, Ekeberg O, Kaasa S. Hodgkin's disease survivors more fatigued than the general population. *J Clin Oncol* 1999; 17: 253– 261.
- Lokey EA, Tran ZV. Effects of exercise training on serum lipid and lipoprotein concentrations in women: a meta-analysis. *Int J Sports Med* 1989; 10: 424–429.
- Lord SR, Ward JA, Williams P, Zivanovic E. The effects of a community exercise program on fracture risk factors in older women. *Osteoporos Int* 1996; 6: 361–367.
- Lundgren F, Dahllof AG, Lundholm K, Schersten T, Volkmann R. Intermittent claudication – surgical reconstruction or physical training? A prospective randomized trial of treatment efficacy. *Ann Surg* 1989a; 209: 346– 355.
- Lundgren F, Dahllof AG, Schersten T, Bylund-Fellenius AC. Muscle enzyme adaptation in patients with peripheral arterial insufficiency: spontaneous adaptation, effect of different treatments and consequences on walking performance. *Clin Sci (Lond)* 1989b; 77: 485–493.
- Lundgren S, Stenstrom CH. Muscle relaxation training and quality of life in rheumatoid arthritis. A randomized controlled clinical trial. *Scand J Rheumatol* 1999; 28: 47–53.
- Lyngberg KK, Harreby M, Bentzen H, Frost B, Danneskiold-Samsøe B. Elderly rheumatoid arthritis patients on steroid treatment tolerate physical training without an increase in disease activity. *Arch Phys Med Rehabil* 1994; 75: 1189–1195.
- MacVicar MG, Winningham ML. Promoting the functional capacity of cancer patients. *Cancer Bulletin* 1986; 38: 235–239.
- Magnusson G, Kaijser L, Sylven C, Karlberg KE, Isberg B, Saltin B. Peak skeletal muscle perfusion is maintained in patients with chronic heart failure when only a small muscle mass is exercised. *Cardiovasc Res* 1997; 33: 297–306.
- Maiorana A, O'Driscoll G, Cheetham C, Collis J, Goodman C, Rankin S, Taylor R, Green D. Combined aerobic and resistance exercise training improves functional capacity and strength in CHF. *J Appl Physiol* 2000; 88: 1565–1570.
- Makimattila S, Mantysaari M, Groop PH, Summanen P, Virkamaki A, Schlenzka A, Fagerudd J, Yki-Jarvinen H. Hyperreactivity to nitrovasodilators in forearm vasculature is related to autonomic dysfunction in insulin-dependent diabetes mellitus. *Circulation* 1997; 95: 618–625.
- Makimattila S, Virkamaki A, Groop PH, Cockcroft J, Utriainen T, Fagerudd J, Yki-Jarvinen H. Chronic hyperglycemia impairs endothelial function and insulin sensitivity via different mechanisms in insulin-dependent diabetes mellitus. *Circulation* 1996; 94: 1276–1282.
- Malkia E, Impivaara O. Intensity of physical activity and respiratory function in subjects with and without bronchial asthma. *Scand J Med Sci Sports* 1998; 8: 27–32.
- Manchanda SC, Narang R, Reddy KS, Sachdeva U, Prabhakaran D, Dharmanand S, Rajani M, Bijlani R. Retardation of coronary atherosclerosis with yoga lifestyle intervention. *J Assoc Physicians India* 2000; 48: 687–694.
- Mandroukas K, Krotkiewski M, Hedberg M, Wroblewski Z, Bjorntorp P, Grimby G. Physical training in obese women. Effects of muscle morphology, biochemistry and function. *Eur J Appl Physiol Occup Physiol* 1984; 52: 355– 361.
- Mann GV, Garrett HL, Farhi A, Murray H, Billings FT. Exercise to prevent coronary heart disease. An experimental study of the effects of training on risk factors for coronary disease in men. *Am J Med* 1969; 46: 12– 27.
- Mannarino E, Pasqualini L, Innocente S, Scricciolo V, Rignanese A, Ciufo G. Physical training and antiplatelet treatment in stage II peripheral arterial occlusive disease: alone or combined? *Angiology* 1991; 42: 513–521.
- Mannerkorpi K, Ahlmen M, Ekdahl C. Six- and 24-month follow-up of pool exercise therapy and education for patients with fibromyalgia. *Scand J Rheumatol* 2002; 31: 306–310.
- Marceau M, Kouame N, Lacourciere Y, Cleroux J. Effects of different training intensities on 24-hour blood pressure in hypertensive subjects. *Circulation* 1993; 88: 2803–2811.
- Marchionni N, Fattoroli F, Fumagalli S, Oldridge N, Del Lungo F, Morosi L, Burgisser C, Masotti G. Improved exercise tolerance and quality of life with cardiac rehabilitation of older patients after myocardial infarction: results of a randomized, controlled trial. *Circulation* 2003; 107: 2201–2206.
- Marcus BH, Simkin LR. The transtheoretical model: applications to exercise behavior. *Med Sci Sports Exerc* 1994; 26: 1400–1404.
- Marshall P, Al Timman J, Riley R, Wright J, Williams S, Hainsworth R, Tan LB. Randomized controlled trial of home-based exercise training to evaluate cardiac functional gains. *Clin Sci (Lond)* 2001; 101: 477–483.
- Martin D, Nodelovitz M. Effects of aerobic training on bone mineral density of postmenopausal women. *J Bone Miner Res* 1993; 8: 931–936.
- Martin JE, Dubbert PM, Cushman WC. Controlled trial of aerobic exercise in hypertension. *Circulation* 1990; 81: 1560–1567.
- Martin L, Nutting A, MacIntosh BR, Edworthy SM, Butterwick D, Cook J. An exercise program in the treatment of fibromyalgia. *J Rheumatol* 1996; 23: 1050–1053.
- Martinsen EW. Comparing aerobic and non-aerobic forms of exercise in the treatment of clinical depression: a randomized trial [abstract]. London: Sports Council and Health Education Authority, 1988: 84–95.
- Martinsen EW, Hoart A, Solberg O. Comparing aerobic with nonaerobic forms of exercise in the treatment of clinical

- depression: a randomized trial. *Compr Psychiatry* 1989; 30: 324–331.
- Martinsen EW, Medhus A, Sandvik L. Effects of aerobic exercise on depression: a controlled study. *Br Med J (Clin Res Ed)* 1985; 291: 109.
- Massie BM, Conway M, Rajagopalan B, Yonge R, Frostick S, Ledingham J, Sleight P, Radda G. Skeletal muscle metabolism during exercise under ischemic conditions in congestive heart failure. Evidence for abnormalities unrelated to blood flow. *Circulation* 1988; 78: 320–326.
- Mayoux-Benhamou MA, Bagheri F, Roux C, Auleley GR, Rabourdin JP, Revel M. Effect of psoas training on postmenopausal lumbar bone loss: a 3-year follow-up study. *Calcif Tissue Int* 1997; 60: 348–353.
- McAllister RM, Hirai T, Musch TI. Contribution of endothelium-derived nitric oxide (EDNO) to the skeletal muscle blood flow response to exercise.
- Med Sci Sports Exerc 1995; 27: 1145–1151.
- McCain GA, Bell DA, Mai FM, Halliday PD. A controlled study of the effects of a supervised cardiovascular fitness training program on the manifestations of primary fibromyalgia. *Arthritis Rheum* 1988; 31: 1135–1141.
- McCann IL, Holmes DS. Influence of aerobic exercise on depression. *J Pers Soc Psychol* 1984; 46: 1142–1147.
- McDonald CD, Burch GE, Walsh JJ. Prolonged bed rest in the treatment of idiopathic cardiomyopathy. *Am J Med* 1972; 52: 41–50.
- McDowell SL, Chaloa K, Housh TJ, Sharp GD, Johnson GO. The effect of exercise intensity and duration on salivary immunoglobulin A. *Eur J Appl Physiol* 1991; 63: 108–111.
- McGavin CR, Gupta SP, Lloyd EL, McHardy GJ. Physical rehabilitation for the chronic bronchitic: results of a controlled trial of exercises in the home. *Thorax* 1977; 32: 307–311.
- McGuire MT, Wing RR, Klem ML, Hill JO. Behavioral strategies of individuals who have maintained long-term weight losses. *Obes Res* 1999; 7: 334–341.
- McKay HA, Petit MA, Schutz RW, Prior JC, Barr SI, Khan KM. Augmented trochanteric bone mineral density after modified physical education classes: a randomized school-based exercise intervention study in prepubescent and early pubescent children. *J Pediatr* 2000; 136: 156–162.
- McMeeken J, Stillman B, Story I, Kent P, Smith J. The effects of knee extensor and flexor muscle training on the timed-up-and-go test in individuals with rheumatoid arthritis. *Physiother Res Int* 1999; 4: 55–67.
- McNally PG, Watt PA, Rimmer T, Burden AC, Hearnshaw JR, Thurston H. Impaired contraction and endothelium-dependent relaxation in isolated resistance vessels from patients with insulin-dependent diabetes mellitus. *Clin Sci (Lond)* 1994; 87: 31–36.
- McNeil JK, LeBlanc EM, Joyner M. The effect of exercise on depressive symptoms in the moderately depressed elderly. *Psychol Aging* 1991; 6: 487–488.
- McVeigh GE, Brennan GM, Johnston GD, McDermott BJ, McGrath LT, Henry WR, Andrews JW, Hayes JR. Impaired endothelium-dependent and independent vasodilation in patients with type 2 (non-insulin-dependent) diabetes mellitus. *Diabetologia* 1992; 35: 771–776.
- Mengshoel AM, Komnaes HB, Forre O. The effects of 20 weeks of physical fitness training in female patients with fibromyalgia. *Clin Exp Rheumatol* 1992; 10: 345–349.
- Meredith IT, Friberg P, Jennings GL, Dewar EM, Fazio VA, Lambert GW, Esler MD. Exercise training lowers resting renal but not cardiac sympathetic activity in humans. *Hypertension* 1991; 18: 575–582.
- Meredith IT, Jennings GL, Esler MD, Dewar EM, Bruce AM, Fazio VA, Korner PI. Time-course of the antihypertensive and autonomic effects of regular endurance exercise in human subjects. *J Hypertens* 1990; 8: 859–866.
- Messier SP, Loeser RF, Mitchell MN, Valle G, Morgan TP, Rejeski WJ, Ettinger WH. Exercise and weight loss in obese older adults with knee osteoarthritis: a preliminary study. *J Am Geriatr Soc* 2000; 48: 1062–1072.
- Messier SP, Thompson CD, Ettinger WH. Effects of long-term aerobic or weight training regimens on gait in an older, osteoarthritic population. *J Appl Biomech* 1997; 13: 205–225.
- Meyer K, Schwaibold M, Westbrook S, Beneke R, Hajric R, Gornandt L, Lehmann M, Roskamm H. Effects of short-term exercise training and activity restriction on functional capacity in patients with severe chronic congestive heart failure. *Am J Cardiol* 1996; 78: 1017–1022.
- Miedema H. *Reuma-Onderzoek Meerdere Echelons (ROME): Basisrapport*. Leiden, The Netherlands: NIPG-TNO, 1994.
- Miller NH, Haskell WL, Berra K, DeBusk RF. Home versus group exercise training for increasing functional capacity after myocardial infarction. *Circulation* 1984; 70: 645–649.
- Miller WC, Koceja DM, Hamilton EJ. A meta-analysis of the past 25 years of weight loss research using diet, exercise or diet plus exercise intervention. *Int J Obes Relat Metab Disord* 1997; 21: 941–947.
- Minor MA, Hewett JE, Webel RR, Anderson SK, Kay DR. Efficacy of physical conditioning exercise in patients with rheumatoid arthritis and osteoarthritis. *Arthritis Rheum* 1989; 32: 1396–1405.
- Minor MA, Hewett JE, Webel RR, Dreisinger TE, Kay DR. Exercise tolerance and disease related measures in patients with rheumatoid arthritis and osteoarthritis. *J Rheumatol* 1988; 15: 905–911.
- Minotti JR, Massie BM. Exercise training in heart failure patients. Does reversing the peripheral abnormalities protect the heart? *Circulation* 1992; 85: 2323–2325.
- Miyai N, Arita M, Miyashita K, Morioka I, Shiraishi T, Nishio I, Takeda S. Antihypertensive effects of aerobic exercise in middle-aged normotensive men with exaggerated blood pressure response to exercise. *Hypertens Res* 2002; 25: 507–514.
- Mock V, Burke MB, Sheehan P, Creaton EM, Winningham ML, McKenney-Tedder S, Schwager LP, Liebman M. A nursing rehabilitation program for women with breast cancer receiving adjuvant chemotherapy. *Oncol Nurs Forum* 1994; 21: 899–907.
- Moreau KL, Degarmo R, Langley J, McMahon C, Howley ET, Bassett DR Jr, Thompson DL. Increasing daily walking



- lowers blood pressure in postmenopausal women. *Med Sci Sports Exerc* 2001; 33: 1825–1831. Mosekilde L. Mechanisms in osteoporosis. *Ugeskr Laeger* 2001; 163: 1243–1246.
- Mourier A, Gautier JF, De Kerviler E, Bigard AX, Villette JM, Garnier JP, Duvallet A, Guezennec CY, Cathelineau G. Mobilization of visceral adipose tissue related to the improvement in insulin sensitivity in response to physical training in NIDDM. Effects of branched-chain amino acid supplements. *Diabetes Care* 1997; 20: 385–391.
- Moy CS, Songer TJ, LaPorte RE, Dorman JS, Kriska AM, Orchard TJ, Becker DJ, Drash AL. Insulin-dependent diabetes mellitus, physical activity, and death. *Am J Epidemiol* 1993; 137: 74–81.
- Muir JF, Pierson DJ, Simonds AK. Pulmonary rehabilitation. In: Simonds AK, Muir JF, Pierson DJ, eds. *BMJ Books*. 1996.
- Munneke M, de Jong Z, Zwinderman AH, Jansen A, Ronday HK, Peter WF, Boonman DC, Van Den Ende CH, Vliet Vlieland TP, Hazes JM. Adherence and satisfaction of rheumatoid arthritis patients with a long-term intensive dynamic exercise program (RAPIT program). *Arthritis Rheum* 2003; 49: 665–672.
- Munneke M, de Jong Z, Zwinderman AH, Ronday HK, Van Den Ende CH, Vliet Vlieland TP, Hazes JM. High intensity exercise or conventional exercise for patients with rheumatoid arthritis? Outcome expectations of patients, rheumatologists, and physiotherapists. *Ann Rheum Dis* 2004; 63: 804–808.
- Murphy MH, Hardman AE. Training effects of short and long bouts of brisk walking in sedentary women. *Med Sci Sports Exerc* 1998; 30: 152–157.
- Mutric N. Exercise as a treatment for moderate depression in the UK National Health Service [abstract]. London: Sports Council and Health Education Authority, 1988: 96–105. Myers J, Prakash M, Froelicher V, Do D, Partington S, Atwood JE. Exercise capacity and mortality among men referred for exercise testing. *N Engl J Med* 2002; 346: 793–801.
- Mynors-Wallis LM, Gath DH, Day A, Baker F. Randomised controlled trial of problem solving treatment, antidepressant medication, and combined treatment for major depression in primary care. *BMJ* 2000; 320: 26–30.
- Myrtek M, Villinger U. Psychological and physiological effects of a 5-week ergometer training in healthy young men (author's transl). *Med Klin* 1976; 71: 1623–1630.
- National Heart, Lung and Blood Institute. Obesity education initiative expert panel: clinical guidelines on the identification, evaluation and treatment of overweight and obesity in adults: the evidence report. 98-4083. Bethesda, MD: NIH, 1998: 1–228.
- National Institute of Health, N.H.L.a.B.I. (1995). Global initiative for asthma. NIH Publication No. 95-3659.
- National Institutes of Health Consensus Development Panel Triglyceride, LDL, and CHD. *JAMA* 1993; 269: 505–520.
- Nelson L, Jennings GL, Esler MD, Korner PI. Effect of changing levels of physical activity on blood-pressure and haemodynamics in essential hypertension. *Lancet* 1986; 2: 473–476.
- Nelson ME, Fiatarone MA, Morganti CM, Trice I, Greenberg RA, Evans WJ. Effects of high-intensity strength training on multiple risk factors for osteoporotic fractures. A randomized controlled trial. *JAMA* 1994; 272: 1909–1914.
- Nicklas BJ, Katzel LI, Busby-Whitehead J, Goldberg AP. Increases in high-density lipoprotein cholesterol with endurance exercise training are blunted in obese compared with lean men. *Metabolism* 1997; 46: 556–561.
- Nordesjo LO, Nordgren B, Wigren A, Kolstad K. Isometric strength and endurance in patients with severe rheumatoid arthritis or osteoarthritis in the knee joints. A comparative study in healthy men and women. *Scand J Rheumatol* 1983; 12: 152–156.
- Normandin EA, McCusker C, Connors M, Vale F, Gerardi D, ZuWallack RL. An evaluation of two approaches to exercise conditioning in pulmonary rehabilitation. *Chest* 2002; 121: 1085–1091.
- Notarius CF, Azevedo ER, Parker JD, Floras JS. Peak oxygen uptake is not determined by cardiac noradrenaline spill-over in heart failure. *Eur Heart J* 2002; 23: 800–805.
- Obrant KJ, Bengner U, Johnell O, Nilsson BE, Sernbo I. Increasing age-adjusted risk of fragility fractures: a sign of increasing osteoporosis in successive generations? *Calcif Tissue Int* 1989; 44: 157–167.
- Oka RK, De Marco T, Haskell WL, Botvinick E, Dae MW, Bolen K, Chatterjee K. Impact of a home-based walking and resistance training program on quality of life in patients with heart failure. *Am J Cardiol* 2000; 85: 365–369.
- Okumiya K, Matsubayashi K, Wada T, Kimura S, Doi Y, Ozawa T. Effects of exercise on neurobehavioral function in community-dwelling older people more than 75 years of age. *J Am Geriatr Soc* 1996; 44: 569–572.
- Oldridge N. Physical activity in primary and secondary prevention – there is a treatment gap. *Eur J Cardiovasc Prev Rehabil* 2003; 10: 317–318.
- Oldridge N, Guyatt G, Jones N, Crowe J, Singer J, Feeny D, McKelvie R, Runions J, Streiner D, Torrance G. Effects on quality of life with comprehensive rehabilitation after acute myocardial infarction. *Am J Cardiol* 1991; 67: 1084–1089.
- Oluseye KA. Cardiovascular responses to exercise in Nigerian women. *J Hum Hypertens* 1990a; 4: 77–79.
- Oluseye KA. Cardiovascular responses to exercise in Nigerian women. *J Hum Hypertens* 1990b; 4: 77–79.
- O'Reilly SC, Muir KR, Doherty M. Effectiveness of home exercise on pain and disability from osteoarthritis of the knee: a randomised controlled trial. *Ann Rheum Dis* 1999; 58: 15–19.
- Orenstein DM. The child and the adolescent athlete. In: Asthma and sports. Bar-Or O, ed. New York: Blackwell Science, 1995: 433–454.
- Ornish D, Brown SE, Scherwitz LW, Billings JH, Armstrong WT, Ports TA, McLanahan SM, Kirkeeide RL, Brand RJ, Gould KL. Can lifestyle changes reverse coronary heart disease? The Lifestyle Heart Trial. *Lancet* 1990; 336: 129–133.
- Owen A, Croucher L. Effect of an exercise programme for elderly patients with heart failure. *Eur J Heart Fail* 2000; 2: 65–70.
- Owens JF, Matthews KA, Wing RR, Kuller LH. Can physical activity mitigate the effects of aging in middle-aged women? *Circulation* 1992; 85: 1265–1270.
- Padilla J, Wallace JP, Park S. Accumulation of physical activity reduces blood pressure in pre- and hypertension. *Med*



- Sci Sports Exerc 2005; 37: 1264–1275.
- Paganini-Hill A, Chao A, Ross RK, Henderson BE. Exercise and other factors in the prevention of hip fracture: the Leisure World study. *Epidemiology* 1991; 2: 16–25.
- Palacio J, Galdiz JB, Bech JJ, Marinan M, Casadevall C, Martinez P, Gea J. Interleukin 10 and tumor necrosis factor alpha gene expression in respiratory and peripheral muscles. Relation to sarcolemmal damage. *Arch Bronconeumol* 2002; 38: 311–316.
- Pan XR, Li GW, Hu YH, Wang JX, Yang WY, An ZX, Hu ZX, Lin J, Xiao JZ, Cao HB, Liu PA, Jiang XG, Jiang YY, Wang JP, Zheng H, Zhang H, Bennett PH, Howard BV. Effects of diet and exercise in preventing NIDDM in people with impaired glucose tolerance. The Da Qing IGT and Diabetes Study. *Diabetes Care* 1997; 20: 537–544.
- Paolisso G, De Riu S, Marrazzo G, Verza M, Varricchio M, D'Onofrio F. Insulin resistance and hyperinsulinemia in patients with chronic congestive heart failure. *Metabolism* 1991; 40: 972–977.
- Park S, Jastremski CA, Wallace JP. Time of day for exercise on blood pressure reduction in dipping and nondipping hypertension. *J Hum Hypertens* 2005; 19: 597–605.
- Parnell MM, Holst DP, Kaye DM. Exercise training increases arterial compliance in patients with congestive heart failure. *Clin Sci (Lond)* 2002; 102: 1–7.
- Pasternak RC, Grundy SM, Levy D, Thompson PD. Spectrum of risk factors for CHD. *J Am Coll Cardiol* 1990; 27: 964–1047.
- Pavlou KN, Krey S, Stee WP. Exercise as an adjunct to weight loss and maintenance in moderately obese subjects. *Am J Clin Nutr* 1989; 49: 1115–1123.
- Pedersen BK, Holman-Goetz L. Exercise and the immune system: regulation, integration and adaptation. *Physiol Rev* 2000; 80: 1055–1081.
- Pedersen BK, Steensberg A, Schjerling P. Muscle-derived interleukin-6: possible biological effects. *J Physiol (London)* 2001; 536: 329–337.
- Penninx BW, Messier SP, Rejeski WJ, Williamson JD, DiBari M, Cavazzini C, Applegate WB, Pahor M. Physical exercise and the prevention of disability in activities of daily living in older persons with osteoarthritis. *Arch Intern Med* 2001; 161: 2309–2316.
- Perri MG, McAdoo WG, McAllister DA, Lauer JB, Yancey DZ. Enhancing the efficacy of behavior therapy for obesity: effects of aerobic exercise and a multicomponent maintenance program. *J Consult Clin Psychol* 1986; 54: 670–675.
- Perri MG, McAllister DA, Gange JJ, Jordan RC, McAdoo G, Nezu AM. Effects of four maintenance programs on the long-term management of obesity. *J Consult Clin Psychol* 1988; 56: 529–534.
- Pescatello LS, Franklin BA, Fagard R, Farquhar WB, Kelley GA, Ray CA. American College of Sports Medicine position stand. Exercise and hypertension. *Med Sci Sports Exerc* 2004; 36: 533–553.
- Peters S, Stanley I, Rose M, Kaney S, Salmon P. A randomized controlled trial of group aerobic exercise in primary care patients with persistent, unexplained physical symptoms. *Fam Pract* 2002; 19: 665–674.
- Peterson MG, Kovar-Toledano PA, Otis JC, Allegrante JP, MacKenzie CR, Gutin B, Kroll MA. Effect of a walking program on gait characteristics in patients with osteoarthritis. *Arthritis Care Res* 1993; 6: 11–16.
- Petrella RJ. How effective is exercise training for the treatment of hypertension? *Clin J Sport Med* 1998; 8: 224–231.
- Petrella RJ, Bartha C. Home based exercise therapy for older patients with knee osteoarthritis: a randomized clinical trial. *J Rheumatol* 2000; 27: 2215–2221.
- Piepoli MF, Davos C, Francis DP, Coats AJ. Exercise training meta-analysis of trials in patients with chronic heart failure (ExTraMATCH). *BMJ* 2004; 328: 189.
- Pinkney JH, Downs L, Hopton M, Mackness MI, Bolton CH. Endothelial dysfunction in Type 1 diabetes mellitus: relationship with LDL oxidation and the effects of vitamin E. *Diabetes Med* 1999; 16: 993–999.
- Pinto BM, Maruyama NC, Engebretson TO, Theborge RW. Participation in exercise, mood, and coping in survivors of early stage breast cancer survivors. *J Psychosocial Oncol* 1998; 16: 45–58.
- Posner JD, Gorman KM, Windsor-Landsberg L, Larsen J, Bleiman M, Shaw C, Rosenberg B, Knebl J. Low to moderate intensity endurance training in healthy older adults: physiological responses after four months. *J Am Geriatr Soc* 1992; 40: 1–7.
- Potempa K, Lopez M, Braun LT, Szidon JP, Fogg L, Tincknell T. Physiological outcomes of aerobic exercise training in hemiparetic stroke patients. *Stroke* 1995; 26: 101–105.
- Powell P, Bentall RP, Nye FJ, Edwards RH. Randomised controlled trial of patient education to encourage graded exercise in chronic fatigue syndrome. *BMJ* 2001; 322: 387–390.
- Pradhan AD, Manson JE, Rifai N, Buring JE, Ridker PM. C-reactive protein, interleukin 6, and risk of developing type 2 diabetes mellitus. *JAMA* 2001; 286: 327–334.
- Preisinger E, Alacamlioglu Y, Pils K, Saradeth T, Schneider B. Therapeutic exercise in the prevention of bone loss. A controlled trial with women after menopause. *Am J Phys Med Rehabil* 1995; 74: 120–123.
- Prince R, Devine A, Dick I, Criddle A, Kerr D, Kent N, Price R, Randell A. The effects of calcium supplementation (milk powder or tablets) and exercise on bone density in postmenopausal women. *J Bone Miner Res* 1995; 10: 1068–1075.
- Prince RL, Smith M, Dick IM, Price RI, Webb PG, Henderson NK, Harris MM. Prevention of postmenopausal osteoporosis. A comparative study of exercise, calcium supplementation, and hormone-replacement therapy. *N Engl J Med* 1991; 325: 1189–1195.
- Proctor DN, O'Brien PC, Atkinson EJ, Nair KS. Comparison of techniques to estimate total body skeletal muscle mass in people of different age groups. *Am J Physiol* 1999; 277: E489–E495.



- Prong NP. Short term effects of exercise on plasma lipids and lipoprotein in humans. *Sports Med* 2003; 16: 431–448.
- Province MA, Hadley EC, Hornbrook MC, Lipsitz LA, Miller JP, Mulrow CD, Ory MG, Sattin RW, Tinetti ME, Wolf SL. The effects of exercise on falls in elderly patients. A preplanned meta-analysis of the FICSIT Trials. Frailty and injuries: cooperative studies of intervention techniques. *JAMA* 1995; 273: 1341–1347.
- Pruitt LA, Taaie DR, Marcus R. Effects of a one-year high-intensity versus low-intensity resistance training program on bone mineral density in older women. *J Bone Miner Res* 1995; 10: 1788–1795.
- Pu CT, Johnson MT, Forman DE, Hausdorff JM, Roubenoff R, Foldvari M, Fielding RA, Singh MA. Randomized trial of progressive resistance training to counteract the myopathy of chronic heart failure. *J Appl Physiol* 2001; 90: 2341–2350.
- Quittan M, Sturm B, Wiesinger GF, Pacher R, Fialka-Moser V. Quality of life in patients with chronic heart failure: a randomized controlled trial of changes induced by a regular exercise program. *Scand J Rehabil Med* 1999; 31: 223–228.
- Rabasa-Lhoret R, Bourque J, Ducros F, Chiasson JL. Guidelines for premeal insulin dose reduction for postprandial exercise of different intensities and durations in type 1 diabetic subjects treated intensively with a basal-bolus insulin regimen (ultralente-lispro). *Diabetes Care* 2001; 24: 625–630.
- Radaelli A, Piepoli M, Adamopoulos S, Pipilis A, Clark SJ, Casadei B, Meyer TE, Coats AJ. Effects of mild physical activity, atenolol and the combination on ambulatory blood pressure in hypertensive subjects. *J Hypertens* 1992; 10: 1279–1282.
- Ram FS, Robinson SM, Black PN. Effects of physical training in asthma: a systematic review. *Br J Sports Med* 2000a; 34: 162–167.
- Ram FS, Robinson SM, Black PN. Physical training for asthma. *Cochrane Database Syst Rev* 2000bCD001116.
- Raz I, Hauser E, Bursztyjn M. Moderate exercise improves glucose metabolism in uncontrolled elderly patients with non-insulin-dependent diabetes mellitus. *Isr J Med Sci* 1994; 30: 766–770.
- Ready AE, Drinkwater DT, Ducas J, Fitzpatrick DW, Brereton DG, Oades SC. Walking program reduces elevated cholesterol in women postmenopause. *Can J Cardiol* 1995; 11: 905–912.
- Ready AE, Naimark B, Ducas J, Sawatzky JV, Boreskie SL, Drinkwater DT, Oosterveen S. Influence of walking volume on health benefits in women post-menopause. *Med Sci Sports Exerc* 1996; 28: 1097–1105.
- Redondo JR, Justo CM, Moraleda FV, Velayos YG, Puche JJ, Zubero JR, Hernandez TG, Ortells LC, Pareja MA. Long-term efficacy of therapy in patients with fibromyalgia: a physical exercise-based program and a cognitive-behavioral approach. *Arthritis Rheum* 2004; 51: 184–192.
- Regensteiner JG, Steiner JF, Hiatt WR. Exercise training improves functional status in patients with peripheral arterial disease. *J Vasc Surg* 1996; 23: 104–115.
- Reid CM, Dart AM, Dewar EM, Jennings GL. Interactions between the effects of exercise and weight loss on risk factors, cardiovascular haemodynamics and left ventricular structure in overweight subjects. *J Hypertens* 1994; 12: 291–301.
- Reid WD, Warren CPW. Ventilatory muscle strength and endurance training in elderly subjects and patients with chronic airflow limitation: a pilot study. *Physioter Can* 1984; 36: 305–311.
- Reilly JJ, Wilson ML, Summerbell CD, Wilson DC. Obesity: diagnosis, prevention, and treatment; evidence based answers to common questions. *Arch Dis Child* 2002; 86: 392–394.
- Reitman JS, Vasquez B, Klimes I, Nagulesparan M. Improvement of glucose homeostasis after exercise training in non-insulin-dependent diabetes. *Diabetes Care* 1984; 7: 434–441.
- Reuter M, Mutric N, Harris DV. Running as an adjunct to counseling in the treatment of depression [dissertation]. Pennsylvania: Pennsylvania State University, 1984.
- Revel M, Mayoux-Benhamou MA, Rabourdin JP, Bagheri F, Roux C. One-year psoas training can prevent lumbar bone loss in postmenopausal women: a randomized controlled trial. *Calcif Tissue Int* 1993; 53: 307–311.
- Richards SC, Scott DL. Prescribed exercise in people with fibromyalgia: parallel group randomised controlled trial. *BMJ* 2002; 325: 185.
- Ringbaek TJ, Broendum E, Hemmingsen L, Lybeck K, Nielsen D, Andersen C, Lange P. Rehabilitation of patients with chronic obstructive pulmonary disease. Exercise twice a week is not sufficient!. *Respir Med* 2000; 94: 150–154.
- Rissanen AM, Heliovaara M, Knekt P, Reunanen A, Aromaa A. Determinants of weight gain and overweight in adult Finns. *Eur J Clin Nutr* 1991; 45: 419–430.
- Robeer GG, Brandsma JW, van den Heuvel SP, Smit B, Oostendorp RA, Wittens CH. Exercise therapy for intermittent claudication: a review of the quality of randomised clinical trials and evaluation of predictive factors. *Eur J Vasc Endovasc Surg* 1998; 15: 36–43.
- Roberts CK, Barnard RJ. Effects of exercise and diet on chronic disease. *J Appl Physiol* 2005; 98: 3–30.
- Robertson MC, Campbell AJ, Gardner MM, Devlin N. Preventing injuries in older people by preventing falls: a meta-analysis of individual-level data. *J Am Geriatr Soc* 2002; 50: 905–911.
- Robillon JF, Sadoul JL, Jullien D, Morand P, Freychet P. Abnormalities suggestive of cardiomyopathy in patients with type 2 diabetes of relatively short duration. *Diabetes Metab* 1994; 20: 473–480.
- Rogers MW, Probst MM, Gruber JJ, Berger R, Boone JB Jr. Differential effects of exercise training intensity on blood pressure and cardiovascular responses to stress in borderline hypertensive humans. *J Hypertens* 1996; 14: 1369–1375.
- Rogind H, Bibow-Nielsen B, Jensen B, Moller HC, Frimodt-Moller H, Bliddal H. The effects of a physical training program on patients with osteoarthritis of the knees. *Arch Phys Med Rehabil* 1998; 79: 1421–1427.
- Rohan TE, Fu W, Hiller JE. Physical activity and survival from breast cancer. *Eur J Cancer Prev* 1995; 4: 419–424.
- Ronnemaa T, Mattila K, Lehtonen A, Kallio V. A controlled randomized study on the effect of long-term physical exercise on the metabolic control in type 2 diabetic patients. *Acta Med Scand* 1986; 220: 219–224.

- Ross R, Janssen I. Physical activity, total and regional obesity: dose-response considerations. *Med Sci Sports Exerc* 2001; 33: S521–S527.
- Ross R, Dagnone D, Jones PJ, Smith H, Paddags A, Hudson R, Janssen I. Reduction in obesity and related comorbid conditions after diet-induced weight loss or exercise-induced weight loss in men. A randomized, controlled trial. *Ann Intern Med* 2000; 133: 92–103.
- Ross R, Janssen I, Dawson J, Kungl AM, Kuk JL, Wong SL, Nguyen-Duy TB, Lee S, Kilpatrick K, Hudson R. Exercise-induced reduction in obesity and insulin resistance in women: a randomized controlled trial. *Obes Res* 2004; 12: 789–798.
- Rossy LA, Buckelew SP, Dorr N, Hagglund KJ, Thayer JF, McIntosh MJ, Hewett JE, Johnson JC. A meta-analysis of fibromyalgia treatment interventions. *Ann Behav Med* 1999; 21: 180–191.
- Roviaro S, Holmes DS, Holmsten RD. Influence of a cardiac rehabilitation program on the cardiovascular, psychological, and social functioning of cardiac patients. *J Behav Med* 1984; 7: 61–81.
- Rowlands AV, Ingledew DK, Eston RG. The effect of type of physical activity measure on the relationship between body fatness and habitual physical activity in children: a meta-analysis. *Ann Hum Biol* 2000; 27: 479–497.
- Ruderman NB, Ganda OP, Johansen K. The effect of physical training on glucose tolerance and plasma lipids in maturity-onset diabetes. *Diabetes* 1979; 28(Suppl. 1:89–92): 89–92.
- Sakai T, Ideishi M, Miura S, Maeda H, Tashiro E, Koga M, Kinoshita A, Sasaguri M, Tanaka H, Shindo M, Arakawa K. Mild exercise activates renal dopamine system in mild hypertensives. *J Hum Hypertens* 1998; 12: 355–362.
- Salman GF, Mosier MC, Beasley BW, Calkins DR. Rehabilitation for patients with chronic obstructive pulmonary disease: meta-analysis of randomized controlled trials. *J Gen Intern Med* 2003; 18: 213–221.
- Salmon P. Effects of physical exercise on anxiety, depression, and sensitivity to stress: a unifying theory. *Clin Psychol Rev* 2001; 21: 33–61.
- Saltin B, Helge JW. [Metabolic capacity of skeletal muscles and health]. *Ugeskr Laeger* 2000; 162: 2159–2164.
- Saltin B, Henriksson J, Nygaard E, Andersen P, Jansson E. Fiber types and metabolic potentials of skeletal muscles in sedentary man and endurance runners. *Ann N Y Acad Sci* 1977; 301(3–29): 3–29.
- Sant M, Capocaccia R, Verdecchia A, Gatta G, Micheli A, Mariotto A, Hakulinen T, Berrino F. Comparisons of colon-cancer survival among European countries: the Eurocare Study. *Int J Cancer* 1995; 63: 43–48.
- Santa-Clara H, Szymanski L, Fernhall B. Effect of exercise training on blood pressure in postmenopausal Caucasian and African-American women. *Am J Cardiol* 2003; 91: 1009–1011, A8.
- Santuz P, Baraldi E, Filippone M, Zacchello F. Exercise performance in children with asthma: is it different from that of healthy controls? *Eur Respir J* 1997; 10: 1254–1260.
- Sarljo-Lahteenkorva S, Rissanen A. Weight loss maintenance: determinants of long-term success. *Eat Weight Disord* 1998; 3: 131–135.
- Sarljo-Lahteenkorva S, Rissanen A, Kaprio J. A descriptive study of weight loss maintenance: 6 and 15 year follow-up of initially overweight adults. *Int J Obes Relat Metab Disord* 2000; 24: 116–125.
- Scalvini S, Marangoni S, Volterrani M, Schena M, Quadri A, Levi GF. Physical rehabilitation in coronary patients who have suffered from episodes of cardiac failure. *Cardiology* 1992; 80: 417–423.
- Schantz PG. Plasticity of human skeletal muscle with special reference to effects of physical training on enzyme levels of the NADH shuttles and phenotypic expression of slow and fast myofibrillar proteins. *Acta Physiol Scand* 1986; 558(Suppl.): 1–62.
- Scherer TA, Spengler CM, Owassapian D, Imhof E, Boutellier U. Respiratory muscle endurance training in chronic obstructive pulmonary disease: impact on exercise capacity, dyspnea, and quality of life. *Am J Respir Crit Care Med* 2000; 162: 1709–1714.
- Schilke JM, Johnson GO, Housh TJ, O'Dell JR. Effects of muscle-strength training on the functional status of patients with osteoarthritis of the knee joint. *Nurs Res* 1996; 45: 68–72.
- Schneider SH, Amorosa LF, Khachaturian AK, Ruderman NB. Studies on the mechanism of improved glucose control during regular exercise in type 2 (non-insulin-dependent) diabetes. *Diabetologia* 1984; 26: 355–360.
- Schuler G, Hambrecht R, Schlierf G, Niebauer J, Hauer K, Neumann J, Hoberg E, Drinkmann A, Bacher F, Grunze M. Regular physical exercise and low-fat diet. Effects on progression of coronary artery disease. *Circulation* 1992; 86: 1–11.
- Schulz KF, Chalmers I, Hayes RJ, Altman DG. Empirical evidence of bias. Dimensions of methodological quality associated with estimates of treatment effects in controlled trials. *JAMA* 1995; 273: 408–412.
- Schulz KH, Szlovak C, Schulz G, Gold S, Brechtel L, Braumann M, Koch U. [Implementation and evaluation of an ambulatory exercise therapy based rehabilitation program for breast cancer patients]. *Psychother Psychosom Med Psychol* 1998; 48: 398–407.
- Schulze PC, Gielen S, Schuler G, Hambrecht R. Chronic heart failure and skeletal muscle catabolism: effects of exercise training. *Int J Cardiol* 2002; 85: 141–149.
- Scott MG. The contributions of physical activity to psychological development. *Res Q* 1960; 31: 307–320.
- Segal R, Evans W, Johnson D, Smith J, Colletta S, Gayton J, Woodard S, Wells G, Reid R. Structured exercise improves physical functioning in women with stages I and II breast cancer: results of a randomized controlled trial. *J Clin Oncol* 2001; 19: 657–665.
- Segar ML, Katch VL, Roth RS, Garcia AW, Portner TI, Glickman SG, Haslanger S, Wilkins EG. The effect of aerobic exercise on self-esteem and depressive and anxiety symptoms among breast cancer survivors. *Oncol Nurs Forum* 1998; 25: 107–113.
- Seki E, Watanabe Y, Sunayama S, Iwama Y, Shimada K, Kawakami K, Sato M, Sato H, Mokuno H, Daida H. Effects



- of phase III cardiac rehabilitation programs on health-related quality of life in elderly patients with coronary artery disease: juntendo Cardiac Rehabilitation Program (J-CARP). *Circ J* 2003; 67: 73–77.
- Sexton H, Maere A, Dahl NH. Exercise intensity and reduction in neurotic symptoms. A controlled follow-up study. *Acta Psychiatr Scand* 1989; 80: 231–235.
- Sharpe MC, Archard LC, Banatvala JE, Borysiewicz LK, Clare AW, David A, Edwards RH, Hawton KE, Lambert HP, Lane RJ. A report – chronic fatigue syndrome: guidelines for research. *J R Soc Med* 1991; 84: 118–121.
- Shaw LW. Effects of a prescribed supervised exercise program on mortality and cardiovascular morbidity in patients after myocardial infarction. The national exercise and heart disease project. *Am J Cardiol* 1981; 48: 39–46.
- Shepherd RJ, Kavanagh T, Mertens DJ. On the prediction of physiological and psychological responses to aerobic training in patients with stable congestive heart failure. *J Cardiopulm Rehabil* 1998; 18: 45–51.
- Sikand G, Kondo A, Foreyt JP, Jones PH, Gotto AM Jr. Two-year follow-up of patients treated with a very-low-calorie diet and exercise training. *J Am Diet Assoc* 1988; 88: 487–488.
- Simpson K, Killian K, McCartney N, Stubbing DG, Jones NL. Randomised controlled trial of weightlifting exercise in patients with chronic airflow limitation. *Thorax* 1992; 47: 70–75.
- Sinaki M, Wahner HW, Orford KP, Hodgson SF. Efficacy of nonloading exercises in prevention of vertebral bone loss in postmenopausal women: a controlled trial. *Mayo Clin Proc* 1989; 64: 762–769.
- Singh NA, Clements KM, Fiatarone MA. A randomized controlled trial of progressive resistance training in depressed elders. *J Gerontol A Biol Sci Med Sci* 1997; 52: M27–M35.
- Sisto SA, LaManca J, Cordero DL, Bergen MT, Ellis SP, Drastal S, Boda WL, Tapp WN, Natelson BH. Metabolic and cardiovascular effects of a progressive exercise test in patients with chronic fatigue syndrome. *Am J Med* 1996; 100: 634–640.
- Sivarajan ES, Bruce RA, Lindsog BD, Almes MJ, Belanger L, Green B. Treadmill test responses to an early exercise program after myocardial infarction: a randomized study. *Circulation* 1982; 65: 1420–1428.
- Skelton DA, Beyer N. Exercise and injury prevention in older people. *Scand J Med Sci Sports* 2003; 13: 77–85.
- Skyrme-Jones RA, O'Brien RC, Luo M, Meredith IT. Endothelial vasodilator function is related to low-density lipoprotein particle size and low-density lipoprotein vitamin E content in type 1 diabetes. *J Am Coll Cardiol* 2000; 35: 292–299.
- Slemenda C, Heilman DK, Brandt KD, Katz BP, Mazza SA, Braunstein EM, Byrd D. Reduced quadriceps strength relative to body weight: a risk factor for knee osteoarthritis in women? *Arthritis Rheum* 1998; 41: 1951–1959.
- Slentz CA, Aiken LB, Houmar JA, Bales CW, Johnson JL, Tanner CJ, Duscha BD, Kraus WE. Inactivity, exercise, and visceral fat. STRIDE: a randomized, controlled study of exercise intensity and amount. *J Appl Physiol* 2005; 99: 1613–1618.
- Slentz CA, Duscha BD, Johnson JL, Ketchum K, Aiken LB, Samsa GP, Houmar JA, Bales CW, Kraus WE. Effects of the amount of exercise on body weight, body composition, and measures of central obesity: stride – a randomized controlled study. *Arch Intern Med* 2004; 164: 31–39.
- Sly RM, Harper RT, Rosselot I. The effect of physical conditioning upon asthmatic children. *Ann Allergy* 1972; 30: 86–94.
- Smidt GL, Lin SY, O'Dwyer KD, Blaupied PR. The effect of high-intensity trunk exercise on bone mineral density of postmenopausal women. *Spine* 1992; 17: 280–285.
- Soo K, Furler SM, Samaras K, Jenkins AB, Campbell LV, Chisholm DJ. Glycemic responses to exercise in IDDM after simple and complex carbohydrate supplementation. *Diabetes Care* 1996; 19: 575–579.
- Sopko G, Leon AS, Jacobs DR Jr., Foster N, Moy J, Kuba K, Anderson JT, Casal D, McNally C, Frantz I. The effects of exercise and weight loss on plasma lipids in young obese men. *Metabolism* 1985; 34: 227–236.
- Specchia G, De Servi S, Scire A, Assandri J, Berzuini C, Angoli L, La Rovere MT, Cobelli F. Interaction between exercise training and ejection fraction in predicting prognosis after a first myocardial infarction. *Circulation* 1996; 94: 978–982.
- Stamano BA, Braun LT, Rosenson RS. The accumulative effects of physical activity in hypertensive postmenopausal women. *J Cardiovasc Risk* 2001; 8: 283–290.
- Stahle A, Lindquist I, Mattsson E. Important factors for physical activity among elderly patients one year after an acute myocardial infarction. *Scand J Rehabil Med* 2000; 32: 111–116.
- Stamler J, Vaccaro O, Neaton JD, Wentworth D. Diabetes, other risk factors, and 12-yr cardiovascular mortality for men screened in the multiple risk factor intervention trial. *Diabetes Care* 1993; 16: 434–444.
- Stefanick ML, Wood PD. Physical activity, lipid and lipid transport. In: Bouchard C, Shepherd RJ, Stephens T, eds. *Physical activity, fitness, health. International proceedings and consensus statement*. Champaign, IL: Human Kinetics, 1994: 417–437.
- Stefanick ML, Mackey S, Sheehan M, Ellsworth N, Haskell WL, Wood PD. Effects of diet and exercise in men and postmenopausal women with low levels of HDL cholesterol and high levels of LDL cholesterol. *N Engl J Med* 1998b; 339: 12–20.
- Stefanick ML, Mackey S, Sheehan M, Ellsworth N, Haskell WL, Wood PD. Effects of diet and exercise in men and postmenopausal women with low levels of HDL cholesterol and high levels of LDL cholesterol. *N Engl J Med* 1998a; 339: 12–20.
- Stern MJ, Gorman PA, Kaslow L. The group counseling v exercise therapy study. A controlled intervention with subjects following myocardial infarction. *Arch Intern Med* 1983; 143: 1719–1725.
- Stewart KJ. Exercise and hypertension. In: Roitman J, ed. *ACSM's resource manual for guidelines for exercise testing and prescription*. Baltimore: Lippincott Williams Wilkins, 2001.

- Stewart KJ. Exercise training and the cardiovascular consequences of type 2 diabetes and hypertension: plausible mechanisms for improving cardiovascular health. *JAMA* 2002; 288: 1622–1631.
- Sullivan MJ, Green HJ, Cobb FR. Skeletal muscle biochemistry and histology in ambulatory patients with long-term heart failure. *Circulation* 1990; 81: 518–527.
- Sullivan MJ, Higginbotham MB, Cobb FR. Exercise training in patients with severe left ventricular dysfunction. Hemodynamic and metabolic effects. *Circulation* 1988; 78: 506–515.
- Sullivan MJ, Higginbotham MB, Cobb FR. Exercise training in patients with chronic heart failure delays ventilatory anaerobic threshold and improves submaximal exercise performance. *Circulation* 1989a; 79: 324–329.
- Sullivan MJ, Knight JD, Higginbotham MB, Cobb FR. Relation between central and peripheral hemodynamics during exercise in patients with chronic heart failure. Muscle blood flow is reduced with maintenance of arterial perfusion pressure. *Circulation* 1989b; 80: 769–781.
- Suter E, Marti B, Tschopp A, Wanner HU, Wenk C, Gutzwiller F. Effects of self-monitored jogging on physical fitness, blood pressure and serum lipids: a controlled study in sedentary middle-aged men. *Int J Sports Med* 1990; 11: 425–432.
- Svendsen OL, Hassager C, Christiansen C. Effect of an energy-restrictive diet, with or without exercise, on lean tissue mass, resting metabolic rate, cardiovascular risk factors, and bone in overweight postmenopausal women. *Am J Med* 1993; 95: 131–140.
- Svendsen OL, Hassager C, Christiansen C. Six months' follow-up on exercise added to a short-term diet in overweight postmenopausal women – effects on body composition, resting metabolic rate, cardiovascular risk factors and bone. *Int J Obes Relat Metab Disord* 1994; 18: 692–698.
- Svendsen OL, Heitmann BL, Mikkelsen KL, Raben A, Rytting KJ, Sørensen TIA, et al. Fedme i Danmark. En rapport fra Dansk task force on obesity. *Ugeskr Laeger* 2001; 163.
- Swann IL, Hanson CA. Double-blind prospective study of the effect of physical training on childhood asthma. In: Oseid S, Edwards AM, eds. *The asthmatic child – in play and sport*. London: Pitman Books Ltd, 1983: 318–325.
- Sylvester KL. Pilot study: investigation of the effect of hydrotherapy in the treatment of osteoarthritic hips. *Clin Rehabil* 1989; 4: 228.
- Takenaka K, Sakamoto T, Amano K, Oku J, Fujinami K, Murakami T, Toda I, Kawakubo K, Sugimoto T. Left ventricular filling determined by Doppler echocardiography in diabetes mellitus. *Am J Cardiol* 1988; 61: 1140–1143.
- Tan KH, De Cossart L, Edwards PR. Exercise training and peripheral vascular disease. *Br J Surg* 2000; 87: 553–562.
- Tan RA, Spector SL. Exercise-induced asthma: diagnosis and management. *Ann Allergy Asthma Immunol* 2002; 89: 226–235.
- Tanabe Y, Urata H, Kiyonaga A, Ikeda M, Tanaka H, Shindo M, Arakawa K. Changes in serum concentrations of taurine and other amino acids in clinical antihypertensive exercise therapy. *Clin Exp Hypertens A* 1989; 11: 149–165.
- Tanaka H, Bassett DR Jr, Howley ET, Thompson DL, Ashraf M, Rawson FL. Swimming training lowers the resting blood pressure in individuals with hypertension. *J Hypertens* 1997; 15: 651–657.
- Tarumi N, Iwasaka T, Takahashi N, Sugiura T, Morita Y, Sumimoto T, Nishiue T, Inada M. Left ventricular diastolic filling properties in diabetic patients during isometric exercise. *Cardiology* 1993; 83: 316–323.
- TASC Management of peripheral arterial disease. *Eur J Vasc Endovasc Surg* 2000; 19: S1–S250.
- Taylor A. Physiological response to a short period of exercise training in patients with chronic heart failure. *Physiother Res Int* 1999; 4: 237–249.
- Taylor CB, Jatulis DE, Winkley MA, Rockhill BJ, Kraemer HC. Effects of life-style on body mass index change. *Epidemiology* 1994; 5: 599–603.
- Taylor CB, Miller NH, Smith PM, DeBusk RF. The effect of a home-based, case-managed, multifactorial risk-reduction program on reducing psychological distress in patients with cardiovascular disease. *J Cardiopulm Rehabil* 1997; 17: 157–162.
- Taylor RS, Brown A, Ebrahim S, Jolliffe J, Noorani H, Rees K, Skidmore B, Stone JA, Thompson DR, Oldridge N. Exercise-based rehabilitation for patients with coronary heart disease: systematic review and meta-analysis of randomized controlled trials. *Am J Med* 2004; 116: 682–692.
- Tessier D, Menard J, Fulop T, Ardilouze J, Roy M, Dubuc N, Dubois M, Gauthier P. Effects of aerobic physical exercise in the elderly with type 2 diabetes mellitus. *Arch Gerontol Geriatr* 2000; 31: 121–132.
- The P.R.E.COR group. Comparison of a rehabilitation programme, a counselling programme and usual care after an acute myocardial infarction: results of a long-term randomized trial. *Eur Heart J* 1991; 12: 612–616.
- Thomas KS, Muir KR, Doherty M, Jones AC, O'Reilly SC, Bassey EJ. Home based exercise programme for knee pain and knee osteoarthritis: randomised controlled trial. *BMJ* 2002; 325: 752.
- Thune I. Physical exercise in rehabilitation program for cancer patients? *J Altern Complement Med* 1998; 4: 205–207.
- Thune I, Furberg AS. Physical activity and cancer risk: dose-response and cancer, all sites and site-specific. *Med Sci Sports Exerc* 2001; 33: S530–S550.
- Thune I, Smeland S. Is physical activity important in treatment and rehabilitation of cancer patients? *Tidsskr Nor Laegeforen* 2000; 120: 3302–3304.
- Tipton CM. Exercise and hypertension. In: Shephard RJ, Miller HJS, eds. *Exercise and the heart in health and disease*. New York: Marcel Dekker Inc, 1999: 463–488.
- Toobert DJ, Glasgow RE, Radcliffe JL. Physiologic and related behavioral outcomes from the women's lifestyle heart trial. *Ann Behav Med* 2000; 22: 1–9.
- Topp R, Woolley S, Hornyak J III, Khuder S, Kahaleh B. The effect of dynamic versus isometric resistance training on pain and functioning among adults with osteoarthritis of the knee. *Arch Phys Med Rehabil* 2002; 83: 1187–1195.



- Tran ZV, Weltman A. Differential effects of exercise on serum lipid and lipoprotein levels seen with changes in body weight. A meta-analysis. *JAMA* 1985; 254: 919–924.
- Tran ZV, Weltman A, Glass GV, Mood DP. The effects of exercise on blood lipids and lipoproteins: a meta-analysis of studies. *Med Sci Sports Exerc* 1983; 15: 393–402.
- Trovati M, Carta Q, Cavalot F, Vitali S, Banaudi C, Lucchina PG, Fiocchi F, Emanuelli G, Lenti G. Influence of physical training on blood glucose control, glucose tolerance, insulin secretion, and insulin action in non-insulin-dependent diabetic patients. *Diabetes Care* 1984; 7: 416–420.
- Tsai JC, Chang WY, Kao CC, Lu MS, Chen YJ, Chan P. Beneficial effect on blood pressure and lipid profile by programmed exercise training in Taiwanese patients with mild hypertension. *Clin Exp Hypertens* 2002a; 24: 315–324.
- Tsai JC, Liu JC, Kao CC, Tomlinson B, Kao PF, Chen JW, Chan P. Beneficial effects on blood pressure and lipid profile of programmed exercise training in subjects with white coat hypertension. *Am J Hypertens* 2002b; 15: 571–576.
- Tsuda K, Yoshikawa A, Kimura K, Nishio I. Effects of mild aerobic physical exercise on membrane fluidity of erythrocytes in essential hypertension. *Clin Exp Pharmacol Physiol* 2003; 30: 382–386.
- Tudor-Locke C, Bell RC, Myers AM, Harris SB, Ecclestone NA, Lauzon N, Rodger NW. Controlled outcome evaluation of the first step program: a daily physical activity intervention for individuals with type II diabetes. *Int J Obes Relat Metab Disord* 2004; 28: 113–119.
- Tudor-Locke C, Myers AM, Rodger NW. Formative evaluation of the first step program: a practical intervention to increase daily physical activity. *Can J Diabetes Care* 2000; 24: 34–38.
- Tudor-Locke C, Myers AM, Bell RC, Harris S, Rodger NW. Preliminary outcome evaluation of the first step program: a daily physical activity intervention for individuals with type 2 diabetes. *Patient Educ Couns* 2002; 47: 23–28.
- Tudor-Locke CE, Myers AM, Rodger NW. Development of a theory-based daily activity intervention for individuals with type 2 diabetes. *Diabetes Educ* 2001; 27: 85–93.
- Tuomilehto J, Lindstrom J, Eriksson JG, Valle TT, Hamalainen H, Ilanne-Parikka P, Keinanen-Kiukkaanniemi S, Laakso M, Louheranta A, Rastas M, Salminen V, Uusitupa M. Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med* 2001; 344: 1343–1350.
- Tuominen JA, Karonen SL, Melamies L, Bolli G, Koivisto VA. Exercise-induced hypoglycaemia in IDDM patients treated with a short-acting insulin analogue. *Diabetologia* 1995; 38: 106–111.
- Tyni-Lenne R, Dencker K, Gordon A, Jansson E, Sylven C. Comprehensive local muscle training increases aerobic working capacity and quality of life and decreases neurohormonal activation in patients with chronic heart failure. *Eur J Heart Fail* 2001; 3: 47–52.
- Tyni-Lenne R, Gordon A, Sylven C. Improved quality of life in chronic heart failure patients following local endurance training with leg muscles. *J Card Fail* 1996; 2: 111–117.
- Tyni-Lenne R, Gordon A, Europe E, Jansson E, Sylven C. Exercise-based rehabilitation improves skeletal muscle capacity, exercise tolerance, and quality of life in both women and men with chronic heart failure. *J Card Fail* 1998; 4: 9–17.
- Tyni-Lenne R, Gordon A, Jansson E, Bermann G, Sylven C. Skeletal muscle endurance training improves peripheral oxidative capacity, exercise tolerance, and health-related quality of life in women with chronic congestive heart failure secondary to either ischemic cardiomyopathy or idiopathic dilated cardiomyopathy. *Am J Cardiol* 1997; 80: 1025–1029.
- Tyni-Lenne R, Gordon A, Jensen-Urstad M, Dencker K, Jansson E, Sylven C. Aerobic training involving a minor muscle mass shows greater efficiency than training involving a major muscle mass in chronic heart failure patients. *J Card Fail* 1999; 5: 300–307.
- US Department of Health and Human Services. Physical activity and health: a report of the surgeon general. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, 1996: 1–278.
- UK Prospective Diabetes Study (UKPDS) Group. Effect of intensive blood-glucose control with metformin on complications in overweight patients with type 2 diabetes (UKPDS 34). *Lancet* 1998; 352: 854–865.
- Urata H, Tanabe Y, Kiyonaga A, Ikeda M, Tanaka H, Shindo M, Arakawa K. Antihypertensive and volume-depleting effects of mild exercise on essential hypertension. *Hypertension* 1987; 9: 245–252.
- van Baar ME, Assendelft WJ, Dekker J, Oostendorp RA, Bijlsma JW. Effectiveness of exercise therapy in patients with osteoarthritis of the hip or knee: a systematic review of randomized clinical trials. *Arthritis Rheum* 1999; 42: 1361–1369.
- van Baar ME, Dekker J, Oostendorp RA, Bijl D, Voorn TB, Bijlsma JW. Effectiveness of exercise in patients with osteoarthritis of hip or knee: nine months' follow up. *Ann Rheum Dis* 2001; 60: 1123–1130.
- van Baar ME, Dekker J, Oostendorp RA, Bijl D, Voorn TB, Lemmens JA, Bijlsma JW. The effectiveness of exercise therapy in patients with osteoarthritis of the hip or knee: a randomized clinical trial. *J Rheumatol* 1998; 25: 2432–2439.
- van Dale D, Saris WH, ten Hoor F. Weight maintenance and resting metabolic rate 18–40 months after a diet/exercise treatment. *Int J Obes* 1990; 14: 347–359.
- Van Den Ende CH, Breedveld FC, le Cessie S, Dijkmans BA, de Mug AW, Hazes JM. Effect of intensive exercise on patients with active rheumatoid arthritis: a randomised clinical trial. *Ann Rheum Dis* 2000a; 59: 615–621.
- Van Den Ende CH, Hazes JM, le Cessie S, Mulder WJ, Belfor DG, Breedveld FC, Dijkmans BA. Comparison of high and low intensity training in well controlled rheumatoid arthritis. Results of a randomised clinical trial. *Ann Rheum Dis* 1996; 55: 798–805.
- Van Den Ende CH, Vliet Vlieland TP, Munneke M, Hazes JM. Dynamic exercise therapy for rheumatoid arthritis. *Cochrane Database Syst Rev* 2000b CD000322.
- van der Poest CE, van der WH, Patka P, Roos JC, Lips P. Long-term consequences of fracture of the lower leg:

- cross-sectional study and long-term longitudinal follow-up of bone mineral density in the hip after fracture of lower leg. *Bone* 1999; 24: 131–134.
- Van der Wiel HE, Lips P, Nauta J, Patka P, Haarman HJ, Teule GJ. Loss of bone in the proximal part of the femur following unstable fractures of the leg. *J Bone Joint Surg Am* 1994; 76: 230–236.
- Van Hoof R, Hespel P, Fagard R, Lijnen P, Staessen J, Amery A. Effect of endurance training on blood pressure at rest, during exercise and during 24 hours in sedentary men. *Am J Cardiol* 1989a; 63: 945–949.
- Van Hoof R, Hespel P, Fagard R, Lijnen P, Staessen J, Amery A. Effect of endurance training on blood pressure at rest, during exercise and during 24 hours in sedentary men. *Am J Cardiol* 1989b; 63: 945–949.
- van Santen M, Bolwijn P, Landewe R, Verstappen F, Bakker C, Hidding A, van Der KD, Houben H, van der LS. High or low intensity aerobic fitness training in fibromyalgia: does it matter? *J Rheumatol* 2002; 29: 582–587.
- Vanninen E, Uusitupa M, Siitonen O, Laitinen J, Lansimies E. Habitual physical activity, aerobic capacity and metabolic control in patients with newly-diagnosed type 2 (non-insulin-dependent) diabetes mellitus: effect of 1-year diet and exercise intervention. *Diabetologia* 1992; 35: 340–346.
- Varray AL, Mercier JG, Prefaut CG. Individualized training reduces excessive exercise hyperventilation in asthmatics. *Int J Rehabil Res* 1995; 18: 297–312.
- Varray AL, Mercier JG, Terral CM, Prefaut CG. Individualized aerobic and high intensity training for asthmatic children in an exercise readaptation program. Is training always helpful for better adaptation to exercise? *Chest* 1991; 99: 579–586.
- Veale D, Le Fevre K. Aerobic exercise in the adjunctive treatment of depression: a randomised controlled trial [abstract]. London: Sports Council and Health Education Authority, 1988: 106–111.
- Veale D, Le Fevre K, Pantelis C, de SV, Mann A, Sargeant A. Aerobic exercise in the adjunctive treatment of depression: a randomized controlled trial. *J R Soc Med* 1992; 85: 541–544.
- Vecchio C, Cobelli F, Opasich C, Assandri J, Poggi G, Grieco R. [Early functional evaluation and physical rehabilitation in patients with wide myocardial infarction (author's transl)]. *G Ital Cardiol* 1981; 11: 419–429.
- Veje K, Hyllested JL, Ostergaard K. Osteoarthritis. Pathogenesis, clinical features and treatment. *Ugeskr Laeger* 2002; 164: 3173–3179.
- Vermeulen A, Lie KI, Durrer D. Effects of cardiac rehabilitation after myocardial infarction: changes in coronary risk factors and long-term prognosis. *Am Heart J* 1983; 105: 798–801.
- Veves A, Saouaf R, Donaghue VM, Mullooly CA, Kistler JA, Giurini JM, Horton ES, Fielding RA. Aerobic exercise capacity remains normal despite impaired endothelial function in the micro- and macrocirculation of physically active IDDM patients. *Diabetes* 1997; 46: 1846–1852.
- Vroman NB, Healy JA, Kertzer R. Cardiovascular response to lower body negative pressure (LBNP) following endurance training. *Aviat Space Environ Med* 1988; 59: 330–334.
- Wadden TA, Vogt RA, Foster GD, Anderson DA. Exercise and the maintenance of weight loss: 1-year follow-up of a controlled clinical trial. *J Consult Clin Psychol* 1998; 66: 429–433.
- Wadell K, Henriksson-Larsen K, Lundgren R. Physical training with and without oxygen in patients with chronic obstructive pulmonary disease and exercise-induced hypoxaemia. *J Rehabil Med* 2001; 33: 200–205.
- Walker KZ, Piers LS, Putt RS, Jones JA, O'Dea K. Effects of regular walking on cardiovascular risk factors and body composition in normoglycemic women and women with type 2 diabetes. *Diabetes Care* 1999; 22: 555–561.
- Wallace BA, Cumming RG. Systematic review of randomized trials of the effect of exercise on bone mass in pre- and postmenopausal women. *Calcif Tissue Int* 2000; 67: 10–18.
- Wallberg-Henriksson H, Gunnarsson R, Henriksson J, Ostman J, Wahren J. Influence of physical training on formation of muscle capillaries in type 1 diabetes. *Diabetes* 1984; 33: 851–857.
- Wallberg-Henriksson H, Gunnarsson R, Rossner S, Wahren J. Long-term physical training in female type 1 (insulin-dependent) diabetic patients: absence of significant effect on glycaemic control and lipoprotein levels. *Diabetologia* 1986; 29: 53–57.
- Walsh MF, Flynn TJ. A 54-month evaluation of a popular very low calorie diet program. *J Fam Pract* 1995; 41: 231–236.
- Wang JS, Jen CJ, Chen HI. Effects of exercise training and deconditioning on platelet function in men. *Arterioscler Thromb Vasc Biol* 1995; 15: 1668–1674.
- Wang JS, Jen CJ, Chen HI. Effects of chronic exercise and deconditioning on platelet function in women. *J Appl Physiol* 1997; 83: 2080–2085.
- Wasserman DH, Zinman B. Exercise in individuals with IDDM. *Diabetes Care* 1994; 17: 924–937.
- Wearden AJ, Morriss RK, Mullis R, Strickland PL, Pearson DJ, Appleby L, Campbell IT, Morris JA. Randomised, double-blind, placebo-controlled treatment trial of fluoxetine and graded exercise for chronic fatigue syndrome. *Br J Psychiatry* 1998; 172(485–90): 485–490.
- Wei J, Xu H, Davies JL, Hemmings GP. Increase of plasma IL-6 concentration with age in healthy subjects. *Life Sci* 1992; 51: 1953–1956.
- Weiner P, Azgad Y, Ganam R. Inspiratory muscle training combined with general exercise reconditioning in patients with COPD. *Chest* 1992; 102: 1351–1356.
- Westby MD, Wade JP, Rangno KK, Berkowitz J. A randomized controlled trial to evaluate the effectiveness of an exercise program in women with rheumatoid arthritis taking low dose prednisone. *J Rheumatol* 2000; 27: 1674–1680.
- Whelton SP, Chin A, Xin X, He J. Effect of aerobic exercise on blood pressure: a meta-analysis of randomized, controlled trials. *Ann Intern Med* 2002; 136: 493–503.
- Wickham CA, Walsh K, Cooper C, Barker DJ, Margetts BM, Morris J, Bruce SA. Dietary calcium, physical activity, and risk of hip fracture: a prospective study. *BMJ* 1989; 299: 889–892.



- Wielenga RP, Huisveld IA, Bol E, Dunselman PH, Erdman RA, Baselier MR, Mosterd WL. Safety and effects of physical training in chronic heart failure. Results of the chronic heart failure and graded exercise study (CHANGE). *Eur Heart J* 1999; 20: 872–879.
- Wiesinger GF, Pleiner J, Quittan M, Fuchsjäger-Mayrl G, Crevenna R, Nuhr MJ, Francesconi C, Seit HP, Francesconi M, Fialka-Moser V, Wolzt M. Health related quality of life in patients with long-standing insulin dependent (type 1) diabetes mellitus benefits of regular physical training. *Wien Klin Wochenschr* 2001; 113:670–675.
- Wigers SH, Stiles TC, Vogel PA. Effects of aerobic exercise versus stress management treatment in fibromyalgia. A 4.5 year prospective study. *Scand J Rheumatol* 1996; 25: 77–86.
- Wijkstra PJ, TenVergert EM, van der Mark TW, Postma DS, Van Altena R, Kraan J, Koeter GH. Relation of lung function, maximal inspiratory pressure, dyspnoea, and quality of life with exercise capacity in patients with chronic obstructive pulmonary disease. *Thorax* 1994; 49: 468–472.
- Wijnen JA, Kool MJ, van Baak MA, Kuipers H, de Haan CH, Verstappen FT, Struijker Boudier HA, Van Bortel LM. Effect of exercise training on ambulatory blood pressure. *Int J Sports Med* 1994; 15: 10–15.
- Wilhelmsen L, Sanne H, Elmfeldt D, Grimby G, Tibblin G, Wedel H. A controlled trial of physical training after myocardial infarction. Effects on risk factors, nonfatal reinfarction, and death. *Prev Med* 1975; 4: 491–508.
- Willenheimer R, Erhardt L, Cline C, Rydberg E, Israelsson B. Exercise training in heart failure improves quality of life and exercise capacity. *Eur Heart J* 1998; 19: 774–781.
- Williamson DF, Madans J, Anda RF, Kleinman JC, Kahn HS, Byers T. Recreational physical activity and ten-year weight change in a US national cohort. *Int J Obes Relat Metab Disord* 1993; 17: 279–286.
- Wilson JR, Mancini DM, Dunkman WB. Exertional fatigue due to skeletal muscle dysfunction in patients with heart failure. *Circulation* 1993; 87: 470–475.
- Wilson MG, Michet CJ Jr., Ilstrup DM, Melton LJ. III. Idiopathic symptomatic osteoarthritis of the hip and knee: a population-based incidence study. *Mayo Clin Proc* 1990; 65: 1214–1221.
- Wing RR, Epstein LH, Paternostro-Bayles M, Kriska A, Nowalk MP, Gooding W. Exercise in a behavioural weight control programme for obese patients with Type 2 (non-insulin-dependent) diabetes. *Diabetologia* 1988; 31: 902–909.
- Wing RR, Venditti E, Jakicic JM, Polley BA, Lang W. Lifestyle intervention in overweight individuals with a family history of diabetes. *Diabetes Care* 1998; 21: 350–359.
- Winningham ML, MacVicar MG. The effect of aerobic exercise on patient reports of nausea. *Oncol Nurs Forum* 1988; 15: 447–450.
- Winocour PH, Durrington PN, Bhatnagar D, Mbewu AD, Ishola M, Mackness M, Arrol S. A cross-sectional evaluation of cardiovascular risk factors in coronary heart disease associated with type 1 (insulin-dependent) diabetes mellitus. *Diabetes Res Clin Pract* 1992; 18: 173–184.
- Wolfe F. The fibromyalgia syndrome: a consensus report on fibromyalgia and disability. *J Rheumatol* 1996; 23: 534–539.
- Wolfe F, Mitchell DM, Sibley JT, Fries JF, Bloch DA, Williams CA, Spitz PW, Hagen M, Kleinheksel SM, Cathey MA. The mortality of rheumatoid arthritis. *Arthritis Rheum* 1994; 37: 481–494.
- Wolfe F, Smythe HA, Yunus MB, Bennett RM, Bombardier C, Goldenberg DL, Tugwell P, Campbell SM, Abeles M, Clark P. The American College of Rheumatology 1990 criteria for the classification of fibromyalgia. Report of the Multicenter Criteria Committee. *Arthritis Rheum* 1990; 33: 160–172.
- Wolke I, van Croonenborg JJ, Kemper HC, Kostense PJ, Twisk JW. The effect of exercise training programs on bone mass: a meta-analysis of published controlled trials in pre- and postmenopausal women. *Osteoporos Int* 1999; 9: 1–12.
- Wood PD, Stefanick ML, Dreon DM, Frey-Hewitt B, Garay SC, Williams PT, Superko HR, Fortmann SP, Albers JJ, Vranizan KM. Changes in plasma lipids and lipoproteins in overweight men during weight loss through dieting as compared with exercise. *N Engl J Med* 1988; 319: 1173–1179.
- Wood RH, Reyes R, Welsch MA, Favaloro-Sabatier J, Sabatier M, Matthew LC, Johnson LG, Hooper PF. Concurrent cardiovascular and resistance training in healthy older adults. *Med Sci Sports Exerc* 2001; 33: 1751–1758.
- World Health Organization. Rehabilitation and comprehensive secondary prevention after acute myocardial infarction. *EURO Rep Stud* 1983; 84.
- Wosornu D, Bedford D, Ballantyne D. A comparison of the effects of strength and aerobic exercise training on exercise capacity and lipids after coronary artery bypass surgery. *Eur Heart J* 1996; 17: 854–863.
- Yamanouchi K, Shinozaki T, Chikada K, Nishikawa T, Ito K, Shimizu S, Ozawa N, Suzuki Y, Maeno H, Kato K. Daily walking combined with diet therapy is a useful means for obese NIDDM patients not only to reduce body weight but also to improve insulin sensitivity. *Diabetes Care* 1995; 18: 775–778.
- Yasuda I, Kawakami K, Shimada T, Tanigawa K, Murakami R, Izumi S, Morioka S, Kato Y, Moriyama K. Systolic and diastolic left ventricular dysfunction in middle-aged asymptomatic non-insulin-dependent diabetics. *J Cardiol* 1992; 22: 427–438.
- Yki-Jarvinen H, DeFronzo RA, Koivisto VA. Normalization of insulin sensitivity in type I diabetic subjects by physical training during insulin pump therapy. *Diabetes Care* 1984; 7: 520–527.
- Yu CM, Li LS, Ho HH, Lau CP. Long-term changes in exercise capacity, quality of life, body anthropometry, and lipid profiles after a cardiac rehabilitation program in obese patients with coronary heart disease. *Am J Cardiol* 2003; 91: 321–325.
- Zavaroni I, Bonini L, Gasparini P, Barilli AL, Zuccarelli A, Dall'Aglio E, Delsignore R, Reaven GM. Hyperinsulinemia in a normal population as a predictor of non-insulin-dependent diabetes mellitus, hypertension, and coronary heart disease: the Barilla factory revisited. *Metabolism* 1999; 48: 989–994.