

Reference

Chapter 1

1. Mackay J, Mensah GA, Mendis S, Greenlund K. World Health Organization. The Atlas of Heart Disease and Stroke. Geneva: World Health Organization; 2004.
2. Rosamond W, Flegal K, Friday, G, et al. Heart disease and stroke statistics 2007 update. A report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation* 2007; 115:e69–e171.
3. Lopez-Yunez A. The management of stroke patients by neurologists: Common questions and new observations. *Semin Neurol* 2002; 22(1):53–61.
4. Feigin V, Lawes CM, Bennett DA, et al. Stroke epidemiology. A review of population-based studies of incidence, prevalence, and case-fatality in the late 20th century. *Lancet Neurol* 2003; 2:43–53.
5. Pendlebury S. Worldwide under-funding of stroke research. *Int J Stroke* 2007; 2:80–84.
6. Wolfe C. The impact of stroke. *Br Med Bull* 2000; 56:275–286.
7. Khadilkar A, Phillips K, Jean N, Lamothe C, Milne S, Sarnecka J. Ottawa panel evidence-based clinical practice guidelines for post-stroke rehabilitation. *Top Stroke Rehabil* 2006; 13(2):1–269.
8. Gordon N, Gulanick M, Costa F, et al. Physical activity and exercise recommendations for stroke survivors: An American Heart Association scientific statement from the Council on Clinical Cardiology, Subcommittee on Exercise, Cardiac Rehabilitation, and Prevention; the Council on Cardiovascular Nursing; the Council on Nutrition, Physical Activity, and Metabolism; and the Stroke Council. *Circulation* 2004; 109(5):2031–2041.
9. Caplan L. *Stroke. A Clinical Approach*, 3rd ed. Boston: Butterworth Heinemann; 2000.
10. Collins C. Pathophysiology and classification of stroke. *Nurs Stand* 2007; 21(28):35–39.
11. Williams LS, Kroenke K, Bakas T, et al. Care management of post-Stroke depression: A randomized, controlled trial. *Stroke* 2007; 38(3):998–1003.
12. Rich DQ, Gaziano JM, Kurth T. Geographic patterns in overall and specific cardiovascular disease incidence in apparently healthy men in the United States. *Stroke* 2007; 38(8):2221–2227.
13. Stewart DG. Stroke rehabilitation. 1. Epidemiologic aspects and acute management. *Arch Phys Med Rehabil* 1999; 80(5 Suppl 1):S4–S7.
14. Kwok T, Lo RS, Wong E, Wai-Kwong T, Mok V, Kai-Sing W. Quality of life of stroke survivors: A 1-year follow-up study. *Arch Phys Med Rehabil* 2006; 87(9):1177–1182; quiz 1287.
15. Lai SM, Studenski S, Richards L, et al. Therapeutic exercise and depressive symptoms after stroke. *J Am Geriatr Soc* 2006; 54(2):240–247.
16. Teasell R, Foley N, Bhogal S, Bagg S, Jutai J. Evidence-based practice and setting basic standards for stroke rehabilitation in Canada. *Top Stroke Rehabil* 2006; 13(3):59–65.
17. Black-Schaffer RM, Kirssteins AE, Harvey RL. Stroke rehabilitation. 2 Co-morbidities and complications. *Arch Phys Med Rehabil* 1999; 80(5 Suppl 1):S8–S16.
18. Rimmer JH, Shenoy SS. Impact of exercise on targeted secondary conditions. In: Field MJ, Jette AM, Martin L, eds. *Workshop on Disability in America*. Washington, DC: National Academies Press; 2006.
19. Roth EJ, Lovell L. Community skill performance and its association with the ability to perform everyday tasks by stroke survivors one year following rehabilitation discharge. *Top Stroke Rehabil* 2007; 14(1):48–56.
20. Ebrahim S. *Clinical Epidemiology of Stroke*. New York: Oxford University Press; 1990.
21. Gordon NF. *Stroke. Your Complete Exercise Guide*. The Cooper Clinic and Research Institute fitness series. Champaign, IL: Human Kinetics; 1993.
22. Kelly JF. Stroke rehabilitation for elderly patients. In: Kemp B, Brummel-Smith K, Ramsdell JW, Eds. *Geriatric Rehabilitation*. Boston: Little, Brown and Company; 1990: 61–89.
23. Pauls JA, Reed KL. *Quick Reference to Physical Therapy*. Gaithersburg, MD: Aspen; 1996.
24. Ramasubbu R, Robinson RG, Flint AJ, Kosier T, Price TR. Functional impairment associated with acute poststroke depression: the Stroke Data Bank Study. *J Neuropsychiatry Clin Neurosci* 1998; 10(1): 26–33.
25. Alberts MJ, Easton JD. Stroke best practices: A team approach to evidence-based care. *JAMA* 2004; 96(4 Supplement):5S–20S.
26. Macko RF, DeSouza CA, Tretter LD, et al. Treadmill aerobic exercise training reduces the energy expenditure and cardiovascular demands of hemiparetic gait in chronic stroke patients. A preliminary report. *Stroke* 1997; 28(2):326–330.
27. Wolf PA, Claggett GP, Easton JP, et al. Preventing Ischemic Stroke in Patients with Prior Stroke and Transient Ischemic Attack: A Statement for Healthcare Professionals from the Stroke Council of the American Heart Association. *Stroke* 1999; 30:1991–1994. CHAPTER 1 Stroke 17
28. Niemi ML, Laaksonen R, Kotila M, Waltimo O. Quality of life 4 years after stroke. *Stroke* 1988; 19:1101–1107.
29. Black-Schaffer RM, Kirssteins AE, Harvey RL. Stroke rehabilitation: Co-morbidities and complications. *Arch Phys Med Rehabil* 1999; 80:S8–S16.
30. Corr S, Bayer A. Poor functional status of stroke patients after hospital discharge: Scope for intervention? *Br J Occup Ther* 1992; 55:383–385.
31. Dean CM, Richards CL, Malouin F. Task-related circuit training improves performance of locomotor tasks in chronic stroke: A randomized controlled pilot trial. *Arch Phys Med Rehabil* 2000; 81(4): 409–417.
32. Hill K, Ellis P, Berhardt J, Maggs P, Hull S. Balance and mobility Outcomes for stroke patients: A comprehensive audit. *Aust J Physiother* 1997; 43:173–180.
33. Goldie PA, Matyas TA, Evans OM. Deficit and change in gait velocity during rehabilitation after stroke. *Arch Phys Med Rehabil* 1996; 77:1074–1082.
34. Duncan PW, Samsa GP, Weinberger M, et al. Health status of individuals with mild stroke. *Stroke* 1997; 28:740–745.
35. Mackay-Lyons MJ, Makrides L. Exercise capacity early after stroke. *Arch Phys Med Rehabil* 2002; 83:1697–1702.
36. Macko RF, Katzell LJ, Yataco A, et al. Low-velocity graded treadmill stress testing in hemiparetic stroke patients. *Stroke* 1997; 28(5): 988–992.
37. Rimmer JH, Wang E. Aerobic exercise training in stroke survivors. *Top Stroke Rehabil* 2005; 12(1):17–30.
38. Ivey F, Macko RF, Ryan AS, Hafer-Macko CE. Cardiovascular health and fitness after stroke. *Top Stroke Rehabil* 2005; 12(1):1–16.
39. Pang M, Eng JJ, Gylfadottir S, et al. The use of aerobic exercise training in improving aerobic capacity in individuals with stroke: A meta-analysis. *Clin Rehabil* 2006; 20:97–111.
40. Duncan P, Studenski S, Richards L, et al. Randomized clinical trial of therapeutic exercise in subacute stroke. *Stroke* 2003; 34(9): 2173–2180.
41. Chu KS, Eng JJ, Dawson AS, Harris JE, Ozkaplan A, Gylfadottir S. Water-based exercise for cardiovascular fitness in people with chronic stroke: A randomized controlled trial. *Arch Phys Med Rehabil* 2004; 85(6):870–874.
42. Potempa K, Braun LT, Szidon JP, Fogg L, Tincknell T. Physiological Outcomes of aerobic exercise training in hemiparetic stroke patients. *Stroke* 1995; 26:101–105.
43. Rimmer JH, Riley B, Creviston T, Nicola T. Exercise training in a predominantly African-American group of stroke survivors. *Med Sci Sports Exerc* 2000; 32(12):1990–1996.
44. Potempa K, Braun LT, Tinknell T, Popovich J. Benefits of aerobic exercise after stroke. *Sports Med* 1996; 21:337–346.
45. Binder EF, Burge SJ, Spina R. Peak aerobic power as an important component of physical performance in older women. *J Gerontol Med Sci* 1999; 54A (M353–M356).
46. Malbut-Shennan K, Young A. The physiology of physical performance and training in old age. *Coron Artery Dis* 1999; 10:37–42.
47. Rimmer JH, Nicola T. Stroke and exercise. In: ACSM's Resources for Clinical Exercise Physiology for Special Populations. Baltimore: Lippincott Williams & Wilkins; 2002.
48. Rimmer JH, Riley B, Creviston C, Nicola T. Exercise training in a Predominantly African-American group of stroke survivors. *Med Sci Sports Exerc* 2000; 32:1990–1996.
49. American College of Sports Medicine. ACSM's Guidelines for Exercise Testing and Prescription, 7th ed. Baltimore: Lippincott Williams & Wilkins; 2006.

50. Teixeira-Salmela LF, Olney SJ, Nadeau S, Brouwer B. Muscle strengthening and physical conditioning to reduce impairment and disability in chronic stroke survivors. *Arch Phys Med Rehabil* 1999;80(10):1211–1218.
51. Duncan PW. Synthesis of intervention trials to improve motor recovery following stroke. *Top Stroke Rehabil* 1997; 3:1–20.
52. Tang A, Sibley KM, Thomas SG, McIlroy WE, Brooks D. Maximal exercise test results in subacute stroke. *Arch Phys Med Rehabil* 2006;87(8):1100–1105.
53. McArdle W, Katch FI, Katch VL. *Exercise Physiology: Energy, Nutrition and Performance*. Baltimore Lippincott Williams & Williams; 2006.
54. Weiss A, Suzuki T, Bean J, Fielding RA. High intensity strength training improves strength and functional performance after stroke. *Am J Phys Med Rehabil* 2000;79(4):369–376.
55. Ouellette M, LeBrasseur NK, Bean JF, et al. High-intensity resistance training improves muscle strength, self-reported function, and disability in long-term stroke survivors. *Stroke* 2004; 35:1404–1409.
56. Rimmer JH, Wang E. Obesity prevalence among a group of Chicago residents with disabilities. *Arch Phys Med Rehabil* 2005; 86(7):1461–1464.
57. Duncan PW, Lai SM, Keighley J. Defining post-stroke recovery: Implications for design and interpretation of drug trials. *Neuropharmacology* 2000;39(5):835–841.
58. Meek C, Pollock A, Potter J, et al. A systematic review of exercisetrials post stroke. *Clin Rehabil* 2003; 17:6–13.
59. Ivey FM, Hafer-Macko CE, Macko RF. Exercise rehabilitation after stroke. *NeuroRx* 2006; 3(4):439–450.
60. Pang MY, Eng JJ, Dawson AS, et al. A community-based fitness and mobility exercise program for older adults with chronic stroke: A randomized, controlled trial. *J Am Geriatr Soc* 2005; 53(10):1667–1674.
61. Patten C, Lexell J, Brown HE. Weakness and strength training in persons with poststroke hemiplegia: rationale, method, and efficacy. *J Rehabil Res Dev* 2004; 41(3A):293–312.
62. Rimmer J. Exercise and physical activity in persons aging with a physical disability. *Phys Med Rehabil Clin N Am* 2005; 16(1):41–56.
63. Macko RF, Ivey FM, Forrester LW, et al. Treadmill exercise rehabilitation improves ambulatory function and cardiovascular fitness in patients with chronic stroke. A randomized, controlled trial. *Stroke* 2005; 26:2206–2211.
64. Macko RF, Ivey FM, Forrester LW, et al. Treadmill exercise rehabilitation improves ambulatory function and cardiovascular fitness in patients with chronic stroke: A randomized, controlled trial. *Stroke* 2005;36(10):2206–2211.
65. Ivey FM, Ryan AS, Hafer-Macko CE, Goldberg AP, Macko RF. Treadmill aerobic training improves glucose tolerance and indices of insulin sensitivity in disabled stroke survivors: a preliminary report. *Stroke* 2007; 38(10):2752–2758.
66. Rimmer JH, Rauworth AE, Wang E, Nicola TL, Hill B. A preliminary study to examine the effects of aerobic and therapeutic (non-aerobic) exercise on cardiorespiratory fitness and coronary risk reduction in stroke survivors. *Arch Phys Med Rehabil*. In press.
67. Macko RF, Smith GV, Dobrovolsky CL, Sorkin JD, Goldberg AP, Silver KH. Treadmill training improves fitness reserve in chronic stroke patients. *Arch Phys Med Rehabil* 2001; 82(7):879–884.
68. Rimmer J. The conspicuous absence of people with disabilities in public fitness and recreation facilities: Lack of interest or lack of access? *Am J Health Promot* 2005; 19:327–329.
69. Rimmer J, Wang E, Smith D. Barriers associated with exercise and community access for individuals with stroke. *J Rehab Res Develop* 2008; 45:315–322.
70. Sullivan K, Brown DA, Klassen T, Mulroy S, Winstein CJ. Effects of task-specific locomotor and strength training in adults who were ambulatory after stroke: Results of the STEPS randomized clinical trial. *Phys Ther*. 87:1580–1602.
71. Yang Y, Wang R, Lin K, Chu M, Chan R. Task-oriented progressive resistance strength training improves muscle strength and functional performance in individuals with stroke. *Clin Rehab* 2006; 20:860–870.

Chapter 2

1. Club L. Memorandum on terminology and classification of ‘cerebral palsy.’ *Cerebral Palsy Bulletin* 1959;1:27–35.
2. Mutch L, Alberman E, Hagberg B, et al. Cerebral palsy epidemiology: Where are we now and where are we going? *Dev Med Child Neurol* 1992;34:574–551.
3. Albright AL. Spasticity and movement disorders in cerebral palsy. *J Child Neurol* 1996; 11(Suppl 1):S1–S4.
4. Nelson KB, Ellenberg JH. Children who “outgrew” cerebral palsy. *Pediatrics* 1982; 69:529–536.
5. Arvidsson J, Hagberg B. Delayed onset dyskinetic ‘cerebral palsy’: A late effect of perinatal asphyxia? *Acta Paediatr Scand* 1990; 79: 1121–1123.
6. Lesny I. The development of athetosis. *Dev Med Child Neurol* 1968; 10:441–446.
7. Murphy CC, Yeargin-Allsopp M, Decoufle P, et al. Prevalence of cerebral palsy among ten year old children in metropolitan Atlanta, 1985 through 1987. *J Pediatr* 1993; 123:S13–S19.
8. Arens LJ, Molteno CD. A comparative study of postnatally-acquired cerebral palsy in Cape Town. *Dev Med Child Neurol* 1989;31: 246–254.
9. Al-Rajah S, Bademosi O, Awada A, et al. Cerebral palsy in Saudi Arabia: A case-control study of risk factors. *Dev Med Child Neurol* 1991; 33:1048–1052.
10. Bhasin TK, Brocksen S, Avchen RN, Braun KVN. Prevalence of four developmental disabilities among children aged 8 years— Metropolitan Atlanta developmental disabilities surveillance program, 1996–2000. *MMWR* 55(SS-1) 2006:1–9.
11. Sinha G, Corry P, Subesinghe D, et al. Prevalence and type of cerebral palsy in a British ethnic community: the role of consanguinity. *Dev Med Child Neurol* 1997;39:259–262.
12. Odding E, Roebroeck ME, Stam HJ. The epidemiology of cerebral palsy: Incidence, impairments and risk factors. *Disabil Rehabil* 2006;28(4):183–191.
13. DeLuca PA. The musculoskeletal management of children with cerebral palsy. *Pediatr Clin North Am* 1996;43(5): 1135–1150.
14. Stanley F, Blair E, Alberman E. *Cerebral Palsies: Epidemiology and Causal Pathways*. Clinics in Developmental Medicine. Number 87. London: Mac Keith Press; 2000:1–51.
15. Saito N, Ebara S, Ohotsuka K, et al. Natural history of scoliosis in spastic cerebral palsy. *Lancet* 1998;351:1687–1692.
16. Del Giudice E, Staiano A, Capano G, et al. Gastrointestinal manifestations in children with cerebral palsy. *Brain Dev* 1999;21: 307–311.
17. Bandini LG, Schoeller DA, Fukagawa NK, et al. Body composition and energy expenditure in adolescents with cerebral palsy or myelodysplasia. *Pediatr Res* 1991;29:70–77.
18. Van den Berg-Emons RJ, van Baak MA, Speth L, et al. Physical training of school children with spastic cerebral palsy: Effects on daily activity, fat mass and fitness. *Int J Rehabil* 1998;21:179–194.
19. Damiano DL, Kelly LE, Vaughn CL. Effects of quadriceps femoris muscle strengthening on crouch gait in children with spastic cerebral palsy. *Phys Ther* 1995;75:658–671.
20. Bar-Or O, Inbar O, Spira R. Physiological effects of a sports rehabilitation program on cerebral palsied and post-poliomyelitic adolescents. *Med Sci Sports Exerc* 1976;8:157–161.
21. Hutzler Y, Chacham A, Bergman U, et al. Effects of a movement and swimming program on vital capacity and water orientation skills of children with cerebral palsy. *Dev Med Child Neurol* 1998; 40: 176–181.
22. Laskin JJ. Cerebral palsy. In: *ACSM’s Exercise Management for Persons with Chronic Diseases and Disabilities*, 2nd ed. Champaign, IL: Human Kinetics; 2003.
23. Bowen TR, Lennon N, Castagno P, et al. Variability of energy consumption measures in children with cerebral palsy. *J Pediatr Orthop* 1998;18:738–742.
24. Pranzatelli MR. Oral pharmacotherapy for the movement disorders of cerebral palsy. *J Child Neurol* 1996;11(Suppl 1):S13–S22.
25. Albright AL. Intrathecal baclofen in cerebral palsy movement disorders. *J Child Neurol* 1996;11(Suppl 1):S29–S35.
26. Penn RD, Kroin JS. Intrathecal baclofen alleviates spinal cord spasticity. *Lancet* 1984; 1:1078.
27. Penn RD, Savoy SM, Corcos D, et al. Intrathecal baclofen for severe spinal spasticity. *N Engl J Med* 1989;320:1517–1521.
28. Goldstein M. The treatment of cerebral palsy: What we know, what we don’t know. *J Pediatr* 2004; 145:S45–S46.

29. Bodensteiner JB. The management of cerebral palsy: Subjectivity and conundrum. *J Child Neurol* 1996;11(2):75–76.
30. Almeida GL, Campbell SK, Girolami GL, et al. Multidimensional assessment of motor function in a child with cerebral palsy following intrathecal administration of baclofen. *Phys Ther* 1997; 77(7): 751–764.
31. Gerszten PC, Albright AL, Johnstone GF. Intrathecal baclofen infusion and subsequent orthopedic surgery in patients with spastic cerebral palsy. *J Neurosurg* 1998;88:1009–1013.
32. Albright AL, Shultz BL. Plasma baclofen levels in children receiving continuous intrathecal baclofen infusion. *J Child Neurol* 1999;14: 408–409.
33. Albright AL. Baclofen in the treatment of cerebral palsy. *J Child Neurol* 1996;11:77–83.
34. Koman LA, Mooney JF, Smith BP. Neuromuscular blockade in the management of cerebral palsy. *J Child Neurol* 1996;11(Suppl 1): S23–S28.
35. Gordon N. The role of botulinum toxin type A in treatment—With special reference to children. *Brain Dev* 1999;21:147–151.
36. Gouch JL, Sandell TV. Botulinum toxin for spasticity and athetosis in children with cerebral palsy. *Arch Phys Med Rehabil* 1996;77: 508–511.
37. Carr LJ, Cosgrove AP, Gringras P, et al. Position paper on the use of botulinum toxin in cerebral palsy. *Arch Dis Child* 1998;79:271–73.
38. Hazneci GB, Vurucu, S, Örs, F, Tan AF, Gençdoğ'an, S, Alp, T. Factors affecting the functional level in children with cerebral palsy. *Turk J Phys Med Rehab* 2006;52:105–109.
39. Massin M, Allington N. Role of exercise testing in the functional assessment of cerebral palsy children after botulinum A toxin injection. *J Pediatr Orthop* 1999;19:362–365.
40. Barwood S, Baillieu C, Boyd R, Brereton K, Low J, Natrass G, Graham HK. Analgesic effects of botulinum toxin A: A randomized, placebo-controlled clinical trial. *Dev Med Child Neurol* 2000; 42:116–121.
41. Jefferson RJ. Botulinum toxin in the management of cerebral palsy. *Dev Med Child Neurol* 2004; 46:491–499.
42. Ubhi T, Bhakta BB, Ives HL, Allgar V, Roussounis SH. Randomised double blind placebo-controlled trial of the effect of botulinum toxin on walking in cerebral palsy. *Arch Dis Child* 2000; 83:481–487.
43. Forssberg H, Tedroff KB. Botulinum toxin treatment in cerebral palsy: Intervention with poor evaluation? *Dev Med Child Neurol* 1997;39:635–640.
44. Lannin N, Scheinberg A, Clark K. AACPD systematic review of the effectiveness of therapy for children with cerebral palsy after botulinum toxin A injections. *Dev Med Child Neurol* 2006; 48:533–539.
45. Reeuwijk A, van Schie PEM. Effects of botulinum toxin type A on upper limb function in children with cerebral palsy: A systematic review. *Clin Rehabil* 2006; 20:375–387.
46. Ade-Hall RA, Moore AP. Botulinum toxin type A in the treatment of lower limb spasticity in cerebral palsy. *The Cochrane Database of Systemic Reviews, The Cochrane Library* 2007;(2).
47. Davis DW. Review of cerebral palsy. Part II: Identification and intervention. 1997;16(4):19–25.
48. Precht HFR. State of the art of a new functional assessment of the young nervous system. An early predictor of cerebral palsy. *Early Hum Dev* 1997;50:1–11.
49. Barry MJ. Physical therapy interventions for patients with movement disorders from cerebral palsy. *J Child Neurol* 1996; 11(Suppl 1):S51–S60.
50. Law M, Pollock N, Rosenbaum P, et al. A comparison of intensive neurodevelopmental therapy plus casting and a regular occupational therapy program for children with cerebral palsy. *Dev Med Child Neurol* 1997;39:664–670.
51. Weindling AM, Hallam P, Gregg J, et al. A randomized controlled trial of early physiotherapy for high-risk infants. *Acta Paediatr* 1996; 85:1107–1111.
52. Bower E, McLellan DL, Arney J, et al. A randomised controlled trial of different intensities of physiotherapy and different goal-setting procedures in 44 children with cerebral palsy. *Dev Med Child Neurol* 1996; 38:226–237.
53. Chad KE, Bailey DA, McKay HA, et al. The effect of a weight-bearing physical activity program on bone mineral content and estimated volumetric density in children with spastic cerebral palsy. *J Pediatr* 1999; 135:115–117.
54. Carlson WE, Vaughan CL, Damiano DL, et al. Orthotic management of gait in spastic diplegia. *Am J Phys Med Rehabil* 1997;76:219–225.
55. Mattsson E, Andersson C. Oxygen cost, walking speed, and perceived exertion in children with cerebral palsy when walking with anterior and posterior walkers. *Dev Med Child Neurol* 1997;39:671–676.
56. Damiano DL, Abel MF. Functional outcomes of strength training in spastic cerebral palsy. *Arch Phys Med Rehabil* 1998;79:119–25.
57. Morton R. New surgical interventions for cerebral palsy and the place of gait analysis. *Dev Med Child Neurol* 1999;41:424–428.
58. Wright FV, Sheil EM, Drake JM, et al. Evaluation of selective dorsal rhizotomy for the reduction of spasticity in cerebral palsy: A randomized controlled trial. *Dev Med Child Neurol* 1998;40:239–247.
59. Steinbok P, Reiner AM, Beauchamp R, et al. A randomized clinical trial to compare selective posterior rhizotomy plus physiotherapy with physiotherapy alone in children with spastic diplegic cerebral palsy. *Dev Med Child Neurol* 1997;39:178–184.
60. Steinbok P, Reiner A, Kestle JRW. Therapeutic electrical stimulation following selective posterior rhizotomy in children with spastic diplegic cerebral palsy: A randomized clinical trial. *Dev Med Child Neurol* 1997;39:515–520.
61. Buckon CE, Thomas SS, Aiona MD, et al. Assessment of upper extremity function in children with spastic diplegia before and after selective dorsal rhizotomy. *Dev Med Child Neurol* 1995;38:967–975.
62. Subramanian N, Vaughan CL, Peter JC, et al. Gait before and 10 years after rhizotomy in children with cerebral palsy spasticity. *J Neurosurg* 1998; 88:1014–1019.
63. Abbott R. Sensory rhizotomy for the treatment of childhood spasticity. *J Child Neurol* 1996; 11(Suppl 1):S36–S42.
64. Engsberg JR, Olree KS, Ross SA, et al. Spasticity and strength changes as a function of selective dorsal rhizotomy. *J Neurosurg* 1998; 88:1020–1026.
65. Engsberg JR, Ross SA, Park TS. Changes in ankle spasticity and strength following selective dorsal rhizotomy and physical therapy for spastic cerebral palsy. *J Neurosurg* 1999; 91:727–732.
66. DeSalles AAF. Role of stereotaxis in the treatment of cerebral palsy. *J Child Neurol* 1996; 11(Suppl 1):S43–S50.
67. Speelman JD, van Manen J. Cerebral palsy and stereotactic neuro-surgery: Long term results. *J Neurol Neurosurg Psychiatry* 1989;52:23–30.
68. Van Heest AE, House JH, Cariello C. Upper extremity surgical treatment of cerebral palsy. *J Hand Surg* 1999; 24A:323–330.
69. Renshaw TS, Green NE, Griffin PP, et al. Cerebral palsy: Orthopaedic management. *Instr Course Lect* 1996; 45:475–490. CHAPTER 2 Cerebral Palsy 33
70. Lonstein JE, Beck K. Hip dislocation and subluxation in cerebral palsy. *J Pediatr Orthop* 1986;6:521–526.
71. Hagglund G, Andersson S, Duppe H, Lauge-Pedersen H, Nordmark E, Westbom L. Prevention of dislocation of the hip in children with cerebral palsy: The first ten years of a population based prevention program. *J Bone Joint Surg (Br)* 2005;87-B:95–101.
72. Terjesen T, Lie GD, Hyldmo AA, Knaus A. Adductor tenotomy in spastic cerebral palsy. *Acta Orthop* 2005;76(1):128–137.
73. Preseido A, Oh C, Dabney KW, Miller F. Soft-tissue releases to treat spastic hip subluxation in children with cerebral palsy. *J Bone Joint Surg* 2005;87-A(4):832–841.
74. Dobson F, Boyd RN, Parrot J, Natrass GR, Graham HK. Hip surveillance in children with cerebral palsy: Impact on the surgical management of spastic hip disease. *J Bone Joint Surg (Br)* 2002;85-B(5):720–726.
75. House JH, Gwaathmey FW, Fidler MO. A dynamic approach to the thumb-in palm deformity in cerebral palsy. *J Bone Joint Surg* 1981; 63A:216–225.
76. Davids JR, Ounpuu S, DeLuca PA, Davis RB. Optimization of walking ability of children with cerebral palsy. *J Bone Joint Surg* 2003; 85-A(11):2224–2234.
77. Teli MG, Cinnella P, Vincitorio F, Lovi A, Grava G, Brayda-Bruno M. Spinal fusion with Cotrel-Dubouset instrumentation for neuropathic scoliosis in patients with cerebral palsy. *Spine* 2006; 31(4):E441–E447.
78. Vialle R, Delecourt C, Morin C. Surgical treatment of scoliosis with pelvic obliquity in cerebral palsy. *Spine* 2006; 31(13): 1461–1466.
79. Onley, SJ, Wright MJ. Cerebral palsy. In: Campbell SK, Vander Linden DW, Palisano RJ, Eds. *Physical Therapy for Children*. Philadelphia: Saunders Elsevier; 2006:625–664.
80. Morgan AM, Aldag JC. Early identification of cerebral palsy using a profile of abnormal motor patterns. *Pediatrics* 1996; 98:692–697.
81. Campbell SK. Quantifying the effects of interventions for movement disorders resulting from cerebral palsy. *J Child Neurol* 1996; 11(Suppl 1):S61–S70.

82. Tecklin JS. *Pediatric Physical Therapy*. 4th ed. Philadelphia: Lippincott Williams & Wilkins; 2007.
83. Laskin JJ. Physiological adaptations to concurrent muscular strength and aerobic endurance training in functionally active people with a physical disability. Unpublished doctoral dissertation, University of Alberta, 2001.
84. Carroll KL, Leiser J, Paisley TS. Cerebral palsy: Physical activity and sport. *Curr Sports Med Rep* 2006;5(6):319–322.
85. Damiano DL. Activity, activity, activity: Rethinking our physical therapy approach to cerebral palsy (III STEP Series). *Phys Ther* 2006; 86(11):1534–1537.
86. American College of Sports Medicine. *ACSM's Guidelines for Exercise Testing and Prescription*, 7th ed. Baltimore: Lippincott Williams & Wilkins; 2005.
87. American College of Sports Medicine. *ACSM's Resource Manual for Guidelines for Exercise Testing and Prescription*, 5th ed. Philadelphia: Lippincott Williams & Wilkins; 2005.
88. DePauw KP, Gavron SJ. *Disability Sport*, 2nd ed. Champaign, IL: Human Kinetics; 2005.
89. Shinohara T, Suzuki N, Oba M, Kawasumi M, Kimizuka M, Mita K. Effect of exercise at the AT point for children with cerebral palsy. *Bull NYU Hosp Jt Dis* 2002;61(1/2):63–67.
90. Unnithan VB, Kenne EM, Logan L, Collier S, Turk M. The effect of partial body weight support on the oxygen cost of walking in children and adolescents with spastic cerebral palsy. *Pediatr Exerc Sci* 2006;18(1):11–22.
91. Bhambhani YN, Holland LJ, Steaward RD. Maximal aerobic power in cerebral palsied wheelchair athletes: Validity and reliability. *Arch Phys Med Rehabil* 1992;73(3):246–252.
92. Bhambhani YN, Holland LJ, Steaward RD. Anaerobic threshold in wheelchair athletes with cerebral palsy: Validity and reliability. *Arch Phys Med Rehabil* 1993;74(3):305–311.
93. Suzuki N, Oshimi Y, Shinohara T, Kawasumi M, Mita M. Exercise intensity based on heart rate while walking in spastic cerebral palsy. *Bull NYU Hosp Jt Dis* 2001;60(1):18–22.
94. Bohannon RW. Is the measurement of muscle strength appropriate in patients with brain lesions? A special communication. *Phys Ther* 1989;69(3):225–229.
95. Bohannon RW, Walsh S. Nature, reliability, and predictive value of muscle performance measures in patients with hemiparesis following stroke. *Arch Phys Med Rehabil* 1992; 73:721–725.
96. Allen J, Dodd K, Taylor N, McBurney H, Larkin H. Strength training can be enjoyable and beneficial for adults with cerebral palsy. *Disabil Rehabil* 2004;26(19):1121–1128.
97. Bohannon RW, Larkin PA, Smith MB, et al. Relationship between static muscle strength deficits and spasticity in stroke patients with hemiparesis. *Phys Ther* 1987;67(7):1068–1071.
98. Bohannon RW. Relative decreases in knee extension torque with increased knee extension velocities in stroke patients with hemiparesis. *Phys Ther* 1987;67(7):1218–1220.
99. Bohannon RW, Smith MB. Assessment of strength deficits in eight paretic upper extremity muscle groups of stroke patients with hemiplegia. *Phys Ther* 1987;67(4):522–525.
100. Bohannon RW, Smith MB. Upper extremity strength deficits in hemiplegic stroke patients: Relationship between admission and discharge assessment and time since onset. *Arch Phys Med Rehabil* 1987; 68:155–157.
101. Bennett SE, Karnes JL. *Neurological Disabilities: Assessment and Treatment*. Philadelphia: Lippincott Williams & Wilkins; 1998.
102. Fernandez JE, Pitetti KH, Betzen MT. Physiological capacities of individuals with cerebral palsy. *Hum Factors* 1990; 32(4):457–466.
103. Parker DF, Carriere L, Hebestreit H, et al. Anaerobic endurance and peak muscle power in children with spastic cerebral palsy. *Am J Dis Child* 1992;146(9):1069–1073.
104. Parker DF, Carriere L, Hebestreit H, et al. Muscle performance and gross motor function of children with spastic cerebral palsy. *Dev Med Child Neurol* 1993;35(1):17–23.

Chapter 3

1. Web site home page-National Multiple Sclerosis Society [Internet]. New York, NY: National MS Society; [cited 2007]. Available from: <http://www.nmms.org>.
2. Rosati G. Descriptive epidemiology of MS in Europe in the 1980's: A critical overview. *Ann Neurol* 1994;36:5164–5174.
3. Weinstenker BG. Epidemiology of MS. *Neurol Clin* 1996; 14:291–308.
4. Sibley WA, Bamford CR, Clark K. Clinical viral infections and MS. *Lancet* 1985;1:1313–1315.
5. Panitch HS. Influence of infection on exacerbation of MS. *Ann Neurol* 1994; 36:525–528.
6. Korn-Labetzki I, Khana E, Cooper G, et al. Activity of MS during pregnancy and puerperium. *Ann Neurol* 1984; 16:229–231.
7. Carriere, W, Baskerville J, Ebers, GC. The natural history of MS: A geographically based study. Applications to planning and interpretation of clinical and therapeutic trials. *Brain* 1991;114:1057–1067.
8. Wynn DR, Rodriguez M, O'Fallon M, Kurland LT. A reappraisal of the epidemiology of MS in Olmstead County, Minnesota. *Neurology* 1990;40:780–786.
9. Lublin FD, Reingold SC. Defining the clinical course of multiple sclerosis: Results of an international survey. National Multiple Sclerosis Society (USA) Advisory Committee on Clinical Trials of New Agents in Multiple Sclerosis. *Neurology* 1996;46:907–911.
10. The International Multiple Sclerosis Genetics Consortium. Risk alleles for multiple sclerosis identified by a genomewide study. *N Engl J Med* [Internet]. 2007 [cited 2007, July 29]. Available from: <http://www.nejm.com>.
11. Haegert DG, Marrosu MG. Genetic susceptibility to MS. *Ann Neurol* 1994; 36:2S04–S210.
12. Mumford CJ, Wood NW, Kellar-Wood H, Thorpe JW, Miller DH, Compston, DA. The British Isles survey of MS in twins. *Neurology* 1994; 44:11–15.
13. Sadovnick AD, Armstrong H, Rice GP, et al. A population-based study of MS in twins: Update. *Ann Neurol* 1993; 33:281–285.
14. Prineas JW. Pathology of MS. In: Cook SD, ed. *Handbook of MS*. New York: Marcel Dekker. 1990:187–218.
15. Sobel RA. The pathology of MS. *Neurol Clin* 1995;13:1–21.
16. Waxman SG. Pathophysiology of MS. In: Coe DS, ed. *Handbook of MS*. New York: Marcel Dekker. 1990:219–249.
17. Prineas JW, Barnard RD, Revesz T, Kwon EE, Sharer L, Cho E-S. MS: Pathology of recurrent lesions. *Brain* 1993; 116:681–693.
18. Gonen O, Patalace I, Babb JS, et al. Total brain N-acetylaspartate, a new measure of disease load in MS. *Neurology* 2000;54:15–19.
19. Poser CM, Paty DW, Scheinberg L, et al. New diagnostic criteria for MS: Guidelines for research protocols. *Ann Neurol* 1983; 13:227–231.
20. Krupp LB, Alvarez LA, LaRocca NG, et al. Fatigue in MS. *Arch Neurol* 1988;45:435–437.
21. Freal JE, Kraft GH, Coryell JK. Symptomatic fatigue in MS. *Arch Phys Med Rehabil* 1984; 65:165–168.
22. Thompson AJ, Kennard C, Swash M, et al. Relative efficacy of intravenous methylprednisolone and ACTH in the treatment of acute relapse in MS. *Neurology* 1989;39:696–971.
23. Beck RW, Cleary PA, Anderson MM, et al. A randomized controlled trial of corticosteroids in the treatment of acute optic neuritis. *N Engl J Med* 1992; 326:581–588.
24. Blaiwas JG. Management of bladder dysfunction in MS. *Neurology* 1980;30:12–18.
25. Bradley WE, Logothetis JL, Timm GW. Cystometric and sphincter abnormalities in MS. *Neurology* 1973; 23:1131–1139.
26. Valleroy ML, Kraft G. Sexual dysfunction in MS. *Arch Phys Med Rehabil* 1984;65:125–128.
27. Svendsen, KB, Jensen, TS, Hansen, HJ, Bach, FW. Sensory function and quality of life in patients with multiple sclerosis and pain. *Pain* 2005; 114:473–481.
28. Osterberg, A, Boivie, J, Thuomas, K-A. Central pain in multiple sclerosis—prevalence and clinical characteristics. *Eur J Pain* 2005; 9:531–542.
29. Herholz, K. Cognitive dysfunction and emotional-behavioural changes in MS: The potential of positron emission tomography. *J Neurol Sci* 2006; 245(1–2):9–12.
30. The IFNB MS study group and the University of British Columbia MS/MRI analysis group. Interferon beta-1b in the treatment of MS: Final outcome of the

- randomized controlled trial. *Neurology* 1995; 45:1277–1285.
31. Gold R, Rieckmann P, Chang P, Abdalla J; the PRISMS Study Group. The long-term safety and tolerability of high-dose interferon -1a in relapsing-remitting multiple sclerosis: 4-year data from the PRISMS study. *Eur J Neurol* 2005; 12:649–656.
 32. Johnson KP, Brooks BR, Cohen JA, et al. Copolymer-1 reduces relapse rate and improves disability in relapsing-remitting MS: Results of a phase III multicenter double-blind placebo-controlled trial. *Neurology* 1995; 45:1268–1276.
 33. Thompson AJ, Noseworthy JH. New treatments for MS: Clinical perspective. *Curr Opin Neurol* 1996; 9:187–198.
 34. Fidler JM, DeJoy SQ, Smith FR 3rd, Gibbons JJ Jr. Selective immunomodulation by the antineoplastic agent mitoxantrone. Non-specific adherent suppressor cells derived from mitoxantrone-treated mice. *J Immunol* 1986; 136:2747–2754.
 35. Katz R. Management of spasticity. *Am J Phys Med Rehabil* 1988; 67:108–116.
 36. Nance DW, Sheremata WA, Lynch SG, et al. Relationships of the antispasticity effect of Tizanidine to plasma concentration in patients with MS. *Arch Neurol* 1997; 54:731–736.
 37. Rudick RA, Goodman A, Herndon RM, Panitch HS. Selecting relapsing-remitting MS patients for treatment: The care for early treatment. *J Neuroimmunol* 1999; 98:22–28.
 38. Ponichtera-Mulcare JA, Glaser RM, Mathews T, Camaione, DN. Maximal aerobic exercise in persons with MS. *Clin Kinesiol* 1983; 46(4):12–21.
 39. Ponichtera-Mulcare JA, Mathews T, Glaser RM, Mathews T Gupta SC. Maximal aerobic exercise of individuals with MS using three modes of ergometry. *Clin Kinesiol* 1995; 49:4–13.
 40. White LJ, McCoy SC, Castellano V, Guterrez G, Stevens JE, Walter GA, Vandeborne K. Resistance training improves strength and functional capacity in persons with multiple sclerosis. *Mult Scler* 2004; 10:668–674.
 41. Surakka J, Romberg A, Ruutiainen J, Aunola S, Virtanen A, Karppi SL, Maentaka K. Effects of aerobic and strength exercise on motor fatigue in men and women with multiple sclerosis: A randomized controlled trial. *Clin Rehabil* 2004;18(7):737–746.
 42. Ng AV, Dao HT, Miller RG, Gelina DF, Kent-Braun JA. Blunted pressor and intramuscular metabolic responses to voluntary isometric exercise in MS. *J Appl Physiol* 2000; 88:871–880.
 43. Lambert CP, Archer RL, Evans WJ. Muscle strength and fatigue during isokinetic exercise in individuals with multiple sclerosis. *Med Sci Sport Exerc* 2001; 33(10):1613–1619.
 44. Ponichtera JA, Rodgers MM, Glaser RM, Mathews, T. Concentric and eccentric isokinetic lower extremity strength in persons with MS. *J Orthop Sport Phys Ther* 1988;16(3):114–122.
 45. Armstrong LE, Winant DM, Swasey PR, Seidle ME, Carter AL, Gehlsen GM. Using isokinetic dynamometry to test ambulatory patients with MS. *Phys Ther* 1983;63:1274–1279.
 46. Chen W-Y, Peirson FM, Burnett CN. Force-time measurements of knee muscle function in MS. *Phys Ther* 1987; 67:934–940.
 47. Rice CL, Volmer TL, Bigland-Ritchie B. Neuromuscular responses of patients with MS. *Muscle Nerve* 1992;15:1123–1132.
 48. McDonald WI, Sears TA. Effect of a demyelinating lesion on conduction in the central nervous system studied in single nerve fibers. *J Physiol (London)* 1970; 207:53–54P.
 49. Kent-Braun JA, Sharma KR, Miller RG, Weiner MW. Postexercise phosphocreatine resynthesis is slowed in multiple sclerosis. *Muscle Nerve* 1994; 17(8):835–841.
 50. Kent-Braun JA, Sharma KR, Weiner MW, Miller RG. Effects of exercise on muscle activation and metabolism in MS. *Muscle Nerve* 1994; 17(10):1162–1169.
 51. Sharma KR, Kent-Braun J, Mynhier MA, Weiner MW, Miller RG. Evidence of an abnormal intramuscular component of fatigue in MS. *Muscle Nerve* 1995;18(12):1403–1411.
 52. Mulcare JA, Webb P, Mathews T, Gupta, SC. Sweat response in persons with multiple sclerosis during submaximal aerobic exercise. *International Journal of MS Care* 2001;3(4):26–33.
 53. Tantucci, C Massucci M, Piperno R, Grassi V, Sorbini CA. Energy cost of exercise in MS patients with low degree of disability. *Mult Scler* 1996; 2(3):161–167.
 54. Vaz Fragoso C, Wirz D, Mashman J. Establishing a physiological basis to multiple sclerosis-related fatigue: A case report. *Arch Phys Med Rehabil* 1995; 76:583–586.
 55. Linden D, Diehl RR, Kretzschmar A, Berlit P. Autonomic evaluation by means of standard tests and power spectral analysis in multiple sclerosis. *Muscle Nerve* 1997; 20(7):809–814.
 56. Flachenecker P, Wolf A, Krauser M, Hartung HP, Reiners K. Cardiovascular autonomic dysfunction in multiple sclerosis: Correlation with orthostatic intolerance. *J Neurol* 1999; 246(7):578–586.
 57. Bonnett M, Mulcare J, Mathews T, Gupta SA, Ahmed N, Yeragani V. Heart rate and QT interval variability in multiple sclerosis: Evidence for decreased sympathetic activity. *J Neurol Sci [Turkish]* 2006; 23(4):248–256.
 58. Pepin EB, Hicks RW, Spencer MK, Tan ZC, Jackson CGR. Pressor response to isometric exercise in patients with multiple sclerosis. *Med Sci Sports Exerc* 1996;23:656–660.
 59. Noble BJ, Robertson RJ. *Perceived Exertion*. Champaign, IL: Human Kinetics; 1996.
 60. Pariser G, Madras D, Weiss E. J Outcomes of an aquatic exercise program including aerobic capacity, lactate threshold, and fatigue in two individuals with multiple sclerosis. *Neur Phys Ther* 2006; 30(2):82–90.
 61. Nieman DC. *Fitness and Sports Medicine: An Introduction*. Palo Alto, CA: Bull Publishing Company. 1990:500.
 62. Petajan JH, Gappmaier E, White AT, Spencer JK, Mino L, Hicks RW. Impact of aerobic training on fitness and quality of life in MS. *Ann Neurol* 1996;34:432–441.
 63. Ponichtera-Mulcare JA, Mathews T, Barrett PJ, Gupta SC. Change in aerobic fitness of patients with MS during 6-month training program. *Sports Medicine, Training, and Rehabilitation* 1997;7:265–272.
 64. Gutierrez GM, Chow JW, Tillman MD, McCoy SC, Castellano V, White J. Resistance training improves gait kinematics in persons with multiple sclerosis. *Arch Phys Med Rehabil* 2005;86:1824–1929.
 65. White LJ, McCoy SC, Castellano V, Guterrez G, Stevens JE, Walter GA, Vandeborne K. Resistance training improves strength and functional capacity in persons with multiple sclerosis. *Mult Scler* 2004;10:668–674
 66. Taylor NF, Dodd KJ, Prasad D, Denisenko S. Progressive resistance exercise for people with multiple sclerosis. *Disabil Rehabil* 2006; 28(18):1119–1126.
 67. American College of Sports Medicine. Progression models in resistance training for healthy adults. *Med Sci Sports Exerc* 2002;34(2): 364–380.
 68. van den Berg M, Dawes H, Wade DT, Newman M, Burridge J, Izadi H, Sackley CM. Treadmill training for individuals with multiple sclerosis: A pilot randomised trial. *J Neurol Neurosurg Psychiatry* 2006;77:531–533.
 69. Hoogvorst ELJ, Eikelenboom MJ, Uitdehaag BMJ, Polman CH. One year changes in disability in multiple sclerosis: Neurological examination compared with patient self report. *J Neurol Neurosurg Psych* 2003;74(4):439–442.
 70. Mostert S, Kesselring J. Effect of a short-term exercise training program on aerobic fitness, fatigue, health perception and activity level of subjects with multiple sclerosis. *Mult Scler* 2002;8:61–168.
 71. Romberg A, Virtanen A, Ruutiainen J. Long-term exercise improves functional impairment but not quality of life in multiple sclerosis. *J Neurol* 2005;252:839–845.
 72. Kileff J, Ashburn A. A pilot study of the effect of aerobic exercise on people with moderate disability multiple sclerosis. *Clin Rehabil* 2005; 19:165–169.
 73. Newman MA, Dawes H, van den Berg M, Wade DT, Burridge J, Izadi H. Can aerobic treadmill training reduce the effort of walking and Fatigue in people with multiple sclerosis: A pilot study. *Mult Scler* 2007;13:113–119.
 74. Giesser B, Beres-Jones J, Budovitch A, Herlihy E, Harkema S. Locomotor training using body weight support on a treadmill improves mobility in persons with multiple sclerosis: A pilot study. *Mult Scler*2007;13:224–231.

75. Berg KO, Wood-Dauphine SL, Williams JI, Maki B. Measuring balance in the elderly: Validation of an instrument. *Can J Public Health* 1992;83[Suppl 2]:S7–S11.
76. Jackson KJ, Mulcare JA, Donahoe-Fillmore B, Fritz HI, Rodgers MM. Home balance training intervention for people with multiple sclerosis. *Int J MS Care* 2007; 9:111–117.
77. Gutierrez GM, Chow JW, Tillman MD, McCoy SC, Castellano V, White LJ. Resistance training improves gait kinematics in persons with multiple sclerosis. *Arch Phys Med Rehabil* 2005;86:1824–1829.
78. White LJ, McCoy SC, Castellano V, Gutierrez G, Stevens JE, Walter GA, Vandenborne K. Resistance training improves strength and functional capacity in persons with multiple sclerosis. *Mult Scler* 2004; 10:668–674.
79. Stuijbergen A. Health promoting behaviors and quality of life among individuals with MS. *Scholarly Inquiry for Nursing Practice* 1995; 9:31–50.
80. Stuijbergen A, Becker H. Predictors of health promoting lifestyles in persons with disabilities. *Res Nurs Health* 1994; 17:3–13.
81. Ng AV, Kent-Braun J. Quantification of lower physical activity in persons with MS. *Med Sci Sports Exerc* 1997; 29(4):517–523.
82. Rimmer JH, Rubin SS, Braddock D. Barriers to exercise in African American women with physical disabilities. *Arch Phys Med Rehabil* 2000; 81(2):182–188.
83. Abramson S, Stein J, Schaufele M, et al. Personal exercise habits and counseling practices of primary care physicians: A national survey. *Clin J Sport Med* 2000; 10(1):40–48.
84. Mulcare JA, Webb P, Mathew T, Gupta SC. The effect of body cooling on the aerobic endurance of persons with MS following a 3-month aerobic training program. *Med Sci Sports Exerc* 1997; 29(5):S83.
85. White AT, Wilson TE, Petajan JH. Effect of pre-exercise cooling on physical function and fatigue in MS patients. *Med Sci Sports Exerc* 1997; 29(5):S83.

Chapter 4

1. Aarsland D, Larsen JP, Tandberg E, Laake K. Predictors of nursing home placement in Parkinson's disease: A population-based, prospective study. *J Am Geriatr Soc* 2000; 48:938–942.
2. Abe K, Takanashi M, Yanagihara T. Fatigue in patients with Parkinson's disease. *Behav Neurol* 2000; 12(3):103–106.
3. Adler CH. Nonmotor complications in Parkinson's disease. *Mov Disord* 2005;20(suppl 11):23–29.
4. Alves G, Larsen JP, Emre M, et al. Changes in motor subtype and risk for incident dementia in Parkinson's disease. *Mov Disord* 2006; 21(8):1123–1130.
5. American College of Sports Medicine. *ACSM's Guidelines for Exercise Testing and Prescription*. 8th ed. Baltimore: Lippincott Williams & Wilkins; 2009.
6. Arle JE, Alterman RL. Surgical options in Parkinson's disease. *Med Clin North Am*. 1999;83(2):483–498.
7. Banks MA, Caird FI. Physiotherapy benefits patients with Parkinson's disease. *Clin Rehabil* 1989;3:11–16.
8. Berg K, Wood-Dauphinee S, Williams JI, et al. Measuring balance in the elderly: Preliminary development of an instrument. *Physiother Can* 1989;41(6):304–311.
9. Berg MJ, Ebert B, Willis DK, et al. Parkinsonism—drug treatment: Part I. *Drug Intell Clin Pharm* 1987;21(1):10–21.
10. Biglan KM, Ravina B. Neuroprotection in Parkinson's disease: An elusive goal. *Semin Neurol* 2007;27:106–112.
11. Bindoff LA, Birch-Machin MA, Carlidge NEF, et al. Mitochondrial function in Parkinson's disease. *Lancet* 1989;2:49.
12. Blin O, Desnuelle C, Rascol O, et al. Mitochondrial respiratory failure in skeletal muscle from patients with Parkinson's disease and multiple system atrophy. *J Neurol Sci* 1994;125:95–101.
13. Bohnen NI, Kaufer DI, Hendrickson R, Constantine GM, Mathis CA, Moore RY. Cortical cholinergic denervation is associated with depressive symptoms in Parkinson's disease and parkinsonian dementia. *J Neurol Neurosurg Psychiatry* 2007;78:641–643.
14. Burn DJ, Rowan EN, Allan LM, Molloy S O'Brien JT, McKeith IG. Motor subtype and cognitive decline in Parkinson's disease, Parkinson's disease with dementia, and dementia with Lew bodies. *J Neurol Neurosurg Psychiatry* 2006;77:585–589.
15. Caird FI. Parkinson's disease and its natural history. In: Caird FI, editor. *Rehabilitation In Parkinson's Disease*. New York: Chapman & Hall; 1991. p. 1–7.
16. Cakit BD, Saracoglu M, Genc H, Erdem HR. The effects of incremental speed-dependent treadmill training on postural instability and fear of falling in Parkinson's disease. *Clin Rehabil* 2007; 21:698–705.
17. Canning CG, Alison JA, Allen NE, et al. Parkinson's disease: An investigation of exercise capacity, respiratory function, and gait. *Arch Phys Med Rehabil* 1997; 78:199–207.
18. Comella CL, Stebbins GT, Brown-Toms N, et al. Physical therapy and Parkinson's disease: A controlled clinical trial. *Neurology* 1994; 44:376–378.
19. Comella C. Sleep disturbances in Parkinson's disease. *Curr Neurol Neurosci Rep* 2003;3:173–180.
20. Corcos DM, Chen CM, Quinn NP, et al. Strength in Parkinson's disease: Relationship to rate of force generation and clinical status. *Ann Neurol* 1996;39:79–88.
21. Diamond SG, Markham ChC. Present mortality in Parkinson's disease: The ratio observed to expected deaths with a method to calculate expected deaths. *J Neural Transm* 1976;36:259–269.
22. Dibble LE, Hale T, Marcus RL, Gerber JP, LaStayo PC. The safety and feasibility of high-force eccentric resistance exercise in persons with Parkinson's disease. *Arch Phys Med Rehabil* 2006;87:1280–1282.
23. Dietz V, Colombo G. Influence of body load on the gait pattern in Parkinson's disease. *Mov Disord* 1998;13(2):255–261.
24. Duncan PW, Weiner DK, Chandler J, et al. Functional reach: A new clinical measure of balance. *J Gerontol* 1990;45:M192–197.
25. Duvoisin RC, Eldridge R, Williams A, et al. Twin study of Parkinson's disease. *Neurology*. 1981; 31:77–80.
26. Duvoisin RC. Genetics of Parkinson's disease. *Adv Neurol* 1986; 45:307–312.
27. Ebadi M, Srinivasan SK, Baxi MD. Oxidative stress and antioxidant therapy in Parkinson's disease. *Prog Neurobiol* 1996;48:1–19.
28. Eckert T, Tang C, Eidelberg D. Assessment of the progression of Parkinson's disease: A metabolic network approach. *Lancet Neurol* 2007;6:926–932.
29. Ellis T, de Goede CJ, Feldman RG, Wolters EC, Kwakkel G, Wagenaar RC. Efficacy of a physical therapy program in patients with Parkinson's disease: A randomized controlled trial. *Arch Phys Med Rehabil* 2005;86:626–632.
30. Erdal KJ. Depressive symptom patterns in patients with Parkinson's disease and other older adults. *J Clin Psychol* 2001;57(12):1559–1569.
31. Factor SA, Feustel PJ, Friedman JH, et al. Logitudinal outcome of Parkinson's disease patients with psychosis. *Neurology* 2003; 60(11): 17546–1761.
32. Fahn S, Elton RL, Members of the UPDRS Development Committee. Unified Parkinson's disease rating scale. In: Stanley F, Marsden CD, Goldstein M, et al., editors. *Recent Developments in Parkinson's disease, volume II*. Florham Park (NJ): MacMillan Healthcare Information; 1987. p. 153–303.
33. Farley B, Koshland GF. Training BIG to move faster: The application of the speed-amplitude relation as a rehabilitation strategy for people with Parkinson's disease. *Exp Brain Res* 2005;167:462–467.
34. Findley L, Aujla M, Bain PG, et al. Direct economic impact of Parkinson's disease: A research survey in the United Kingdom. *Mov Disord* 2003;18(10):1139–1145.
35. Flewitt B, Capildeo R, Rose FC. Physiotherapy and assessment in Parkinson's disease using the polarised lightgoniometer. In: Rose FC, Capildeo R, editors. *Research Progress in Parkinson's disease*. Kent: Pitman Medical; 1981. p. 404–413.
36. Frenkel-Toledo S, Giladi N, Peretz C, Herman T, Gruendlinger L, Hausdorff JM. Treadmill walking as an external pacemaker to improve gait rhythm and stability in Parkinson's disease. *Mov Disord* 2005;20:1109–1114.
37. Friedman J, Friedman H. Fatigue in Parkinson's disease. *Neurology* 1993;43(10):2016–2018.
38. Friedman JH, Friedman H. Fatigue in Parkinson's disease: A nine-year follow-up. *Mov Disord* 2001;16(6):1120–1122.
39. Gershanik OS, Nygaard TG. Parkinson's disease beginning before age 40. In: Streifler MB, Korczyn AD, Melamed E, et al., editors. *Advances In Neurology: Parkinson's Disease: Anatomy, Pathology, and Therapy*. New York: Raven Press, Ltd.; 1990; p. 251–258.
40. Gerstenbrand F, Poewe WH. The classification of Parkinson's disease. In: Stern G, editor. *Parkinson's Disease*. Baltimore: John Hopkins University Press; 1990. p. 315–331.
41. Gibberd FB, Page NGR, Spencer KM, et al. A controlled trial of physiotherapy for Parkinson's disease. In: Rose FC, Capildeo R, editors. *Research Progress in*

- Parkinson's disease. Kent: Pitman Medical; 1981.p. 401–403.
42. Gupta M. Parkinson's disease pathology: Multineurotransmitter systems defects. In: Schneider JS, Gupta M, editors. *Curren Concepts in Parkinson's disease Research*. Seattle: Hogufe & Haber Publishers; 1993. p. 21–40.
 43. Hardy J, Cai H, Cookson MR, Gwing-Hardy K, Aingleton A. Genetics of Parkinson's disease and parkinsonism. *Ann Neurol* 2006; 60:389–398.
 44. Hauser RA. Levodopa/carbidopa/entacapone (stalevo). *Neurology* 2004;62(Suppl 1):S64–S71.
 45. Hennekens CH, Buring JE. *Epidemiology in Medicine*. Boston: Little, Brown and Co.; 1987. p. 57.
 46. Herlofson K, Larsen JP. Measuring fatigue in patients with Parkinson's Disease—the Fatigue Severity Scale. *Eur J Neurol* 2002;9(6):595–600.
 47. Hirsch EC. Does oxidative stress participate in nerve cell death in Parkinson's disease? *Eur Neurol* 1993;33(Suppl 1):52–59.
 48. Hirsch MA, Toole T, Maitland CG, Rider RA. The effects of balance training and high-intensity resistance training on persons with idiopathic Parkinson's disease. *Arch Phys Med Rehabil* 2003; 84:1109–1117.
 49. Hobson P, Meara J. Risk and incidence of dementia in a cohort of older subjects with Parkinson's disease in the United Kingdom. *Mov Disord* 2004; 19(9):1043–1049.
 50. Hoehn M, Yahr MD. Parkinsonism: Onset, progression, and mortality. *Neurology* 1967; 17(5):427–442.
 51. Honey C, Gross RE, Lozano AM. New developments in the surgery for Parkinson's disease. *Can J Neurol Sci* 1999;26(Suppl 2):S45–S52.
 52. Hurwitz LJ. Improving mobility in severely disabled parkinsonian patients. *Lancet* 1964:953–955.
 53. Jahn K, Deutschlander A, Stephan T, Kalla R, Wiesmann M, Strupp M, Brandt T. Imaging human suprsprinal locomotor centers in brainstem and cerebellum. *NeuroImage* 2008;39:786–792.
 54. Jankovic J. Parkinsonism—plus syndromes. *Mov Disord* 1989; 4(suppl 1):S95–S119.
 55. Janvin CC, Larsen JP, Aarsland D, et al. Subtypes of mild cognitive imparment in Parkinson's disease: Progression to dementia. *Mov Disord* 2006;21(9):1343–1349.
 56. Jenner P, Olanow CW. Oxidative stress and the pathogenesis of Parkinson's disease. *Neurology* 1996;47(suppl 3):S161–S170.
 57. Kakinuma S, Hiroshi N, Pramanik B, et al. Muscle weakness in Parkinson's disease: Isokenetic study of the lower limbs. *Eur Neurol* 1998;39:218–222.
 58. Karlsen K, Larsen JP, Tandberg E, Jorgensen K. Fatigue in patients with Parkinson's disease. *Mov Disord* 1999;14(2):237–241.
 59. Kenney C, Jankovic J. Rotigotine transdermal patch in the treatment of Parkinson's disease and restless legss syndrom. *Expert Opin Pharmacother* 2007;8(9):1329–1335.
 60. Kessler II. Epidemiologic studies of Parkinson's disease II. A hospital based survey. *Am J Epidemiol* 1972;95(4):308–318.
 61. Kessler II. Epidemiologic studies of Parkinson's disease III. A community-based survey. *Am J Epidemiol* 1972;96(4):242–254.
 62. Kessler II, Diamond EL. Epidemiologic studies of Parkinson's disease I. Smoking and Parkinson's disease: A survey and explanatory hypothesis. *Am J Epidemiol* 1972;94(1):16–25.
 63. Keus SHJ, Bloem BR, Hendriks EJM, Bredero-Cohen AB, Munneke M. Evidence-based analysis of physical therapy in Parkinson's disease with recommendation for practice and research. *Mov Disord* 2007;22:451–460.
 64. Koller WC, Hubble JP. Classification of Parkinsonism. In: Koller WC, editor. *Handbook of Parkinson's disease*. 2nd ed. New York: Marcel Dekker, Inc.; 1992. p. 59–103.
 65. Kondo K, Kurland LT, Schull WJ. Parkinson's disease, genetic analysis and evidence of a multifactorial etiology. *Mayo Clin Proc* 1973;48:465–475.
 66. Koseoglu F, Inan L, Ozel S, et al. The effects of a pulmonary rehabilitation program on pulmonary function tests and exercise tolerance in patients with Parkinson's disease. *Funct Neurol* 1997;12: 319–325.
 67. Leentjens AF, Van den Akker M, Metsmakers JF, Lousberg R, Verhey FR. Higher incidence of depression preceding onset of Parkinson's disease: A register study. *Mov Disord* 2003;18(4):414–418.
 68. Lew MR, Pahwa R, Leehey M, Bertoni J, Kricorian G, the Sydis Selegiline Study Group. Safety and efficacy of newly formulated selegiline orally disintegrating tablets as an adjunct to levodopa in the management of “off” episodes in patients with Parkinson's disease. *Curr Med Res Opin* 2007;23(4):741–750.
 69. LeWitt PA, Bharucha A, Chitrit I, et al. Perceived exertion and muscle efficiency in Parkinson's disease: L-dopa effects. *Clin Neuropharmacol* 1994;17(5):454–459.
 70. LeWitt PA, Lyons KE, Pahwa R, SP 650 Study Group. Advanced Parkson disease treated with rotigotine transdermal system: PREFER study. *Neurology* 2007;68:1262–1267.
 71. Lim I, van Wegen E, de Goede C, et al. Effects of external rhythmical cueing on gait in patients with Parkinson's disease: A systematic review. *Clin Rehabil* 2005;19:695–713.
 72. Liston RAL, Brouwer BJ. Reliability and validity of measures obtained from stroke patient using the balance master. *Arch Phys Med Rehabil* 1996;77:425–430.
 73. Lou JS, Kearns G, Oken B, Sexton G, Nutt J. Exacerbated physical fatigue and mental fatigue in Parkinson's disease. *Mov Disord* 2001;16(2):190–196.
 74. Ludin SM, Steiger MJ, Ludin. Autonomic disturbances and cardio- vascular reflexes in idiopathic Parkinson's disease. *J Neurol* 1987; 235:10–15.
 75. Maguire GH. Occupational therapy. In: Abrams WB, Berkow R, Fletcher AJ, et al., editors. *The Merck Manual of Geriatrics*. Rathway (NJ): Merck Sharp and Dohme Research Laboratories; 1990. p. 274–278.
 76. Maria B, Sophia S, Michalis M, et al. Sleep breathing disorders in patients with idopathic Parkinson's disease. *Respir Med* 2003;97(10):1151–1157.
 77. Marras C, Rochon P, Lang AE. Predicting motor decline and disability in Parkinson disease: A systematic review. *Arch Neurol* 2002;59:1724–1728.
 78. Marsden CD. Parkinson's disease in twins. *J Neurol Neurosurg Psychiatry* 1987;50:105–106.
 79. Martignoni E, Micieli G, Cavallini A, et al. Autonomic disorders in idiopathic Parkinsonism. *J Neural Transm* 1986; 22(suppl):149–61.
 80. Marttila RJ, Rinne UK. Epidemiology of Parkinson's disease in Finland. *Acta Neurol Scand* 1976;53:80–102.
 81. Marttila RJ, Rinne UK, Siirtola T, et al. Mortality of patients with Parkinson's disease treated with levodopa. *J Neurol* 1977; 216:147–153.
 82. Marttila RJ, Rinne UK. Epidemiology of Parkinson's disease—an overview. *J Neural Trans* 1981;51:135–148.
 83. Marttila RJ, Hennekens CH, Buring JE. *Epidemiology in Medicine*. Boston: Little, Brown and Co.; 1987. p. 57.
 84. Martilla RJ, Kaprio J, Kistenvuo MD, et al. Parkinson's disease in a nationwide twin cohort. *Neurology* 1988;38:1217– 1219.
 85. Marttila RJ. Epidemiology. In: Koller WC, editor. *Handbook of Parkinson's disease*. 2nd ed. New York: Marcel Dekker, Inc.; 1992. p. 35–57.
 86. Montgomery EB, Lipsy RJ. Treatment of Parkinson's disease. In: Bressler R, Kane MD, editors. *Geriatric Pharmacology*. New York: McGraw Hill, Inc.; 1993. p. 309–328.
 87. Morizane A, Li JY, Brundin P. From bench to bed: The potential of stem cells for the treatment of Parkinson's disease. *Cell Tissue Res* 2008;331:323–336.
 88. Nagayama H, Hamamoto M, Ueda M, Nagashima J, Katayama Y. Reliability of MIBG myocardial scintigraphy in the diagnosis of Parkinson's disease. *J Neurol Neurosurg Psychiatry* 2005;76:249–251.
 89. Nogaki H, Kakinuma S, Morimatsu M. Movement velocity dependent muscle strength in Parkinson's disease. *Acta Neurol Scand* 1999; 99:152–157.
 90. Olanow CW. An introduction to the free radical hypothesis in Parkinson's disease. *Ann Neurol* 1992; 32(Suppl):S2–S9.
 91. O'Sullivan SB. Parkinsons disease. In: O'Sullivan SB, Schmitz TJ, editors. *Physical Rehabilitation*. 5th ed. Philadelphia: FA Davis Company; 2007. p. 853–893.

Chapter 5

1. Nelson MR, Rott EJ. Spina bifida. In: Grabois M, Garrison SJ, Hart KA, et al., eds. *Physical Medicine and Rehabilitation: The Complete Approach*. Malden, MA: Blackwell Science; 2000:1414–1432.
2. Cardenas DD, Burns SP, Chan L. Rehabilitation of spinal cord injury. In: Grabois M, Garrison SJ, Hart KA, et al, eds. *Physical Medicine and Rehabilitation: The Complete Approach*. Malden, MA:Blackwell Science; 2000:1305–1324.
3. Yarkony GM, Chen D. Rehabilitation of patients with spinal cord injuries. In: Braddom RJ, ed. *Physical Medicine and Rehabilitation*. Philadelphia: W.B. Saunders; 1996:1149–1179.
4. Mallory B. Autonomic dysfunction in spinal cord disease. In: Lin VW, ed. *Spinal Cord Medicine: Principles and Practice*. New York: Demos; 2003:477–500.
5. Goshgarian HG. Anatomy and function of the spinal cord. In: Lin VW, ed. *Spinal Cord Medicine: Principles and Practice*. New York: Demos; 2003:15–34.
6. ASIA. International Standards for Neurological Classification of SCI. Atlanta: American Spinal Injury Association; 2002.
7. ASIA. Standard neurological classification of spinal cord injury. Atlanta: American Spinal Injury Association; 2006. (accessed online, 2-17-08: http://www.asia-spinalinjury.org/publications/2006_Classif_worksheet.pdf)
8. Consortium for Spinal Cord Medicine. Outcomes following traumatic spinal cord injury: Clinical practice guidelines for healthcare professions. Washington, DC: Paralyzed Veterans of America; 1999.
9. Furrher MJ. Rehabilitation and research training center in community-oriented services for persons with spinal cord injury: A progress report. Houston, TX: The Baylor College of Medicine and The Institute for Rehabilitation Research; 1991.
10. Whitneck GG. Learning from recent empirical investigations. In: Whitneck GG, Charlifue SW, Gerhart KS, et al., eds. *Aging with Spinal Cord Injury*. New York: Demos Publications; 1993.
11. Anson CA, Shepherd C. Incidence of secondary complications in spinal cord injury. *Int J Rehabil Res* 1996;19:55–66.
12. Johnson RL, Gerhart KA, McCray J, et al. Secondary conditions following spinal cord injury in a population-based sample. *Spinal Cord* 1998;36:45–50.
13. Bauman WA, Spungen AM. Metabolic changes in persons after spinal cord injury. *Phys Med Rehabil Clin N Am* 2000;11:109–140.
14. Washburn RA, Figoni SF. Physical activity and chronic cardiovascular disease prevention: A comprehensive literature review. *Top Spinal Cord Injury Rehabil* 1998;3:16–32.
15. Davis GM. Exercise capacity of individuals with paraplegia. *Med Sci Sports Exerc* 1993;25:423–432.
16. Figoni SF. Exercise responses and quadriplegia. *Med Sci Sports Exerc* 1993;25:433–441.
17. Price MJ. Thermoregulation during exercise in individuals with spinal cord injuries. *Sports Med* 2006;36:863–879.
18. Hopman MTE, Nommensen E, van Asten WNJC, et al. Properties of the venous vascular system in the lower extremities on individuals with paraplegia. *Paraplegia* 1994;32:810–816.
19. Bauman WA. Endocrinology and metabolism after spinal cord injury. In: Kirshblum S, Campagnolo DI, DeLisa JA, eds. *Spinal Cord Medicine*. Philadelphia, PA: Lippincott Williams & Wilkins; 2002:164–180.
20. Kelly LE. Spinal cord disabilities. In: Winnick JP, ed. *Adapted Physical Education and Sport*. 4th ed. Champaign, IL: Human Kinetics; 2005:275–306.
21. Hart AL, Malone TR, English T. Shoulder function and rehabilitation implications for the wheelchair athlete. *Top Spinal Cord Injury Rehabil* 1998;3:50–65.
22. Winnick JP, Short FX. The physical fitness of youngsters with spinal neuromuscular conditions. *Adapted Phys Activ Q* 1984;1:37–51.
23. Shephard RJ. *Fitness in Special Populations*. Champaign, IL: Human Kinetics, 1990.
24. Winnick JP, Short FX. *The Brockport Physical Fitness Test Manual*. Champaign, IL: Human Kinetics; 1999.
25. Andrade CK, Kramer J, Garber M, et al. Changes in self-concept, cardiovascular endurance and muscular strength of children with spina bifida aged 8 to 13 years in response to a 10-week physical-activity programme: A pilot study. *Child Care Health Dev* 1991;17:183–196.
26. Morrison SA, Melton-Rogers SL, Hooker SP. Changes in physical capacity and physical strain in persons with acute spinal cord injury. *Top Spinal Cord Injury Rehabil* 1997;3:1–15.
27. Hjeltnes N, Jansen T. Physical endurance capacity, functional status and medical complications in spinal cord injured subjects with long-standing lesions. *Paraplegia* 1990;28:428–432.
28. Stewart MW, Melton-Rogers SL, Morrison S, et al. The measurement properties of fitness measures and health status for person with spinal cord injuries. *Arch Phys Med Rehabil* 2000;81:394–400.
29. Raymond J, Davis GM, Fahey A, et al. Oxygen uptake and heart rate responses during arm vs combined arm/electrically stimulated leg exercise in people with paraplegia. *Spinal Cord* 1997;35:680–685.
30. Haisma JA, van der Woude LH, Stam HJ, Bergen MP, Sluis TA, Bussmann JB. Physical capacity in wheelchair-dependent persons with a spinal cord injury: A critical review of the literature. *Spinal Cord* 2006;44(11):642–652.
31. Glaser RM. Exercise and locomotion for the spinal cord injured. *Exerc Sports Sci Rev* 1985;13:263–303.
32. Dreisinger TE, Londeree BR. Wheelchair exercise: A review. *Paraplegia* 1982;20:20–34.
33. Fuhr L, Langbein E, Edwards LC, et al. Diagnostic wheelchair exercise testing. *Top Spinal Cord Injury Rehabil* 1997;3:34–48.
34. Pitetti KH, Barrett PJ, Campbell KD, et al. The effect of lower body positive pressure on the exercise capacity of individuals with spinal cord injury. *Med Sci Sports Exerc* 1994;26:463–468.
35. Sawka MN, Latzka WA, Pandolf KB. Temperature regulation during upper body exercise: Able-bodied and spinal cord injured. *Med Sci Sports Exerc* 1989;21:S132–S140.
36. Houtman S, Oeseburg B, Hopman MTE. Blood volume and hemoglobin after spinal cord injury. *Am J Phys Med Rehabil* 2000;29:260–265.
37. Veeger HEJ, Hajd Yahmed M, van der Woude LHV, et al. Peak oxygen uptake and maximal power output of Olympic wheelchair-dependent athletes. *Med Sci Sports Exerc* 1991;23:1201–1209.
38. Coutts KD, Rhodes EC, McKenzie DC. Maximal exercise responses of tetraplegics and paraplegics. *J Appl Physiol* 1971;31:479–482.
39. Gass GC, Camp EM. Physiological characteristics of training Australian paraplegic and tetraplegic subjects. *Med Sci Sports Exerc* 1979;11:256–265.
40. Van Loan MD, McCluer S, Loftin JM, et al. Comparison of physiological responses to maximal arm exercise among able-bodied, paraplegics, and quadriplegics. *Paraplegia* 1987;25:397–405.
41. Figoni SF, Boileau RA, Massey BH, et al. Physiological responses of quadriplegic and able-bodied men during exercise at the same VO₂. *Adapted Phys Activ Q* 1988;5:130–139.
42. Hjeltnes N. Capacity for physical work and training after spinal injuries and strokes. *Scand J Rehabil Med* 1982;29:245–251.
43. Hjeltnes N. Control of medical rehabilitation of para- and tetraplegics by repeated evaluation of endurance capacity. *Int J Sports Med* 1984;5:171–174.
44. Figoni SF, Glaser RM. Arm and leg exercise stress testing in a person with quadriplegia. *Clin Kinesiol* 1993;47:25–36.
45. Claydon VE, Hol AT, Eng JJ, Krassioukov AV. Cardiovascular responses and postexercise hypotension after arm cycling exercise in subjects with spinal cord injury. *Arch Phys Med Rehabil* 2006; 87(8):1106–1114.
46. Figoni SF, Gupta SC, Glaser RM. Effects of posture on arm exercise performance of adults with tetraplegia. *Clin Exerc Physiol* 1999;1:74–85.
47. Hopman MTE, Dueck C, Monroe M, et al. Limits of maximal performance in individuals with spinal cord injury. *Int J Sports Med* 1998;19:98–103.
48. Ready AE. Responses of quadriplegic athletes to maximal and submaximal exercise. *Physiother Can* 1984;36:124–128.
49. Hooker SP, Figoni SF, Rodgers MM, et al. Metabolic and hemodynamic responses to concurrent voluntary arm crank and electrical stimulation leg cycle exercise in quadriplegics. *J Rehabil Res Dev* 1992;29:1–11.
50. Takahashi M, Sakaguchi A, Matsukawa K, Komine H, Kawaguchi K, Onari K. Cardiovascular control during voluntary static exercise in humans with tetraplegia. *J Appl Physiol* 2004;97:2077–82.
51. Krebs P, Eickelberg W, Krobath H, et al. Effects of physical exercise on peripheral vision and learning in children with spina bifida manifestation. *Perceptual*

- and Motor Skills 1989;68:167–174.
52. Agre JC, Findley TW, McNally MC, et al. Physical activity capacity in children with myelomeningocele. *Arch Phys Med Rehabil* 1987;68:372–377.
53. Janssen TWJ, van Oers CAJM, van der Woude LHV, et al. Physical strain in daily life of wheelchair users with spinal cord injuries. *Med Sci Sports Exerc* 1994;26:661–670.
54. Dallmeijer AJ, Hopman MTE, van As HHJ, et al. Physical capacity and physical strain in persons with tetraplegia: The role of sport activity. *Spinal Cord* 1996; 34:729–735.
55. Davis GM, Shephard RJ, Leenen FH. Cardiac effects of short term arm crank training in paraplegics: Echocardiographic evidence. *Eur J Appl Physiol Occup Physiol* 1987;56:90–96.
56. Devillard X, Rimaud D, Roche F, Calmels P. Effects of training programs for spinal cord injury. *Ann Readapt Med Phys* 2007;50(6): 490–498.
57. Valent L, Dallmeijer A, Houdijk H, Talsma E, van der Woude L. The effects of upper body exercise on the physical capacity of people with a spinal cord injury: A systematic review. *Clin Rehabil* 2007 Apr;21(4):315–330.
58. Knutsson E, Lewenhaupt-Olsson E, Thorsen M. Physical work capacity and physical conditioning in paraplegic patients. *Paraplegia* 1973;11:205–216.
59. Nilsson S, Staff PH, Pruett ED. Physical work capacity and the effect of training on subjects with long-standing paraplegia. *Scand J Rehabil Med* 1975;7:51–56.
60. Davis GM, Shephard RJ. Strength training for wheelchair users. *Br J Sports Med* 1990;24:25–30.
61. McLean KP, Skinner JS. Effect of body training position on outcomes of an aerobic training study on individuals with quadriplegia. *Arch Phys Med Rehabil* 1995;76:139–150.
62. Lassau-Wray ER, Ward GR. Varying physiological response to arm-crank exercise in specific spinal injuries. *J Physiol Anthropol Appl Human Sci* 2000;19:5–12.
63. Hooker SP, Figoni SF, Rodgers MM, et al. Physiologic effects of electrical stimulation leg cycle exercise training in spinal cord injured persons. *Arch Phys Med Rehabil* 1992;73:470–476.
64. Nash MS, Bilsker S, Marcillo AE, et al. Reversal of adaptive left ventricular atrophy following electrically stimulated exercise training in human tetraplegia. *Paraplegia* 1992;29:590–599.
65. Nash MS. Exercise reconditioning of the heart and peripheral circulation after spinal cord injury. *Top Spinal Cord Injury Rehabil* 1998;4:1–15.
66. Faghri PD, Glaser RM, Figoni SF. Functional electrical stimulation leg cycle ergometer exercise: Training effect on cardiorespiratory responses of spinal cord injured subjects at rest and during submaximal exercise. *Arch Phys Med Rehabil* 1992;73:1085–1093.
67. Janssen TWJ, Glaser RM, Shuster DB. Clinical efficacy of electrical stimulation exercise training: Effects on health, fitness, and function. *Top Spinal Cord Injury Rehabil* 1998;3:33–49.
68. Figoni SF, Glaser RM, Collins SR. Training effects of hybrid exercise on peak physiologic responses in quadriplegics. *Proceedings of the 12th International Congress of World Confederation for Physical Therapy*. Alexandria, VA: American Physical Therapy Association, 1995, Paper no. PO-SI-0028T on CD-ROM.
69. Thijssen DH, Ellenkamp R, Smits P, Hopman MT. Rapid vascular adaptations to training and detraining in persons with spinal cord injury. *Arch Phys Med Rehabil* 2006;87(4):474–481.
70. Cooney MM, Walker JB. Hydraulic resistance exercise benefits cardiovascular fitness of spinal cord injured. *Med Sci Sports Exerc* 1986;18:522–525.
71. Jacobs PL, Nash MS, Rusinkowski JW Jr. Circuit training provides cardiorespiratory and strength benefits in persons with paraplegia. *Med Sci Sports Exerc* 2001;33:711–717.
72. Nash MS, van de Ven I, van Elk N, Johnson BM. Effects of circuit resistance training on fitness attributes and upper-extremity pain in middle-aged men with paraplegia. *Arch Phys Med Rehabil* 2007;88(1):70–75.
73. Curtis KA, Tyner TM, Zachary L, et al. Effect of a standard exercise protocol on shoulder pain in long-term wheelchair users. *Spinal Cord* 1999;37(6):421–429.
74. Nawoczenski DA, Ritter-Soron JM, Wilson CM, Howe BA, Ludewig PM. Clinical trial of exercise for shoulder pain in chronic spinal injury. *Phys Ther* 2006;86(12):1604–1618.
75. Hicks AL, Martin KA, Ditor DS, Latimer AE, Craven C, Bugaresti J, McCartney N. Long-term exercise training in persons with spinal cord injury: Effects on strength, arm ergometry performance and psychological well-being. *Spinal Cord* 2003;41(1):34–43.
76. Barbeau H, Nadeau S, Garneau C. Physical determinants, emerging concepts, and training approaches in gait of individuals with spinal cord injury. *J Neurotrauma* 2006;23(3–4):571–585.
77. Field-Fote EC, Lindley SD, Sherman AL. Locomotor training approaches for individuals with spinal cord injury: A preliminary report of walking-related outcomes. *J Neurol Phys Ther* 2005;29(3): 127–137.
78. Nash MS, Jacobs PL, Johnson BM, Field-Fote E. Metabolic and cardiac responses to robotic-assisted locomotion in motor complete tetraplegia: A case report. *J Spinal Cord Med* 2004;27(1):78–82.
79. Dobkin B, Barbeau H, Deforge D, et al; Spinal Cord Injury Locomotor Trial Group. The evolution of walking-related outcomes over the first 12 wk of rehabilitation for incomplete traumatic spinal cord injury: The multicenter randomized Spinal Cord Injury Locomotor Trial. *Neurorehabil Neural Repair* 2007;21(1):25–35.
80. Ekblom B, Lundberg A. Effect of physical training on adolescents with severe motor handicaps. *Acta Paediatr Scand* 1968;57:17–23.
81. O’Connell DG, Barnhart R. Improvement in wheelchair propulsion in pediatric wheelchair users through resistance training: A pilot study. *Arch Phys Med Rehabil* 1995;76:368–372.
82. Yarkony GM, ed. *Spinal Cord Injury: Medical Management and Rehabilitation*. Gaithersburg, MD: Aspen Publishers; 1994.
83. Nesathurai S. *The Rehabilitation of People with Spinal Cord Injury*, 2nd ed. Williston, VT: Blackwell Science; 2000.
84. Stass W, et al. Rehabilitation of the spinal cord injured patient. In: Delisa JA, Gans BM, Bockenek WL, eds. *Rehabilitation Medicine: Principles and Practice*. Baltimore: Lippincott William & Wilkins, 1993:891.
85. Bodner DR, Perkash I. Urological management in spinal cord injury. In: Lin VW, ed. *Spinal Cord Medicine: Principles and Practice*. New York: Demos; 2003:299–306.
86. Physicians’ Desk Reference (PDR). (accessed online, 2-17-08, <http://www.pdr.net>)
87. Figoni SF. Spinal cord injury. In: Durstine JL, ed. *ACSM’s Exercise Management for Persons with Chronic Diseases and Disabilities*. Champaign, IL: Human Kinetics; 1997:175–179.
88. Kofsky PR, Davis GM, Jackson RW, et al. Field testing—Assessment of physical fitness of disabled adults. *Eur J Appl Physiol* 1983;51:109–120.
89. Franklin BA, Swantek KI, Grais SL, et al. Field test estimation of maximal oxygen consumption in wheelchair users. *Arch Phys Med Rehabil* 1990;71:574–578.
90. Pare G, Noreau L, Simard C. Prediction of maximal aerobic power from a submaximal exercise test performed by paraplegics on a wheelchair ergometer. *Paraplegia* 1993;31:584–592.
91. Glaser RM, Sawka MN, Brune MF, et al. Physiological responses to maximal effort wheelchair and arm crank ergometry. *J Appl Physiol* 1980;48:1060–1064.
92. Glaser RM. Arm exercise training for wheelchair users. *Med Sci Sports Exerc* 1989;21(Suppl 5):S149–S157.
93. Rimmer JH, Connor-Kuntz F, Winnick JP, et al. Feasibility of the target aerobic movement test in children and adolescents with spina bifida. *Adapted Phys Activ Q* 1997;14:147–155.
94. UDS: Uniform Data System for Medical Rehabilitation. Accessed 2-17-08 at <http://www.fimssystem.com>
95. Cole B, Finch, Gowland C, et al. *Physical Rehabilitation Outcome Measures*. Baltimore: Williams & Wilkins; 1995.
96. Rimaud D, Calmels P, Devillard X. Training programs in spinal cord injury. *Ann Readapt Med Phys* 2005;48(5):259–269.
97. Jacobs PL, Nash MS. Exercise recommendations for individuals with spinal cord injury. *Sports Med* 2004;34:727–751.
98. Hooker SP, Greenwood JD, Hatae DT, et al. Oxygen uptake and heart rate relationship in persons with spinal cord injury. *Med Sci Sport Exerc*. 1993;25:1115–1119.
99. Irizawa M, Yamasaki M, Muraki S, et al. Relationship between heart rate and oxygen uptake during submaximal arm cranking in paraplegics and quadriplegics. *Ann Physiol Anthropol* 1994;13(5): 275–280.

100. Hayes AM, Myers JN, Ho M, Lee MY, Perakash I, Kiratli BJ. Heart rate as a predictor of energy expenditure in people with spinal cord injury. *J Rehabil Res Dev* 2005;42(5):617–624.
101. Borg G. Borg's Perceived Exertion and Pain Scales. Champaign, IL: Human Kinetics; 1998.
102. Haskell WL, Lee I-M, Pate RR, et al. Physical activity and public health: Updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc* 2007;39(8):1423–1434.
103. Coyle CP, Santiago MC. Aerobic exercise training and depressive symptomatology in adults with physical disabilities. *Arch Phys Med* 1995;76:647–652.
104. Edwards PA. Health promotion through fitness for adolescents and young adults following spinal cord injury. *SCI Nurs* 1996; 13:69–73.
105. Nichols S, Brasile FM. The role of recreational therapy in physical medicine. *Top Spinal Cord Injury Rehabil* 1998;3:89–98.
106. Fioni SF, McClain L, Bell AA, et al. Accessibility of physical fitness facilities in the Kansas City metropolitan area. *Top Spinal Cord Injury Rehabil* 1998;3:66–78.
107. Scelza WM, Kalpakjian CZ, Zemper ED, Tate DG. Perceived barriers to exercise in people with spinal cord injury. *Am J Phys Med Rehabil*. 2005 Aug;84(8):576–583.
108. Morrison SA, Stanwyck DJ. The effect of shorter lengths of stay on functional outcomes of spinal cord injury rehabilitation. *Top Spinal Cord Injury Rehabil* 1999;4:44–55.
109. Johnson KA, Klaas SJ. Recreation issues and trends in pediatric spinal cord injury. *Top Spinal Cord Injury Rehabil* 1997;3:79–84.
110. Steadward R. Musculoskeletal and neurological disabilities: Implications for fitness appraisal, programming, and counseling. *Can J Appl Physiol* 1998;23:131–165.
111. Sallis JF, Hovell MF, Hofstetter CR. Predictors of adoption and maintenance of vigorous physical activity in men and women. *Prev Med* 1992;21:237–251.
112. North Carolina Office on Disability and Health. Removing Barriers to Health Clubs and Fitness Facilities. Chapel Hill, NC: Frank Porter Graham Child Development Center, 2001. (accessed online, 2-17-08, <http://www.fpg.unc.edu/~ncodh/>)
113. Elliott TR, Rank RG. Depression following spinal cord injury. *Arch Phys Med Rehabil* 1996;77:816–823.
114. Lowe C. Basic training (fitness, exercise, and sports). In: Lutkenhoff M, Oppenheimer SG, eds. *Spinabilities: A Young Person's Guide to Spina Bifida*. Bethesda, MD: Woodbine House; 1997:123–131.
115. Lockette KF, Keyes AM. Conditioning with Physical Disabilities. Champaign, IL: Human Kinetics; 1994.
116. Miller P, ed. *Fitness Programming and Physical Disability*. Champaign, IL: Human Kinetics, 1994.
117. Apple DF, ed. *Physical Fitness: A Guide for Individuals with Spinal Cord Injury*. Washington, DC: VA Rehabilitation Research and Development Service; 1996.
118. Youngbauer J, ed. *Deconditioning and Eight Gain. Secondary Conditions Prevention & Treatment. B Series, No. 3*. Lawrence, KS: Research and Training Center on Independent Living, University of Kansas; 1996.
119. Chase TM. Physical fitness strategies. In: Lanig IS, ed. *A Practical Guide to Health Promotion After Spinal Cord Injury*. Gaithersburg, MD: Aspen Publishers; 1996.
120. Laskin JJ, James SA, Cantwell BM. A fitness and wellness program for people with spinal cord injury. *Top Spinal Cord Injury Rehabil* 1997;3:16–33.
121. Virgilio SJ. Fitness education for children with disabilities. In: Virgilio SJ. *Fitness Education for Children*. Champaign, IL: Human Kinetics; 1997:47–58.
122. Rimmer JH. *Fitness and Rehabilitation Programs for Special Populations*. Madison, WI: Wm. C. Brown; 1994.
123. Wheelchair Sports USA website. (accessed online, 2-17-08, <http://www.wsusa.org>)

Chapter 6

1. Thornsteinsson G. Management of postpolio syndrome. *Mayo Clin Pros* 1997;72:627–628.
2. Dalakas MC, Hallett M. The Post-Polio Syndrome. In: Plum F, ed. *Advances in contemporary neurology*. Philadelphia: F.A. Davis, 1988; 15–95.
3. Widar M, Ahlstrom G. Pain in persons with post-polio. *Scand J Caring Sci* 1999;13:33–40.
4. Bruno RL, Crengel SJ, Frick NM. Parallels between post-polio fatigue and chronic fatigue syndrome: a common pathophysiology? *Am J Med* 1998;105:665–738.
5. Horemans HL, Johannes BJ, Beelen A, Stam HJ, Nollet F. Walking in postpoliomyelitis syndrome: The relationships between timescored tests, walking in daily life and perceived mobility problems. *J Rehabil Med* 2005;37:142–146.
6. Aurlen D, Strandjord RE, Hegland O. Postpolio syndrome -a critical comment to the diagnosis. *Acta Neurol Scand* 1999;100:76–80.
7. Birk TJ. Poliomyelitis and the post-polio syndrome: exercise capacities and adaptation-current research, future directions, and widespread applicability. *Med Sci Sports Exerc* 1993;25:466–472.
8. Miller A, Ali OE, Sinert R. Gullian-Barré syndrome. <http://www.emedicine.com/EMERG/topic222.htm>
9. Willen C, Cider A, Summerhagen KS. Physical performance in individuals with late effects of polio. *Scand J Rehabil Med* 1999;31: 244–249.
10. Stanghelle JK, Festvag L, Aksnes A. Pulmonary function and symptom-limited exercise stress testing in subjects with late sequelae of postpoliomyelitis. *Scand J Rehab Med* 1993;25:125–129.
11. Birk TJ. Polio and post-polio syndrome. In: *ACSM's exercise management for persons with chronic diseases and disabilities*. Champaign, IL: Human Kinetics, 1997.
12. Stanghellod JK, Festvag LV. Postpolio syndrome: a 5 year follow-up. *Spinal Cord* 1997;35:503–508.
13. Agre JC, Rodriguez AA, Franke TM. Strength, endurance, and work capacity after muscle strengthening exercise in postpolio subjects. *Arch Phys Med Rehabil* 1997;78:681–687.
14. Spector SA, Gordon PL, Feuerstein IM, et al. Strength gains without muscle injury after strength training in patients with postpolio muscular atrophy. *Muscle & Nerve* 1996;19:1282–1290.
15. Einarsson G, Grindy G. Strengthening exercise program in postpolio subjects. In: Halstead LS, Wiechers DO, eds. *Research and clinical aspects of late effects of poliomyelitis*. White Plains, NY: March of Dimes Birth Defects Foundation, 1987.
16. Allen GM, Middleton J, Katrak PH, Lord SR, Gandevia SC. Prediction of voluntary activation, strength and endurance of elbow flexors in postpolio patients. *Muscle & Nerve* 2004;30:172–181.
17. Chan KM, Amirjani N, Sumrain M, Clarke A, Strohschein FJ. Randomized controlled trial of strength training in post-polio patients. *Muscle & Nerve* 2003;27:332–338.
18. Prevots DR, Sutter RW. Assessment of Gullian-Barré syndrome mortality and morbidity in the United States: implications for acute flaccid paralysis surveillance. *J Infect Dis* 1997;175:S151–S155.
19. Meythaler JM. Rehabilitation of Gullian-Barré syndrome. *Arch Phys Med Rehabil* 1997;78(8):872–879.
20. Dalakas MC, Bartfeld H, Kurland LT. The postpolio syndrome: advances in the pathogenesis and treatment. *Ann N Y Acad Sci* 1995; 753:1–411.
21. Trojan DA, Collet JP, Shapiro S, et al. A multicenter, randomized, double-blinded trial of pyridostigmine in postpolio syndrome. *Neurology* 1999;53:1225–1233.
22. Horemans HL, Nollet F, Beelen A, Drost G, et al. Pyridostigmine in postpolio syndrome: no decline in fatigue and limited functional performance. *J Neurol Neurosurg Psychiatry* 2003;74:1655–1661.
23. Shahar E. Current therapeutic options in severe Guillain-Barre syndrome. *Neuropharmacol* 2006;29:45–51
24. Nollet F, Beelen A, Prins MH. Disability and functional assessment in former polio patients with and without postpolio syndrome. *Arch Phys Med Rehabil* 1999;80:136–143.
25. Farbu E, Gilhus NE, Barnes MP, et al. EFNS guideline on diagnosis and management of post-polio syndrome. Report of an EFNS task force. *European J Neurol* 2006;13:795–801.
26. Farbu E, Rekand T, Vik-Mo E, Lygren H, Gilhus NE, Aarli JA. Postpolio syndrome patients treated with intravenous immunoglobulin: a double-blinded randomized

- controlled pilot study. *European J Neurol* 2007;14:60–65.
27. Heim M, Yaacobi E, Azaria M. A pilot study to determine the efficiency of lightweight carbon fibre orthoses in the management of patients suffering from post-polio myelitis syndrome. *Clin Rehab* 1997;11:302–305.
 28. Julien J, Leparco-Goffart I, Lina B, et al. Postpolio syndrome: poliovirus persistence is involved in the pathogenesis. *J Neurol* 1999; 246:472–476.
 29. Samii A, Lopez-Devine J, Wasserman EM, et al. Normal postexercise facilitation and depression of motor evoked potentials in post-polio patients. *Muscle Nerve* 1998;21:948–950.
 30. Cywinska-Wasilewska G, Ober JJ, Koczocik-Przedpelska J. Power spectrum of the surface EMG in post-polio syndrome. *Electromyogr Clin Neurophysiol* 1998;38:463–466.
 31. Roeleveld K, Sandberg A, Stalberg EV, et al. Motor unit size estimation of enlarged motor units with surface electromyography. *Muscle Nerve* 1998;21:878–886.
 32. Rodriguez AA, Agre JC, Franke TM. Electromyographic and neuromuscular variables in unstable postpolio subjects, stable postpolio subjects, and control subjects. *Arch Phys Med Rehabil* 1997;78: 986–991.
 33. Sunnerhagen KS, Bengtsson BA, Lundberg PA, Landin K, Lindstedt G, Grimby G. Normal concentrations of serum insulin-like growth factor- in late polio. *Arch Phys Med Rehabil* 1995;76:732–735.
 34. Nordgren B, Falck B, Stalberg E, et al. Postpolio muscular dysfunction: relationships between muscle energy metabolism, subjective symptoms, magnetic resonance imaging, electromyography, and muscle strength. *Muscle Nerve* 1997;20:1341–1351.
 35. Hughes RA, Wijidicks EF, Benson E. Supportive care for patients with Guillain-Barré syndrome. *Arch Neurol* 2005;62:1194–1198.
 36. American College of Sports Medicine. ACSM's guidelines for exercise testing and prescription, 7th ed. Baltimore: Lippincott Williams & Wilkins, 2007.
 37. Finch LE, Ventruini A, Mayo NE, Trojan DA. Effort-limited treadmill walk test-reliability and validity in subjects with postpolio syndrome. *Am J Phys Med Rehabil* 2004;83:613–623.
 38. Pitetti KH, Barrett PJ, Abbas D. Endurance exercise training in Guillain-Barré syndrome. *Arch Phys Med Rehabil* 1993;74: 761–765.
 39. Halstead L, Gawne AC. NRH proposal for limb classification and exercise prescription. *Disabil Rehabil* 1996;18:311–316.
 40. Strumse YAS, Stanghelle JK, Utne L, Ahlvin P, Svendsby EK. Treatment of patients with postpolio syndrome in warm climate. *Disabil Rehabil* 2003;25:77–84.
 41. Willen C, Scherman MH, Grimby G. Dynamic water exercise in individuals with late poliomyelitis. *Arch of Phys Med Rehabil* 2001; 82:66–72.
 42. Staff. Strenuous exercise may impair muscle function in Guillain-Barré patients. *JAMA* 1970;214:468–469.
 43. Steinberg JS. Guillain-Barré syndrome (acute idiopathic polyneuritis): an overview for the lay person. Wynnwood, PA: The Guillain-Barré Syndrome Support Group International, 1987.
 44. Karper WB. Effects of low-intensity aerobic exercise on one subject with Chronic Relapsing Guillain-Barré syndrome. *Rehabil Nurs* 1991;16(2):96–98.
 45. Packer TL, Martins I, Krefling L, Brouwer B. Activity and post-polio fatigue. *Orthopedics* 1991;14:1223–1226.
 46. Klefbeck B, Lagerstrand L, Mattsson E. Inspiratory muscle training in patients with prior polio who use part-time assisted ventilation. *Arch Phys Med Rehabil* 2000;81:1065–1071.
 47. Burrows DS. Residual subclinical impairment in patients who totally recovered from Guillain-Barré syndrome: impact on military performance. *Mil Med* 1990;155:438–440.
 48. Melillo EM, Sethi JM, Mohsenin V. Guillain-Barré syndrome: rehabilitation outcome and recent developments. *Yale J Biol Med* 1998; 71(5):383–389.
 49. Bernsen RA, de Jager AE, Schmitz PI, et al. Residual physical outcome and daily living 3 to 6 years after Guillain-Barré syndrome. *Neurology* 1999;53(2):409–410.
 50. Soryal I, Sinclair E, Hornby J, et al. Impaired joint mobility in Guillain-Barré syndrome: a primary or a secondary phenomenon? *J Neurol Neurosurg Psychiatry* 1992;22(11):1014–1017.
 51. Berman AT, Tom L. The Guillain-Barré syndrome in children. *Clin Ortho Related Resp* 1976;116:61–65.

Chapter 7

- Angelini C. The role of corticosteroids in muscular dystrophy: A critical appraisal. *Muscle Nerve* 2007; May 31.
- Abramson AS, Rogoff J. An approach to rehabilitation of children with muscular dystrophy. *Proceedings of the First and Second Medical Conferences of the MDAA, Inc.* New York, MDAA; 1953, p. 123–124
- Aitkens SG, McCrory MA, Kilmer DD, et al. Moderate resistance exercise program: Its effect in slowly progressive neuromuscular disease. *Arch Phys Med Rehabil* 1993;74:711–715.
- Argov Z, Arnold DL. MR Spectroscopy and imaging in metabolic myopathies. *Neurol Clin* 2000;18:35–52.
- Belanger AY, McComas AJ. Extent of motor unit activation during effort. *J Appl Physiol* 1981;51:1131–1135.
- Bonsett CA. Pseudohypertrophic muscular dystrophy: Distribution of degenerative features as revealed by anatomical study. *Neurology* 1963;13:728–738.
- Buchthal F, Rosenfalck P. Action potential parameters in different human muscles. *Acta Physiol Scand* 1955;30:125–131.
- DeLateur BJ, Giaconni RM. Effect on maximal strength of submaximal exercise in Duchenne muscular dystrophy. *Am J Phys Med* 1979;58:26–36.
- DiMarco AF, Kelling J, Sajovic M, et al. Respiratory muscle training in muscular dystrophy. *Clin Res* 1982; 30:427A.
- Dubowitz V. *Muscle Disorders in Childhood*. 2nd ed. Philadelphia: W.B. Saunders; 1995:1–133.
- Escalante A, Miller L, Beardmore TD. Resistive exercise in the rehabilitation of polymyositis/dermatomyositis. *J Rheumatol* 1993;20:1340–1344.
- Gozal D, Thiriet P. Respiratory muscle training in neuromuscular disease: long-term effects on strength and load perception. *Med Sci Sports Exerc* 1999; 31:1522–1527.
- Grange RW, Call JA. Recommendations to define exercise prescription for Duchenne muscular dystrophy. *Exerc Sport Sci Rev* 2007;35:12–17.
- Haller RG, Lewis SF. Glucose-induced exertional fatigue in muscle phosphofructokinase deficiency. *N Engl J Med* 1991;324:364–369.
- Haller RG, Wyrick P, Taivassalo T, Vissing J. Aerobic conditioning: An effective therapy in McArdle's disease. *Ann Neurol* 2006;59(6):922–928.
- Hickok RJ. Physical therapy as related to peripheral nerve lesions. *Phys Ther Rev* 1961; 41:113–117.
- Hicks JE, Miller F, Plotz P, et al. Isometric exercise increases strength and does not produce sustained creatinine phosphokinase increases in a patient with polymyositis. *J Rheumatol* 1993; 20:1399–1401.
- Hicks JE. Role of rehabilitation in the management of myopathies. *Curr Opin Rheumatol* 1998;10:548–555.
- Hoberman M. Physical medicine and rehabilitation: Its value and limitations in progressive muscular dystrophy. *Am J Phys Med* 1955;34:109–115.
- Jeppesen TD, Schwartz M, Olsen DB, et al. Aerobic training is safe and improves exercise capacity in patients with mitochondrial myopathy. *Brain* 2006;129:3402–3412.
- Johnson EW, Braddom R. Over-work weakness in facioscapulohumeral muscular dystrophy. *Arch Phys Med Rehabil* 1971;52:333–336.
- Kilmer DD, McCrory MA, Wright NC, et al. The effect of a high resistance exercise program in slowly progressive neuromuscular disease. *Arch Phys Med Rehabil* 1994;75:560–563.
- Kilmer DD. The role of exercise in neuromuscular disease. *Phys Med Rehabil Clin N Am* 1998;9:115–125.
- Kilmer DD. Response to resistive strengthening exercise training in humans with neuromuscular disease. *Am J Phys Med Rehabil* 2002; 81(11 Suppl):S121–S126.
- Kley RA, Vorgerd M, Tarnopolsky MA. Creatine for treating muscle disorders. *Cochrane Database Syst Rev* 2007(1):CD004760.
- Lenman JA. A clinical and experimental study of the effects of exercise on motor weakness in neurological disease. *J Neurol Neurosurg Psychiatry* 1959;22:182–194.
- Lewis SF, Haller RG. Skeletal muscle disorders and associated factors that limit exercise performance. *Exerc Sport Sci Rev* 1989;17: 67–113.
- Lewis SF, Haller RG, Cook JD, et al. Muscle fatigue in McArdle's disease studied by ³¹P-NMR: Effect of glucose infusion. *J Appl Physiol* 1985;59:1991–1994.
- Lindeman E, Leffers P, Spaans F, et al. Strength training in patients with myotonic dystrophy and hereditary motor and sensory neuropathy: A randomized clinical trial. *Arch Phys Med Rehabil* 1995;76:612–620.

- McCartney N, Moroz D, Garner SH, et al. The effects of strength training in patients with selected neuromuscular disorders. *Med Sci Sports Exerc* 1988;20:362–368.
- McCool FD, Tzelepis GE. Inspiratory muscle training in the patient with neuromuscular disease. *Phys Ther* 1995;75:1006–1014.
- Mendell JR, Moxley RT, Griggs RC, et al. Randomized, double-blind six-month trial of prednisone in Duchenne's muscular dystrophy. *N Engl J Med* 1989;320:1592–1597.
- Milner-Brown HS, Miller RG. Muscle strengthening through high-resistance weight training in patients with neuromuscular disorders. *Arch Phys Med Rehabil* 1988; 69:14–19.
- Olsen DB, Orngreen MC, Vissing J. Aerobic training improves exercise performance in facioscapulohumeral muscular dystrophy. *Neurology* 2005 22;64(6):1064–1066.
- Orngreen MC, Olsen DB, Vissing J. Aerobic training in patients with myotonic dystrophy type 1. *Ann Neurol* 2005; 57(5):754–757.
- Radcliffe KA, Campbell WW. Statin myopathy. *Curr Neurol Neurosci Rep* 2008;8(1):66–72.
- Spector SA, Lemmer JT, Koffman BM, et al. Safety and efficacy of strength training in patients with sporadic inclusion body myositis. *Muscle Nerve* 1997;20:1242–1248.
- Sveen ML, Jeppesen TD, Hauerslev S, Krag TO, Vissing J. Endurance training: An effective and safe treatment for patients with LGMD2I. *Neurology* 2007;68(1):59–61.
- Tarnopolsky M. Exercise testing as a diagnostic entity in mitochondrial myopathies. *Mitochondrion* 2004;4(5-6):529–542.
- Tarnopolsky MA, Raha S. Mitochondrial myopathies: Diagnosis, exercise intolerance, and treatment options. *Med Sci Sports Exerc* 2005;37(12):2086–2093.
- Tarnopolsky MA, Roy BD, MacDonald JR. A randomized, controlled trial of creatine monohydrate in patients with mitochondrial cytopathies. *Muscle Nerve* 1997;20:1502–1509.
- Tarnopolsky MA, Maguire J, Myint T, et al. Clinical, physiological, and histological features in a kindred with the T3271C melas mutation. *Muscle Nerve* 1998;21:25–33.
- Tarnopolsky M, Mahoney D, Vajsar J, et al. Creatine monohydrate increases strength and body composition in boys with duchenne muscular dystrophy. *Neurology* 2004;62(10):1771–1777.
- Tein I. Metabolic myopathies. *Semin Pediatr Neurol* 1996;3:59–98.
- Tollbäck A, Eriksson S, Wredenberg A, et al. Effects of high resistance training in patients with myotonic dystrophy. *Scand J Rehabil Med* 1999;31:9–16.
- Van Beekvelt MC, van Engelen BG, Wevers RA, et al. Quantitative nearinfrared spectroscopy discriminates between mitochondrial myopathies and normal muscle. *Ann Neurol* 1999;46:667–670.
- van der Kooi EL, Lindeman E, Riphagen I. Strength training and aerobic exercise training for muscle disease. *Cochrane Database Syst Rev* 2005(1):CD003907.
- Vignos PJ Jr. Physical models of rehabilitation in neuromuscular disease. *Muscle Nerve* 1983;6:323–338.
- Vignos PJ, Watkins MP. The effect of exercise in muscular dystrophy. *JAMA* 1966;197:89–96.
- Vissing J, Haller RG. The effect of oral sucrose on exercise tolerance in patients with McArdle's disease. *N Engl J Med* 2003;349(26): 2503–2509.
- Wanke T, Toifl K, Merkle M, et al. Inspiratory muscle training in patients with Duchenne muscular dystrophy. *Chest* 1994;105:475–482.
- Wratney MJ. Physical therapy for muscular dystrophy children. *Phys Ther Rev* 1958;38:26–32.
- Wiesinger GF, Quittan M, Aringer M, et al. Improvement of physical fitness and muscle strength in polymyositis/dermatomyositis patients by a training programme. *Br J Rheumatol* 1998;37:196–200.

Chapter 8

- Dyck PJ. Causes, classification and treatment of peripheral neuropathy. *N Eng J Med* 1982;307:283–286.
- Scadding JW. Peripheral neuropathies. In: Wall PD, Melzack R, eds. *Textbook of Pain*. Edinburgh: Churchill Livingstone; 1999:1815.
- Vaillancourt PD, Langevin HM. Painful peripheral neuropathies. *Med Clin North Am* 1999;83:627.
- Perle SM, Schneider MJ, Seaman DR. Chiropractic management of peripheral neuropathy: Pathophysiology, assessment, and treatment. *Top Clin Chiropr* 1999;6:6–19.
- Seaman DR, Cleveland C. Spinal pain syndromes: Nociceptive, neuropathic, and psychologic mechanisms. *J Manipulative Physiol Ther* 1998;22:458–472.
- Belgrade MJ. Following the clues to neuropathic pain. *Postgraduate Med* 1999;106:127–140.
- Bennett GF. Neuropathic pain. In: Wall PD, Melzack R, eds. *Textbook of Pain*. Edinburgh: Churchill Livingstone; 1994:201–224.
- Fields HL, Baron R, Rowbotham MC. Peripheral neuropathic pain: An approach to management. In: Wall PD, Melzack R, eds. *Textbook of Pain*. Edinburgh: Churchill Livingstone; 1999:1523–1533.
- Boulton AJ. Diabetic neuropathy: Classification, measurement and treatment. *Curr Opin Endocrinol Diabetes* 2007;14:141–145.
- Tracy JA, Dyck PJ. The spectrum of diabetic neuropathies. *Phys Med Rehabil Clin N Am* 2008;19:1–26.
- Pollmann W, Feneberg W. Current management of pain associated with multiple sclerosis. *CNS Drugs* 2008; 22:291–324.
- Loar, C. Peripheral nervous system pain. In: Raj P. *Pain Medicine: A Comprehension Review*. St. Louis: Mosby Year Book; 1996:453–459.
- Oh SJ, ed. *Clinical Electromyography: Nerve Conduction Studies*. Baltimore: Williams & Wilkins; 1993.
- Fisher MA. Peripheral neuropathy. In: Weiner WJ, Goetz CG, eds. *Neurology for the Non-neurologist*, 3rd ed. Philadelphia: JB Lippincott; 1994:154–170.
- Hogan QH. Back pain and radiculopathy. In: Abram S, Haddox J, eds. *The Pain Clinic Manual*, 2nd ed. Philadelphia: Lippincott Williams & Wilkins; 2000:157–166.
- O'Connor PJ, Cook DB. Exercise and pain: The neurobiology, measurement, and laboratory study of pain in relation to exercise in humans. *Exerc Sports Sci Rev* 1999;27:119–166.
- Cardenas DD, Egan KJ. Management of chronic pain. In: Kottke FJ, Lehmann JF, eds. *Krusen's Handbook of Physical Medicine and Rehabilitation*. Philadelphia: WB Saunders; 1990:1162–1191.
- Davis VP, Fillingim RB, Doleys DM, et al. Assessment of aerobic power in chronic pain patients before and after a multidisciplinary treatment program. *Arch Phys Med Rehabil* 1992;73: 726–729.
- Ruhland JL, Shields RK. The effects of a home exercise program on impairment and health-related quality of life in persons with chronic peripheral neuropathies. *Phys Ther* 1997;77:1026–1039.
- Rozmaryn LM, Dovel S, Rothman ER, et al. Nerve and tendon gliding exercises and the conservative management of carpal tunnel syndrome. *J Hand Ther* 1998;11:171–179.
- Seradge H, Bear C, Bithell D. Preventing carpal tunnel syndrome and cumulative trauma disorder: Effect of carpal tunnel decompression exercises. An Oklahoma experience. *J Okla State Med Assoc* 2000;93:150–153.
- Lincoln AE, Bernick JS, Ogatis S, et al. Interventions for the primary prevention of work-related carpal tunnel syndrome. *Am J Prev Med* 2000;18:37–50.
- Albright A, Franz M, Hornsby G, et al. American College of Sports Medicine Position Stand. Exercise and type 2 diabetes. *Med Sci Sports Exerc* 2000;32:1324–1360.
- Richardson JK, Sandman D, Vela S. A focused exercise regimen improves clinical measures of balance in patients with peripheral neuropathy. *Arch Phys Med Rehabil* 2001;82:205–209.
- Attal N. Pharmacologic treatment of neuropathic pain. *Acta Neurol Belg* 2001;101:53–64.
- Dray A. Neuropathic pain: Emerging treatments. *Br J Anaesth* 2008; 101:48–58.
- Julien RM. *A Primer of Drug Action. A Concise, Nontechnical Guide to the Actions, Uses, and Side Effects of Psychoactive Drugs*, 8th ed. New York: W. H. Freeman; 1998:181–222.

28. Pariente A, Danan G. Gastrointestinal disorders. In: Benichou C, eds. *Adverse Drug Reactions: A Practical Guide to Diagnosis and Management*. New York: John Wiley & Sons; 1994:77–86.
29. Davis DP, Videen JS, Marino A, et al. Exercise-associated hyponatremia in marathon runners: A two-year experience. *J Emerg Med* 2001;21:47–57.
30. Hamilton ME, Gershwin ME. Treatment of pain. In: Gershwin ME, Hamilton ME, eds. *The Pain Management Handbook*. Totowa, NJ: Humana Press; 1998:283–235.
31. Waslick BD, Walsh BT, Greenhill LL, et al. Cardiovascular effects of desipramine in children and adults during exercise testing. *J Am Acad Child Adolesc Psychiatry* 1999;38:179–186.
32. Lipman AG. Analgesic drugs for neuropathic and sympathetically maintained pain. *Clin Geriatr Med* 1996;12:501–515.
33. Harke H, Gretenkort P, Ladleif HU, et al. The response of neuropathic pain and pain in complex regional pain syndrome I to carbamazepine and sustained-release morphine in patients pretreated with spinal cord stimulation: A double-blinded randomized study. *Anesth Analg* 2001;92:488–495.
34. Minami T, Bakoshi S, Nakano H, et al. The Effects of capsaicin cream on prostaglandin-induced allodynia. *Anesth Analg* 2001;93: 419–423.
35. Ho J, DeLuca KG. Neurologic assessment of the pain patient. In: Benzon H, Raja S, Borsook D, et al., eds. *Essentials of Pain Medicine and Regional Anesthesia*. New York: Churchill Livingstone; 1999: 14–15.
36. Galluzzi KE. Managing neuropathic pain. *J Am Osteopath Assoc* 2007;107:ES39-ES48.
37. Tan JC. *Practical Manual of Physical Medicine and Rehabilitation*. St. Louis: Mosby Yearbook; 1998:133–155, 607–644.
38. Williams V, Pappagallo M. Entrapment neuropathies. In: Benzon H, Raja S, Borsook D, et al., eds. *Essentials of Pain Medicine and Regional Anesthesia*. Philadelphia: Churchill Livingstone; 1999:298.
39. Rathmell J, Katz J. Diabetic and other peripheral neuropathies. In: Benzon H, Raja S, Borsook D, et al., eds. *Essentials of Pain Medicine and Regional Anesthesia*. Philadelphia: Churchill Livingstone; 1999: 288–294.
40. Backonia M. Anticonvulsants for neuropathic pain syndromes. *Clin J Pain* 2000;16:S67–S72.
41. Tan JC. *Practical Manual of Physical Medicine and Rehabilitation*. St. Louis: Mosby Yearbook; 1998:133–155, 607–644.
42. Abram SE. Neural blockade for neuropathic pain. *Clin J Pain* 2000; 16:S56–S61.
43. Raj P. *Neural Blockade in Clinical Anesthesia and Management of Pain*. St. Louis: Mosby Year Book; 1996:899–934.
44. Levy R. Neuroablative procedures for treatment of intractable pain. In: Benzon H, Raja S, Borsook D, et al., eds. *Essentials of Pain Medicine and Regional Anesthesia*. Philadelphia: Churchill Livingstone; 1999:104–110.
45. Carter GT. Rehabilitation management of peripheral neuropathy. *Semin Neurol* 2005;25:229-237.
46. Kemp SS, Rogg JM. In: Latchaw, ed. *MR and Imaging of the Head, Neck, and Spine*. St. Louis: Mosby Year Book; 1991:1109–1157.
47. Deusch AL, Mink JH. Magnetic resonance imaging of musculoskeletal disorders. *Radiol Clin North Am* 1989;27:983–1002.
48. Daffner RH, Rothfus WB. In: Latchaw, ed. *MR and Imaging of the Head, Neck, and Spine*. St. Louis: Mosby Year Book; 1999:1225–1255.
49. Intracosa JH, Christopherson LA. *Radiology of the spine*. In: Benzon H, Raja S, Borsook D, et al., eds. *Essentials of Pain Medicine and Regional Anesthesia*. New York: Churchill Livingstone; 1999: 20–26.
50. Nishida T, Miniek M. Role of neurophysiologic testing for pain. In: Benzon H, Raja S, Borsook D, et al., eds. *Essentials of Pain Medicine and Regional Anesthesia*. New York: Churchill Livingstone; 1999:27–33.
51. Kimura J. *Electrodiagnosis in Diseases of Nerve and Muscle: Principles and Practice*, 2nd ed. Philadelphia: FA Davis; 1989.
52. Wilborn AJ. The electrodiagnostic exam of patients with radiculopathies. *Muscle Nerve* 1998;32:1612–1631.
53. Dumitru D. *Electrodiagnostic Medicine*. Philadelphia: Hanley and Belfus; 995.
54. Griffin JW, Hiesch ST, McArthur JC, et al. Laboratory testing in peripheral neuropathy. *Neurol Clin* 1996; 14:119–133.
55. Huskisson EC. Measurement of pain. *Lancet* 1974;2:1127–1131.
56. Huskisson EC. Measurement of pain. *J Rheumatol* 1982;9:768–769.
57. Dixon JS. Reproducibility along a 10 cm vertical visual analogue scale. *Ann Rheumatol Dis* 1981;40:87–89.
58. Scott J, Huskisson EC. Vertical or horizontal visual analogue scales. *Ann Rheumatol Dis* 1979;38:560.
59. Maxwell C. Sensitivity and accuracy of the visual analogue scale. *Br J Clin Pharmacol* 1978;6:15.
60. Melzack R, Katz J. Pain measurement in persons in pain. In: Wall PD, Melzack R, eds. *Textbook of Pain*. Edinburgh: Churchill Livingstone; 1999:409–426.
61. Skogsbergh DR, Chapman SA. Dealing with the chronic patient. In: Mootz RD, Vernon HT, eds. *Best Practices in Clinical Chiropractic*. Gaithersburg, MD: Aspen; 1999:120–129.
62. Cardenas DD, Egan KJ. Management of chronic pain. In: Kottke FJ, Lehmann JF, eds. *Krusen's Handbook of Physical Medicine and Rehabilitation*. Philadelphia: WB Saunders; 1990:1162–1191.
63. Yeh C, Gonyea M, Lemke J, et al. Physical therapy. In: Aronoff GM, ed. *Evaluation and Treatment of Chronic Pain*. Baltimore: Williams & Wilkins. 985:251–261.
64. Kilmer DD, Aitkens S. Neuromuscular disease. In: Frontera WR, Dawson DM, Slovic DM, eds. *Exercise in Rehabilitative Medicine*. Champaign, IL: Human Kinetics; 1999:253–266.

Chapter 9

1. American College of Sports Medicine. *Guidelines for Exercise Testing and Prescription*. Philadelphia: Lippincott Williams & Wilkins; 2001.
2. Ashman TA, Gordon WA, Cantor JB, Hibbard MR. Neurobehavioral consequences of traumatic brain injury. *Mt. Sinai J Med* 2006; 73(7):999–1005.
3. Barnard P, Dill H, Eldredge P, Held JM, Judd DL, Nalette E. Reduction of hypertonicity by early casting in a comatose head-injured individual. A case report. *Phys Ther* 1984;64(10):1540–1542.
4. Bateman A, Culpán FJ, Pickering AD, Powell JH, Scott OM, Greenwood RJ. The effect of aerobic training on rehabilitation outcomes after recent severe brain injury: A randomized controlled evaluation. *Arch Phys Med Rehabil* 2001;82(2):174–182.
5. Bell SE, Hlatky R. Update in the treatment of traumatic brain injury. *Curr Treat Options Neurol* 2006;8(2):167–175.
6. Bhambhani Y, Maikala R, Farag M, Rowland G. Reliability of nearinfrared spectroscopy measures of cerebral oxygenation and blood volume during handgrip exercise in nondisabled and traumatic brain-injured subjects. *J Rehabil Res Dev* 2006;43(7):845–856.
7. Bhambhani Y, Rowland G, Farag M. Effects of circuit training on body composition and peak cardiorespiratory responses in patients with moderate to severe traumatic brain injury. *Arch Phys Med Rehabil* 2005;86(2):268–276.
8. Bhambhani Y, Rowland G, Farag M. Reliability of peak cardiorespiratory responses in patients with moderate to severe traumatic brain injury. *Arch Phys Med Rehabil* 2003;84(11):1629–1636.
9. Binder S, Corrigan JD, Langlois JA. The public health approach to traumatic brain injury: An overview of CDC's research and programs. *J Head Trauma Rehabil* 2005;20(3):189–195.
10. Brain Trauma Foundation, American Association of Neurological Surgeons, Congress of Neurological Surgeons and Joint Section on Neurotrauma and Critical Care, AANS/CNS. Guidelines for the management of severe traumatic brain injury. XIV. Hyperventilation. *J Neurotrauma* 2007;24(Suppl 1):S87–S90.
11. Brown TH, Mount J, Roulund BL, Kautz KA, Barnes RM, Kim J. Body weight-supported treadmill training versus conventional gait training for people with chronic traumatic brain injury. *J Head Trauma Rehabil* 2005;20(5):402–415.
12. Butefisch CM. Neurobiological bases of rehabilitation. *Neurol Sci* 2006;27(Suppl 1):S18–S23.
13. Bhambhani Y, Coutts K, Gillespie M, et al. Inclusive Fitness and Lifestyle Services of All Disabilities. Canadian Society for Exercise Physiology; 2002.
14. Cantu RC. Posttraumatic retrograde and anterograde amnesia: Pathophysiology and implications in grading and safe return to play. *Journal of Athletic Training* 2001;36:244–248.
15. Centers for Disease Control and Prevention. *Traumatic Brain Injury in the United States: A Report to Congress*. Atlanta, Georgia: Centers for Disease Control

- and Prevention; 1999.
16. Coles JP. Imaging after brain injury. *Br J Anaesth* 1997;99(1):49–60.
 17. Convertino VA, Bloomfield SA, Greenleaf JE. An overview of the issues: Physiological effects of bed rest and restricted physical activity. *Med Sci Sports Exerc* 1997;29(2):187–190.
 18. Dault MC, Dugas C. Evaluation of a specific balance and coordination programme for individuals with a traumatic brain injury. *Brain Inj* 2002;16(3):231–244.
 19. Dawes H, Bateman A, Culpán J, Scott O, Wade DT, Roach N, Greenwood R. The effect of increasing effort on movement economy during incremental cycling exercise in individuals early after acquired brain injury. *Clin Rehabil* 2003;17(5):528–534.
 20. Dawes H, Scott OM, Roach NK, Wade DT. Exertional symptoms and exercise capacity in individuals with brain injury. *Disabil Rehabil* 2006;28(20):1243–1250.
 21. Dawes HN, Barker KL, Cockburn J, Roach N, Scott O, Wade D. Borg's rating of perceived exertion scales: Do the verbal anchors mean the same for different clinical groups? *Arch Phys Med Rehabil* 2005;86(5):912–916.
 22. Driver S, O'Connor J, Lox C, Rees K. Evaluation of an aquatics programme on fitness parameters of individuals with a brain injury. *Brain Inj* 2004;18(9):847–859.
 23. Giacino JT, Kalmar K, Whyte J. The JFK Coma Recovery Scale-Revised: Measurement characteristics and diagnostic utility. *Arch Phys Med Rehabil* 2004;85(12):2020–2029.
 24. Goldstein LB. Prescribing of potentially harmful drugs to patients admitted to hospital after head injury. *J Neurol Neurosurg Psychiatry* 1995;58(6):753–755.
 25. Gordon WA, Zafonte R, Cicerone K, et al. Traumatic brain injury rehabilitation: State of the science. *Am J Med Rehabil* 2006;85(4):343–382.
 26. Gracies JM. Physical modalities other than stretch in spastic hypertonia. *Phys Med Rehabil Clin North Am* 2001;12(4):769–792, vi. 27. Gray DS. Slow-to-recover severe traumatic brain injury: A review of outcomes and rehabilitation effectiveness. *Brain Inj* 2000;14(11): 1003–1014.
 28. Gray DS, Burnham RS. Preliminary outcome analysis of a long-term rehabilitation program for severe acquired brain injury. *Arch Phys Med Rehabil* 2000;81(11):1447–1456.
 29. Henson MB, De Castro JM, Stringer AY, Johnson C. Food intake by brain-injured humans who are in the chronic phase of recovery. *Brain Inj* 1993;7(2):169–178.
 30. Horn S, Watson M, Wilson BA, McLellan DL. The development of new techniques in the assessment and monitoring of recovery from severe head injury: A preliminary report and case history. *Brain Inj* 1992;6(4):321–325.
 31. Hunter M, Tomberlin J, Kirkikis C, Kuna ST. Progressive exercise testing in closed head-injured subjects: Comparison of exercise apparatus in assessment of a physical conditioning program. *Phys Ther* 1990;70(6):363–371.
 32. Irdesel J, Aydiner SB, Akgoz S. Rehabilitation outcome after traumatic brain injury. *Neurocirugia (Astur)* 2007;18(1):5–15.
 33. Iverson GL, Gaetz M, Lovell MR, Collins MW. Cumulative effects of concussion in amateur athletes. *Brain Inj* 2004;18(5):433–443.
 34. Jackson D, Turner-Stokes L, Culpán J, Bateman A, Scott O, Powell J, Greenwood R. Can brain-injured patients participate in an aerobic exercise programme during early inpatient rehabilitation? *Clin Rehabil* 2001;15(5):535–544.
 35. Jankowski LW, Sullivan SJ. Aerobic and neuromuscular training: Effect on the capacity, efficiency, and fatigability of patients with traumatic brain injuries. *Arch Phys Med Rehabil* 1990;71(7):500–504.
 36. Jankowski LW, Sullivan SJ. Aerobic and neuromuscular training: Effect on the capacity, efficiency, and fatigability of patients with traumatic brain injuries. *Arch Phys Med Rehabil* 1990;71(7):500–504.
 37. Jorge RE, Robinson RG, Moser D, Tateno A, Crespo-Facorro B, Arndt S. Major depression following traumatic brain injury. *Arch Gen Psychiatry* 2004;61(1):42–50.
 38. Kraus JF, McArthur DL. Epidemiologic aspects of brain injury. *Neurol Clin* 1996;14(2):435–450.
 39. Laatsch L, Krisky C. Changes in fMRI activation following rehabilitation of reading and visual processing deficits in subjects with traumatic brain injury. *Brain Inj* 2006;20(13–14):1367–1375.
 40. LaRoche SM, Helmers SL. The new antiepileptic drugs: Clinical applications. *JAMA* 2004;291(5):615–620.
 41. Lequerica AH, Rapport LJ, Loehner K, Axelrod BN, Vangel SJ, Jr, Hanks RA. Agitation in acquired brain injury: Impact on acute rehabilitation therapies. *J Head Trauma Rehabil* 2007;22(3): 177–183.
 42. Lim HB, Smith M. Systemic complications after head injury: A clinical review. *Anaesthesia* 2007;62(5):474–482.
 43. McKinlay WW, Watkiss AJ. Cognitive and behavioral effects of brain injury. In: Rosebthal M, Kreutzer J, Griffith E, Pentland B, eds. *Rehabilitation of the Adult and Child with Traumatic Brain Injury*. Philadelphia: FA Davis; 1999:74–86.
 44. Mehrholz J, Major Y, Meissner D, Sandi-Gahun S, Koch R, Pohl M. The influence of contractures and variation in measurement stretching velocity on the reliability of the Modified Ashworth Scale in patients with severe brain injury. *Clin Rehabil* 2005;19(1):63–72.
 45. Mossberg KA. Reliability of a timed walk test in persons with acquired brain injury. *Am J Med Rehabil* 2003;82(5):385–90; quiz 391–392.
 46. Mossberg KA, Ayala D, Baker T, Heard J, Masel B. Aerobic capacity after traumatic brain injury: Comparison with a nondisabled cohort. *Arch Phys Med Rehabil* 2007;88(3):315–320.
 47. Mossberg KA, Greene BP. Reliability of graded exercise testing after traumatic brain injury: Submaximal and peak responses. *Am J Med Rehabil* 2005;84(7):492–500.
 48. Painter P, Durstine JL, Rimmer J, Morgan D, Franklin B, Pitteti K. Increasing physical activity in disabled populations. *Medicine and Science in Sports and Exercise* 1998;30(5, Suppl):S86.
 49. Parnes LS, Agrawal SK, Atlas J. Diagnosis and management of benign paroxysmal positional vertigo (BPPV). *CMAJ* 2003;169(7): 681–693.
 50. Putnam S, Fichtenberg N. Neuropsychological examination of the patient with traumatic brain injury. In: Rosenthal M, Kreutzer J, Griffith E, Pentland B, eds. *Rehabilitation of the Adult and Child with Traumatic Brain Injury*. Philadelphia: FA Davis; 1999:147–166.
 51. Ragueneau JL, Gambini D, Levante A, Riche F, de Vernejoul P, Echter E. [Monitoring of extra and intra-cellular compartment through total body impedance (author's transl)]. *Anesth Analg (Paris)* 1979;36(9–10):439–443.
 52. Ruff R. Two decades of advances in understanding of mild traumatic brain injury. *J Head Trauma Rehabil* 2005;20(1):5–18. 53. Salazar AM, Jabbari B, Vance SC, Grafman J, Amin D, Dillon JD. Epilepsy after penetrating head injury. I. Clinical correlates: A report of the Vietnam Head Injury Study. *Neurology* 1985;35(10): 1406–1414.
 54. Satkunam LE. Rehabilitation medicine: 3. Management of adult spasticity. *CMAJ* 2003;169(11):1173–1179.
 55. Scherzer BP. Rehabilitation following severe head trauma: Results of a three-year program. *Arch Phys Med Rehabil* 1986;67(6):366–374.
 56. Seel RT, Wright G, Wallace T, Newman S, Dennis L. The Utility of the FIM + FAM for Assessing Traumatic Brain Injury Day Program Outcomes. *J Head Trauma Rehabil* 2007;22(5):267–277.
 57. Singer BJ, Jegasothy GM, Singer KP, Allison GT, Dunne JW. Incidence of ankle contracture after moderate to severe acquired brain injury. *Arch Phys Med Rehabil* 2004;85(9):1465–1469.
 58. Taber KH, Warden DL, Hurley RA. Blast-related traumatic brain injury: What is known? *J Neuropsychiatry Clin Neurosci* 2006;18(2): 141–145.
 59. Teasdale G, Jennett B. Assessment of coma and impaired consciousness. *Lancet* 1974;2:81–84.
 60. Temkin NR, Dikmen SS, Anderson GD, et al. Valproate therapy for prevention of posttraumatic seizures: A randomized trial. *J Neurosurg* 1999;91(4):593–600.
 61. Thurman DJ, Coronado V, Selassie A. The epidemiology of brain injury: Implications for public health. In: Zasler ND, Katz DI, Zafonte RD, eds. *Brain Injury Medicine, Principles and Practice*. New York: Demos Medical Publishing; 2007:13:45–55.
 62. Varghese G. Heterotopic ossification. *Phys Med Rehabil Clin North Am* 1992:407–415.
 63. Vaynman S, Gomez-Pinilla F. License to run: Exercise impacts functional plasticity in the intact and injured central nervous system by using neurotrophins. *Neurorehabil Neural Repair* 2005;19(4):283–295.
 64. Verweij BH, Amelink GJ, Muizelaar JP. Current concepts of cerebral oxygen transport and energy metabolism after severe traumatic brain injury. *Prog Brain Res* 2007;161:111–124.

65. Vitale AE, Jankowski LW, Sullivan SJ. Reliability for a walk/run test to estimate aerobic capacity in a brain-injured population. *Brain Inj* 1997;11(1):67–76.
66. Vitale AE, Sullivan SJ, Jankowski LW. Underestimation of subjects' monitored radial pulse rates following traumatic brain injury. *Percept Mot Skills* 1995;80(1):57–58.
67. Vitale AE, Sullivan SJ, Jankowski LW, Fleury J, Lefrancois C, Lebouhiller E. Screening of health risk factors prior to exercise or a fitness evaluation of adults with traumatic brain injury: A consensus by rehabilitation professionals. *Brain Inj* 1996;10(5):367–375.
68. Wehman P, Targett P, West M, Kregel J. Productive work and employment for persons with traumatic brain injury: What have we learned after 20 years? *J Head Trauma Rehabil* 2005;20(2):115–127.
69. Wilson DJ, Powell M, Gorham JL, Childers MK. Ambulation training with and without partial weightbearing after traumatic brain injury: Results of randomized, controlled trial. *Am J Med Rehabil* 2006;85(1):68–74.
70. Wolman RL, Cornall C. Aerobic training in brain-injured patients. *Clinical Rehabilitation* 1994;8:253–257.
71. Yamada T, Yeh M, Kimura J. Fundamental principles of somatosensory evoked potentials. *Phys Med Rehabil Clin North Am*. 2004;15:19–42.
72. Zhu XL, Poon WS, Chan CC, Chan SS. Does intensive rehabilitation improve the functional outcome of patients with traumatic brain injury (TBI)? A randomized controlled trial. *Brain Inj* 2007; 21(7):681–690.
73. Ziino C, Ponsford J. Measurement and prediction of subjective fatigue following traumatic brain injury. *J Int Neuropsychol Soc* 2005; 11(4):416–425.

Chapter 10

1. Centers for Disease Control and Prevention (CDC). Prevalence of disabilities and associated health conditions among adults: United States, 1999. *MMWR* 2001;50:120–5.
2. Theis KA, Murphy L, Hootman JM, Helmick CG, Yelin E. Prevalence and correlates of arthritis-attributable work limitation in the US population among persons ages 18–64:2002 National Health Interview Survey data. *Arthritis Care Res* 2007;57(3):355–63.
3. Klippel JH, Stone JH, Crofford LJ, White PH, eds. *Primer on the Rheumatic Diseases*. 13th ed. New York: Springer; 2008.
4. Lawrence RC, Felson DT, Helmick CG, et al. Estimates of the prevalence of arthritis and other rheumatic conditions in the United States. Part II. *Arthritis Rheum* 2008;58:26–35.
5. Peyron JG, Altman RD. The epidemiology of osteoarthritis. In: Moskowitz RW, Howell DS, Goldberg VM, Mankin HJ, eds. *Osteoarthritis: Diagnosis and Medical/Surgical Management*. 2nd ed. Philadelphia: WB Saunders; 1992:15–37.
6. Moskowitz RW. Clinical and laboratory findings in osteoarthritis. In: Koopman WJ, editor. *Arthritis and Allied Conditions: A Textbook of Rheumatology*. 13th ed. Baltimore: Williams & Wilkins; 1997: 1985–2011.
7. Kellgren JH, Lawrence JS, Bier F. Genetic factors in generalized osteoarthrosis. *Ann Rheum Dis* 1963;22:237–55.
8. Felson DT, Naimark A, Anderson JJ, Kazis L, Castelli W, Meenan RF. The prevalence of knee osteoarthritis in the elderly: The Framingham Osteoarthritis Study. *Arthritis Rheum* 1987;30:914–8.
9. Felson DT. Epidemiology of the rheumatic diseases. In: Koopman WJ, editor. *Arthritis and Allied Conditions: A Textbook of Rheumatology*. 13th ed. Baltimore: Williams & Wilkins; 1997:3–34.
10. Peyron JG. Epidemiologic and etiologic approach to osteoarthritis. *Semin Arthritis Rheum* 1979;8:288–306.
11. Anderson J, Felson DT. Factors associated with knee osteoarthritis (OA) in the HANES I survey: Evidence of an association with overweight, race and physical demands of work. *Am J Epidemiol* 1988;128:179–89.
12. Felson DT, Zhang Y, Hannan MT, Naimark A, Weissman B, Aliabadi P, Levy P. Risk factors for incident radiographic knee osteoarthritis in the elderly: The Framingham study. *Arthritis Rheum* 1997;40:728–33.
13. Altman R, Alarcón G, Applerouth D, et al. The American College of Rheumatology criteria for classification and reporting of osteoarthritis of the hand. *Arthritis Rheum* 1990;33:1601–10.
14. Altman R, Asch E, Bloch D, et al. Development of criteria for classification and reporting of osteoarthritis. Classification of osteoarthritis of the knee. *Arthritis Rheum* 1986;29:1039–49.
15. Altman R, Alarcón G, Applerouth D, et al. Criteria for classification and reporting of osteoarthritis of the hip. *Arthritis Rheum* 1991;34:505–15.
16. Brandt KD. Nonsurgical management of osteoarthritis, with an emphasis on nonpharmacologic measures. *Arch Fam Med* 1995;4: 1057–64.
17. Albani S, Carson DA. Etiology and pathogenesis of rheumatoid arthritis. In: Koopman WJ, ed. *Arthritis and Allied Conditions: A Textbook of Rheumatology*. 13th ed. Baltimore: Williams & Wilkins; 1997:979–92.
18. Helmick CG, Felson DT, Lawrence RC, et al. Estimates of the prevalence of arthritis and other rheumatic conditions in the United States. Part I. *Arthritis Rheum* 2008;58:15–25.
19. Wolfe F, Mitchell DM, Sibley JT, et al. The mortality of rheumatoid arthritis. *Arthritis Rheum* 1994;37:481–94.
20. Hochberg MC, Chang RW, Dwosh I, Lindsey S, Pincus T, Wolfe F. The American College of Rheumatology 1991 revised criteria for the classification of global functional status in rheumatoid arthritis. *Arthritis Rheum* 1992;35:498–502.
21. Hale LP, Haynes BF. Pathology of rheumatoid arthritis and associated disorders. In: Koopman WJ, ed. *Arthritis and Allied Conditions: A Textbook of Rheumatology*. 13th ed. Baltimore: Williams & Wilkins; 1997:993–1016.
22. Goldenberg DL. Treatment of fibromyalgia syndrome. *Rheum Dis Clin North Am* 1989;15:61–71.
23. Buckelew SP. Fibromyalgia: A rehabilitation approach. *Am J Phys Med Rehabil* 1989;68:37–42.
24. Goldenberg DL, Burckhardt C, Crofford L. Management of fibromyalgia syndrome. *JAMA* 2004;292:2388–95.
25. Bradley LA, Alarcón GS. Fibromyalgia. In: Koopman WJ, ed. *Arthritis and Allied Conditions: A Textbook of Rheumatology*. 13th ed. Baltimore: Williams & Wilkins; 1997:1619–40.
26. Lawrence RC, Helmick CG, Arnett FC, et al. Estimates of the prevalence of arthritis and selected musculoskeletal disorders in the United States. *Arthritis Rheum* 1998;41:778–99.
27. Wolfe F, Smythe HA, Yunus MB, et al. The American College of Rheumatology 1990 Criteria for the Classification of Fibromyalgia: Report of the Multicenter Criteria Committee. *Arthritis Rheum* 1990;33:160–72.
28. Mease P. Fibromyalgia syndrome. Review of clinical presentation, pathogenesis, outcome measures and treatment. *J Rheumatol* 2005;32:6–21.
29. Burckhardt CS, O'Reilly CA, Wien AN. Assessing depression in fibromyalgia patients. *Arthritis Care Res* 1994;1:35–9.
30. Arthritis Foundation, Association of State and Territorial Health Officials, and Centers for Disease Control and Prevention. *National Arthritis Action Plan: A Public Health Strategy*. Arthritis Foundation National Office, Atlanta, GA; 1999.
31. Mannerkorpi K, Burckhardt CS, Bjelle A. Physical performance characteristics of women with fibromyalgia. *Arthritis Care Res* 1994;7:123–9.
32. Minor MA, Hewett JE, Webel RR, Dreisinger TE, Kay DR. Exercise tolerance and disease related measures in patients with rheumatoid arthritis and osteoarthritis. *J Rheumatol* 1988;15:905–11.
33. Jones KD, Burckhardt CS, Clark SR, Bennett RM, Potempa KM. A randomized controlled trial of muscle strengthening versus flexibility training in fibromyalgia. *J Rheumatol* 2002;29(5):1041–8.
34. Hurley MV, Scott DL. Improvements in quadriceps sensorimotor function and disability of patients with knee osteoarthritis following a clinically practicable exercise regimen. *Br J Rheumatol* 1998;37:1181–7.
35. Ettinger WH Jr., Afable RF. Physical disability from knee osteoarthritis: The role of exercise as an intervention. *Med Sci Sports Exerc* 1994;26:1435–40.
36. Slemenda C, Brandt KD, Heilman DK, Mazuca S, Braunstein EM, Katz BP, Wolinsky FD. Quadriceps weakness and osteoarthritis of the knee. *Ann Intern Med* 1997;127:97–104.
37. Tan J, Balci N, Sepici V, Gener FA. Isokinetic and isometric strength in osteoarthrosis of the knee. *Am J Phys Med Rehabil* 1995;74:364–9.

38. Beals CA, Lampman RM, Figley Banwell B, Braunstein EM, Albers JW, Castor CW. Measurement of exercise tolerance in patients with rheumatoid arthritis and osteoarthritis. *J Rheumatol* 1985;12:458–61.
39. Danneskiold-Samsøe B, Lyngberg K, Risum T, Telling M. The effect of water exercise given to patients with rheumatoid arthritis. *Scand J Rehab Med* 1987;19:31–5.
40. Ekdahl C, Broman G. Muscle strength, endurance, and aerobic capacity in rheumatoid arthritis: A comparative study with healthy subjects. *Ann. Rheum Dis* 1992;51:35–40.
41. Hsieh LF, Didenko B, Schumacher HR Jr., Torg JS. Isokinetic and isometric testing of knee musculature in patients with rheumatoid arthritis with mild knee involvement. *Arch Phys Med Rehabil* 1987;68:294–7.
42. Fisher NM, Pendergast DR. Reduced muscle function in patients with osteoarthritis. *Scand J Rehab Med* 1997;29:213–21.
43. Valkeinen H, Alen M, Hannonen P, Häkkinen A, Airaksinen O, Häkkinen K. Changes in knee extension and flexion force, EMG and functional capacity during strength training in older females with fibromyalgia and healthy controls. *Rheumatology* 2004;43: 225–8.
44. Häkkinen A, Häkkinen K, Hannonen P, Alen M. Strength training induced adaptations in neuromuscular function of premenopausal women with fibromyalgia: A comparison with healthy women. *Ann Rheum Dis* 2001;60(1):21–6.
45. Valkeinen H, Häkkinen A, Hannonen P, Häkkinen K, Alen M. Acute heavy-resistance exercise-induced pain and neuromuscular fatigue in elderly women with fibromyalgia and in healthy controls: Effects of strength training. *Arthritis Rheum* 2006;54(4):1334–9.
46. Häkkinen K, Pakarinen A, Hannonen P, Häkkinen A, Airaksinen O, Valkeinen H, Alen M. Effects of strength training on muscle strength, cross-sectional area, maximal electromyographic activity, and serum hormones in premenopausal women with fibromyalgia. *J Rheumatol* 2002;29(6):1287–95.
47. Thyberg I, Hass UAM, Nordenskiöld U, Gerdle B, Skogh T. Activity limitation in rheumatoid arthritis correlates with reduced grip force regardless of sex: The Swedish TIRA Project. *Arthritis Care Res* 2005;53(6):886–96.
48. Häkkinen A, Hannonen P, Nyman K, Häkkinen K. Aerobic and neuromuscular performance capacity of physically active females with early and long-term rheumatoid arthritis compared to matched healthy women. *Scand J Rheumatol* 2002;31:345–50.
49. Ling SM, Xue QL, Simonsick EM, Tian J, Bandeen-Roche K, Fried LP, Bathon JM. Transitions to mobility difficulty associated with lower extremity osteoarthritis in high functioning older women: Longitudinal data from the Women's Health and Aging Study II. *Arthritis Care Res* 2006; 55:256–63.
50. Mengshoel AM, Vollestad NK, Forre O. Pain and fatigue induced by exercise in fibromyalgia patients and sedentary healthy subjects. *Clin Exp Rheumatol* 1995;13:477–82.
51. Philbin EF, Groff GD, Ries MD, Miller TE. Cardiovascular fitness and health in patients with end-stage osteoarthritis. *Arthritis Rheum* 1995;38:799–805.
52. Fisher NM, Pendergast DR. Effects of a muscle exercise program on exercise capacity in subjects with osteoarthritis. *Arch Phys Med Rehabil* 1994;75:792–7.
53. Richards SCM, Scott DL. Prescribed exercise in people with fibromyalgia: Parallel group randomised controlled trial. *BMJ* 2002;325:185–9.
54. Valim V, Oliveira L, Suda A, et al. Aerobic fitness effects in fibromyalgia. *J Rheumatol* 2003;30(5):1060–9.
55. Sutbeyaz ST, Sezer N, Koseoglu BF, Ibrahimoglu F, Tekin D. Influence of knee osteoarthritis on exercise capacity and quality of life in obese adults. *Obesity* 2007;15:2071–6.
56. Fisher NM, Gresham GE, Pendergast DR. Quantitative progressive exercise rehabilitation for osteoarthritis of the knee. *Phys Med Rehabil Clin North Am* 1994;5:785–802.
57. Fisher NM, Pendergast DR, Gresham GE, Calkins E. Muscle rehabilitation: Its effects on muscular and functional performance of patients with knee osteoarthritis. *Arch Phys Med Rehabil* 1991;72: 367–74.
58. Fisher DR, Pendergast DR. Application of quantitative and progressive exercise rehabilitation to patients with osteoarthritis of the knee. *Journal of Back and Musculoskeletal Rehabilitation* 1995;5:33–53.
59. Fisher NM, Gresham GE, Abrams M, Hicks J, Horrigan D, Pendergast DR. Quantitative effects of physical therapy on muscular and functional performance in subjects with osteoarthritis of the knees. *Arch Phys Med Rehabil* 1993;74:840–7.
60. Fisher NM, Gresham G, Pendergast DR. Effects of a quantitative progressive rehabilitation program applied unilaterally to the osteoarthritic knee. *Arch Phys Med Rehabil* 1993;74:1319–26.
61. Fisher NM, Kame VD, Rouse L, Pendergast DR. Quantitative evaluation of a home exercise program on muscle and functional capacity of patients with osteoarthritis. *Am J Phys Med Rehabil* 1994;73:413–20.
62. Fisher NM, White SC, Yack HJ, Smolinski RJ, Pendergast DR. Muscle function and gait in patients with knee osteoarthritis before and after muscle rehabilitation. *Disability Rehabil* 1997;19:47–55.
63. Gowans SE, deHueck A, Voss S, Silaj A, Abbey SE, Reynolds WJ. Effect of a randomized, controlled trial of exercise on mood and physical function in individuals with fibromyalgia. *Arthritis Care Res* 2001;45:519–29.
64. DaCosta D, Abrahamowicz M, Lowensteyn I, Bernatsky S, Dritsa M, Fitzcharles MA, Dobkin PL. A randomized clinical trial of an individualized home-based exercise programme for women with fibromyalgia. *Rheumatology* 2005;44:1422–7.
65. Bennett RM, Clark SR, Goldberg L, Nelson D, Bonafede RP, Porter J, Specht D. Aerobic fitness in patients with fibrositis: A controlled study of respiratory gas exchange and 133 xenon clearance from exercising muscle. *Arthritis Rheum* 1989;32:454–60.
66. Sokka T, Häkkinen A, Kautiainen H, et al. Physical inactivity in patients with rheumatoid arthritis: Data from twenty-one countries in a cross-sectional, international study. *Arthritis Care Res* 2008;59:42–50.
67. Mancuso CA, Rincon M, Sayles W, Paget SA. Comparison of energy expenditure from lifestyle physical activities between patients with rheumatoid arthritis and healthy controls. *Arthritis Care Res* 2007;57:672–8.
68. van den Berg MH, de Boer IG, le Cessie S, Breedveld FC, Vliet Vlieland TPM. Are patients with rheumatoid arthritis less physically active than the general population? *J. Clin Rheumatol* 2007; 13:181–6.
69. Shih M, Hootman JM, Kruger J, Helmick CG. Physical activity in men and women with arthritis. National Health Interview Survey, 2002. *Am J Prev Med* 2006;30(5):385–93.
70. McNair PJ, Marshall RN, Maguire K. Swelling of the knee joint: Effects of exercise on quadriceps muscle strength. *Arch Phys Med Rehabil* 1996;77:896–9.
71. Hurley MV. The role of muscle weakness in the pathogenesis of osteoarthritis. *Rheum Dis Clinics North Am* 1999;25:283–98.
72. Jones KD, Adams D, Winters-Stone K, Burckhardt CS. A comprehensive review of 46 exercise treatment studies in fibromyalgia (1988–2005). *Health Qual Life Outcomes* 2006;4:67.
73. Gowans SE, deHueck A. Effectiveness of exercise in management of fibromyalgia. *Curr Opin Rheumatol* 2004;16:138–42.
74. Rooks DS. Fibromyalgia treatment update. *Curr Opin Rheumatol* 2007;19:111–7.
75. Busch AJ, Barber KAR, Overend TJ, Peloso PMJ, Schachter CL. Exercise for treating fibromyalgia syndrome. *Cochrane Database Syst Rev* 2008;3.
76. Mannerkorpi K. Exercise in fibromyalgia. *Curr Opin Rheumatol* 2005;17:190–4.
77. Westby MD. A health professional's guide to exercise prescription for people with arthritis: A review of aerobic fitness activities. *Arthritis Care Res* 2001;45:501–511.
78. Stenstrom CH, Minor MA. Evidence for the benefit of aerobic and strengthening exercise in rheumatoid arthritis. *Arthritis Care Res* 2003;49:428–34.
79. Mayoux Benhamou MA. Reconditioning in patients with rheumatoid arthritis. *Ann Readapt Med Phys* 2007;50(6):382–5.
80. Vliet Vlieland TPM. Non-drug care for RA—Is the era of evidencebased practice approaching? *Rheumatology* 2007;46:1397–1404.
81. van den Ende CHM, Vliet Vlieland TPM, Munneke M, Hazes JMW. Dynamic exercise therapy for treating rheumatoid arthritis. *Cochrane Database Syst Rev* 2008;3.
82. Roddy E, Zhang W, Doherty M. Aerobic walking or strengthening exercise for osteoarthritis of the knee? A systematic review. *Ann Rheum Dis* 2005;64:544–8.
83. Bartels EM, Lund H, Hagen KB, Dagfinrud H, Christensen R, Danneskiold-Samsøe B. Aquatic exercise for the treatment of knee and hip osteoarthritis. *Cochrane Database Syst Rev* 2008;3.

84. Fransen M, McConnell S, Bell M. Exercise for osteoarthritis of the hip or knee. *Cochrane Database Syst Rev* 2008;3.
85. Hall J, Skevington SM, Maddison PJ, Chapman K. A randomized and controlled trial of hydrotherapy in rheumatoid arthritis. *Arthritis Care Res* 1996;9:206–15.
86. van den Ende CH, Hazes JM, le Cessie S, Mulder WJ, Belfor DG, Breedveld FC, Dijkmans BA. Comparison of high and low intensity training in well controlled rheumatoid arthritis. Results of a randomized clinical trial. *Ann. Rheum Dis* 1996;55:798–805.
87. Dellhag B, Wollersjo I, Bjelle A. Effect of active hand exercise and wax bath treatment in rheumatoid arthritis patients. *Arthritis Care Res* 1992;5:87–92.
88. Huang M-H, Yang R-C, Lee C-L, Chen T-W, Wang M-C. Preliminary results of integrated therapy for patients with knee osteoarthritis. *Arthritis Care Res* 2005;53:812–20.
89. Wang TJ, Belza B, Thompson FE, Whitney JD, Bennett K. Effects of aquatic exercise on flexibility, strength and aerobic fitness in adults with osteoarthritis of the hip or knee. *J Adv Nurs* 2007; 57(2):141–52.
90. Messier SP, Loeser RF, Miller GD, et al. Exercise and dietary weight loss in overweight and obese older adults with knee osteoarthritis: The Arthritis, Diet, and Activity Promotion Trial. *Arthritis Rheum* 2004;50:1501–10.
91. Wyatt FB, Milam S, Manske RC, Deere R. The effects of aquatic and traditional exercise programs on persons with knee osteoarthritis. *J Strength Cond Res* 2001;15:337–40.
92. Peloquin L, Bravo G, Gauthier P, Lacombe G, Billiard J-S. Effects of a cross-training exercise program in persons with osteoarthritis of the knee. A randomized controlled trial. *J Clin Rheumatol* 1999;5: 126–36.
93. Petrella RJ, Bartha C. Home based exercise therapy for older patients with knee osteoarthritis: A randomized clinical trial. *J Rheumatol* 2000;27:2215–21.
94. Mengshoel AM, Komnaes HB, Forre O. The effects of 20 weeks of physical fitness training in female patients with fibromyalgia. *Clin Exp Rheumatol* 1992;10:345–9.
95. O'Reilly SC, Muir KR, Doherty M. Effectiveness of home exercise on pain and disability from osteoarthritis of the knee: A randomized controlled trial. *Ann Rheum Dis* 1999;58:15–19.
96. Røgind H, Bibow-Nielsen B, Jensen B, Møller HC, Frimodt-Møller H, Bliddal H. The effects of a physical training program on patients with osteoarthritis of the knees. *Arch Phys Med Rehabil* 1998;79:1421–7.
97. Maurer BT, Stern AG, Kinossian B, Cook KD, Schumacher HR. Osteoarthritis of the knee: Isokinetic quadriceps exercise versus an educational intervention. *Arch Phys Med Rehabil* 1999;80: 1293–9.
98. Machover S, Sapecky AJ. Effect of isometric exercise on the quadriceps muscle in patients with rheumatoid arthritis. *Arch Phys Med Rehabil* 1966;47:737–41.
99. Lyngberg KB, Danneskiold-Samsøe B, Halskov O. The effect of physical training on patients with rheumatoid arthritis: Changes in disease activity, muscle strength and aerobic capacity. A clinically controlled minimized cross-over study. *Clin Exp Rheumatol* 1988;6:253–60.
100. Ekdahl C, Andersson SI, Moritz U, Svensson B. Dynamic versus static training in patients with rheumatoid arthritis. *Scand J Rheumatol* 1990;19:17–26.
101. Brighton SW, Lubbe JE, van der Merwe CA. The effect of a longterm exercise programme on the rheumatoid hand. *Br J Rheumatol* 1993;32:392–5.
102. Hansen TM, Hansen G, Langgaard AM, Rasmussen JO. Long-term physical training in rheumatoid arthritis. A randomized trial with different training programs and blinded observers. *Scand J Rheumatol* 1993;22:107–12.
103. Lyngberg KB, Ramsing BU, Nawrocki A, Harreby M, Danneskiold Samsøe B. Safe and effective isokinetic knee extension training in rheumatoid arthritis. *Arthritis Rheum* 1994;37:623–8.
104. Hakkinen A, Hakkinen K, Hannonen P. Effects of strength training on neuromuscular function and disease activity in patients with recent-onset inflammatory arthritis. *Scand J Rheumatol* 1994;23:237–42.
105. Rall LC, Meydani SN, Kehayias JJ, Dawson-Hughes B, Roubenoff R. The effect of progressive resistance training in rheumatoid arthritis. Increased strength without changes in energy balance or body composition. *Arthritis Rheum* 1996;39:415–26.
106. Komatireddy GR, Leitch RW, Cella K, Browning G, Minor M. Efficacy of low load resistive muscle training in patients with rheumatoid arthritis functional class II and III. *J Rheumatol* 1997;24: 1531–9.
107. Lyngberg KK, Harreby M, Bentzen H, Frost B, Danneskiold-amsøe B. Elderly rheumatoid arthritis patients on steroid treatment tolerate physical training without an increase in disease activity. *Arch Phys Med Rehabil* 1994;75:1189–95.
108. McMeeken J, Stillman B, Story I, Kent P, Smith J. The effects of knee extensor and flexor muscle training on the timed-up-and-go test in individuals with rheumatoid arthritis. *Physiother Res Int* 1999;4:55–67.
109. Geel SE, Robergs RA. The effect of graded resistance exercise on fibromyalgia symptoms and muscle bioenergetics: A pilot study. *Arthritis Care Res* 2002;47:82–6.
110. Rooks DS, Silverman CB, Kantrowitz FG. The effects of progressive strength training and aerobic exercise on muscle strength and cardiovascular fitness in women with fibromyalgia: A pilot study. *Arthritis Rheum* 2002;47(1):22–8.
111. Valkeinen H, Häkkinen K, Pakarinen A, et al. Muscle hypertrophy, strength development, and serum hormones during strength training in elderly women with fibromyalgia. *Scand J Rheumatol* 2005;34:309–14.
112. Kingsley JD, Panton LB, Toole T, Sirithienthad P, Mathis R, McMillan V. The effects of a 12-week strength-training program on strength and functionality in women with fibromyalgia. *Arch Phys Med Rehabil* 2005;86:1713–21.
113. Gusi N, Tomas-Carus P, Häkkinen A, Häkkinen K, Ortega-Alonso A. Exercise in waist-high warm water decreases pain and improves health-related quality of life and strength in the lower extremities in women with fibromyalgia. *Arthritis Care Res* 2006;55(1):66–73.
114. Callahan LF, Mielenz T, Freburger J, et al. A randomized controlled trial of the People with Arthritis Can Exercise Program: Symptoms, function, physical activity, and psychosocial outcomes. *Arthritis Care Res* 2008;59:92–101.
115. Stenstrom CH, Lindell B, Swanberg E, Swanberg P, Harms-Ring-dahl K, Nordemar R. Intensive dynamic training in water for rheumatoid arthritis functional class II—A long-term study of effects. *Scand J Rheumatol* 1991;20:358–65.
116. Neuberger GB, Press AN, Lindsley HB, et al. Effects of exercise on fatigue, aerobic fitness, and disease activity measures in persons with rheumatoid arthritis. *Res Nurs Health* 1997;20:195–204.
117. van den Ende CHM, Breedveld FC, le Cessie S, Dijkmans BAC, de Mug AW, Hazes JW. Effect of intensive exercise on patients with active rheumatoid arthritis: A randomised clinical trial. *Ann Rheum Dis* 2000;59:615–21.
118. Hakkinen A, Sokka T, Kotaniemi A, Hannonen P. A randomized two-year study of the effects of dynamic strength training on muscle strength, disease activity, functional capacity, and bone mineral density in early rheumatoid arthritis. *Arthritis Rheum* 2001;44(3): 515–22.
119. Hakkinen A. Effectiveness and safety of strength training in rheumatoid arthritis. *Curr Opin Rheumatol* 2004;16:132–7.
120. Hakkinen A, Sokka T, Kautiainen H, Kotaniemi A, Hannonen P. Sustained maintenance of exercise induced muscle strength gains and normal bone mineral density in patients with early rheumatoid arthritis: A 5 year follow-up. *Ann Rheum Dis* 2004;63:910–6.
121. Neuberger GB, Aaronson LS, Gajewski B, Embretson SE, Cagle PE, Loudon JK, Miller PA. Predictors of exercise and effects of exercise on symptoms, function, aerobic fitness, and disease outcomes of rheumatoid arthritis. *Arthritis Care Res* 2007;57:943–52.
122. Rooks DS, Huang J, Bierbaum BE, et al. Effect of preoperative exercise on measures of functional status in men and women undergoing total hip and knee arthroplasty. *Arthritis Care Res* 2006;55: 700–8.
123. Iwamoto J, Takeda T, Sato Y. Effect of muscle strengthening exercises on the muscle strength in patients with osteoarthritis of the knee. *Knee* 2007;15:224–30.
124. Hinman RS, Heywood SE, Day AR. Aquatic physical therapy for hip and knee osteoarthritis: Results of a single-blind randomized controlled trial. *Phys Ther* 2007;87:32–43.
125. Rogers MW, Wilder FV. The effects of strength training among persons with hand osteoarthritis: A two-year follow-up study. *J Hand Ther* 2007;20:244–50.
126. Foley A, Halbert J, Hewitt T, Crotty M. Does hydrotherapy improve strength and physical function in patients with osteoarthritis- A randomised controlled trial comparing a gym based and a hydrotherapy based strengthening programme. *Ann. Rheum Dis* 2003;62:1162–7.

127. Baker KR, Nelson ME, Felson DT, Layne JE, Sarno R, Roubenoff R. The efficacy of home based progressive strength training in older adults with knee osteoarthritis: A randomized controlled trial. *J Rheumatol* 2001;28:1655–65.
128. Hopman-Rock M, Westhoff M. The effects of a health educational and exercise program for older adults with osteoarthritis of the hip or knee. *J Rheumatol* 2000;27:1947–54.
129. McCain GA, Bell DA, Mai FM, Halliday PD. A controlled study of the effects of a supervised cardiovascular fitness training program on the manifestations of primary fibromyalgia. *Arthritis Rheum* 1988;31:1135–41.
130. Martin L, Nutting A, MacIntosh BR, Edworthy SM, Butterwick D, Cook J. An exercise program in the treatment of fibromyalgia. *J Rheumatol* 1996;23:1050–3.
131. Wigers SH, Stiles TC, Vogel PA. Effects of aerobic exercise versus stress management treatment in fibromyalgia. A 4.5 year prospective study. *Scand J Rheumatol* 1996;25:77–86.
132. Mangione KK, McCully K, Gloviak A, Lefebvre I, Hofmann M, Craik R. The effects of high-intensity and low-intensity cycle ergometry in older adults with knee osteoarthritis. *J Gerontol Med Sci* 1999;54A:M184–90.
133. Minor MA, Hewett JE, Webel RR, Anderson SK, Kay DR. Efficacy of physical conditioning exercise in patients with rheumatoid arthritis and osteoarthritis. *Arthritis Rheum* 1989;32:1396–405.
134. Ekblom B, Lovgren O, Alderin M, Fridström M, Sätterström G. Effect of short-term physical training on patients with rheumatoid arthritis I. *Scand J Rheumatol* 1975;4:80–6.
135. Harkcom TM, Lampman RM, Banwell BF, Castor CW. Therapeutic value of graded aerobic exercise training in rheumatoid arthritis. *Arthritis Rheum* 1985;28:32–9.
136. Baslund BK, Lyngberg K, Andersen V, Halkjaer Kristensen J, Hansen M, Klokke M, Pedersen BK. Effect of 8 weeks of bicycle training on the immune system of patients with rheumatoid arthritis. *J Appl Physiol* 1993;75:1691–5.
137. Minor MA, Hewett JE. Physical fitness and work capacity in women with rheumatoid arthritis. *Arthritis Care Res* 1995;8:146–254.
138. Noreau L, Martineau H, Roy L, Belzile M. Effects of a modified dance-based exercise on cardiorespiratory fitness, psychological state and health status of persons with rheumatoid arthritis. *Am J Phys Med Rehabil* 1995;74:19–27.
139. Jentoft ES, Kvalvik AG, Mengshoel AM. Effects of pool-based and land-based aerobic exercise on women with fibromyalgia/chronic widespread muscle pain. *Arthritis Rheum* 2001;45:42–7.
140. Burckhardt CS, Mannerkorpi K, Hedenberg L, Bjelle A. A randomized, controlled clinical trial of education and physical training for women with fibromyalgia. *J Rheumatol* 1994;21:714–20.
141. Gowans SE, deHueck A, Voss S, Richardson M. A randomized, controlled trial of exercise and education for individuals with fibromyalgia. *Arthritis Care Res* 1999;12:120–8.
142. Deyle GD, Henderson NE, Matekel RL, Ryder MG, Garber MB, Allison SC. Effectiveness of manual physical therapy and exercise in osteoarthritis of the knee. A randomized, controlled trial. *Ann Intern Med* 2000;132:173–81.
143. Ettinger WH Jr, Burns R, Messier SP, et al. A randomized trial comparing aerobic exercise and resistance exercise with a health education program in older adults with knee osteoarthritis. The Fitness Arthritis and Seniors Trial (FAST). *JAMA* 1997;277:25–31.
144. Kovar PA, Allegrante JP, MacKenzie CR, Peterson, MG, Gutin B, Charlson ME. Supervised fitness walking in patients with osteoarthritis of the knee. A randomized, controlled trial. *Ann Intern Med* 1992;116:529–34.
145. Rejeski WJ, Ettinger WH, Martin K, Morgan T. Treating disability in knee osteoarthritis with exercise therapy: A central role for self-efficacy and pain. *Arthritis Care Res* 1998;11:94–101.
146. Perlman SG, Connell KJ, Clark A, et al. Dance-based aerobic exercise for rheumatoid arthritis. *Arthritis Care Res.* 1990;3:29–35.
147. Stenstrom CH, Arge B, Sundbom A. Dynamic training versus relaxation training as home exercise for patients with inflammatory rheumatic diseases. A randomized controlled study. *Scand J Rheumatol* 1996;25:28–33.
148. Stenstrom CH, Arge B, Sundbom A. Home exercise and compliance in inflammatory rheumatic diseases—Prospective clinical trial. *J Rheumatol* 1997;24:470–6.
149. Rooks DS, Gautam S, Romeling M, et al. Group exercise, education, and combination self-management in women with fibromyalgia: A randomized trial. *Arch Intern Med* 2007;167(20):2192–200.
150. Mannerkorpi K, Nyberg B, Ahlman M, Ekdahl C. Pool exercise combined with an education program for patients with fibromyalgia syndrome. A prospective, randomized study. *J Rheumatol* 2000; 27(10):2473–81.
151. Mannerkorpi K, Ahlman M, Ekdahl C. Six- and 24-month followup of pool exercise therapy and education for patients with fibromyalgia. *Scand J Rheumatol* 2002;31(5):306–10.
152. Gowans SE, deHueck A. Pool exercise for individuals with fibromyalgia. *Curr Opin Rheumatol* 2007;19:168–73.
153. Gowans SE, deHueck A, Voss S, Silaj A, Abbey SE. Six-month and one-year follow-up of 23 weeks of aerobic exercise for individuals with fibromyalgia. *Arthritis Care Res* 2004;51(6):890–8.
154. Tomas-Carus P, Häkkinen A, Gusi N, Leal A, Häkkinen K, Ortega-Alonso A. Aquatic training and detraining on fitness and quality of life in fibromyalgia. *Med Sci Sports Exerc* 2007;39(7): 1044–50.
155. King SJ, Wessel J, Bhambhani Y, Sholter D, Maksymowych W. The effects of exercise and education, individually or combined, in women with fibromyalgia. *J Rheumatol* 2002;29(12):2620–7.
156. Schachter CL, Busch AJ, Peloso PM, Sheppard MS. Effects of short versus long bouts of aerobic exercise in sedentary women with fibromyalgia: A randomized controlled trial. *Phys Ther* 2003; 83: 340–58.
157. Eversden L, Maggs F, Nightingale P, Jobanputra P. A pragmatic randomised controlled trial of hydrotherapy and land exercises on overall well being and quality of life in rheumatoid arthritis. *BMC Musculoskelet Disord* 2007;8:23.
158. Hughes SL, Seymour RB, Campbell R, Pollak N, Huber G, Sharma L. Impact of the fit and strong intervention on older adults with osteoarthritis. *Gerontologist* 2004;44:217–28.
159. Fransen M, Nairn L, Winstanley J, Lam P, Edmonds J. Physical activity for osteoarthritis management: A randomized controlled clinical trial evaluating hydrotherapy or tai chi classes. *Arthritis Care Res* 2007;57:407–14.
160. Hurley MV, Walsh NE, Mitchell HL, et al. Clinical effectiveness of a rehabilitation program integrating exercise, self-management, and active coping strategies for chronic knee pain: A cluster randomized trial. *Arthritis Care Res* 2007;57:1211–9.
161. Cochrane T, Davey RC, Matthes Edwards SM. Randomised controlled trial of the cost-effectiveness of water-based therapy for lower limb osteoarthritis. *Health Technol Assess* 2005;9(31):iii–xi,1.
162. Fransen M, Crosbie J, Edmonds J. Physical therapy is effective for patients with osteoarthritis of the knee: A randomized controlled trial. *J Rheumatol* 2001;28:156–64.
163. Topp R, Woolley S, Horuyak J, Khuder S, Kahaleh B. The effect of dynamic versus isometric resistance training on pain and functioning among adults with osteoarthritis of the knee. *Arch Phys Med Rehabil* 2002;83:1187–95.
164. van Baar ME, Dekker J, Oostendorp RA, Bijl D, Voorn TB, Lemmens JA, Bijlsma JW. The effectiveness of exercise therapy in patients with osteoarthritis of the hip or knee: A randomized clinical trial. *J Rheumatol* 1998;25:2432–9.
165. Assis MR, Silva LE, Alves AMB, et al. A randomized controlled trial of deep water running: Clinical effectiveness of aquatic exercise to treat fibromyalgia. *Arthritis Care Res* 2006;55(1): 57–65.
166. Altan L, Bingol U, Aykac M, Koc Z, Yurtkuran M. Investigation of the effects of pool-based exercise on fibromyalgia syndrome. *Rheumatol Int* 2004;24(5):272–7.
167. Lemstra M, Olszynski WP. The effectiveness of multidisciplinary rehabilitation in the treatment of fibromyalgia. A randomized controlled trial. *Clin J Pain*

- 2005;21:166–74.
168. Talbot LA, Gaines JM, Huynh TN, Metter EJ. A home-based pedometer-driven walking program to increase physical activity in older adults with osteoarthritis of the knee: A preliminary study. *J Am Geriatr Soc* 2003;51:387–92.
 169. Quilty B, Tucker M, Campbell R, Dieppe P. Physiotherapy, including quadriceps exercises and patellar taping, for knee osteoarthritis with predominant patello-femoral joint involvement: Randomized controlled trial. *J Rheumatol* 2003;30:1311–7.
 170. Yip YB, Sit JWH, Fung KKY, Wong DYS, Chong SYC, Chung LH, Ng TP. Effects of a self-management arthritis programme with an added exercise component for osteoarthritic knee: Randomized controlled trial. *J Adv Nurs* 2007;59:20–8.
 171. Stener-Victorin E, Kruse-Smidje C, Jung K. Comparison between electro-acupuncture and hydrotherapy, both in combination with patient education and patient education alone, on the symptomatic treatment of osteoarthritis of the hip. *Clin J Pain* 2004;20(3):179–85.
 172. Bautch JC, Malone DG, Vailas AC. Effects of exercise on knee joints with osteoarthritis: A pilot study of biologic markers. *Arthritis Care Res* 1997;10:48–55.
 173. Thomas KS, Muir KR, Doherty M, Jones AC, O'Reilly SC, Bassey EJ. Home based exercise programme for knee pain and knee osteoarthritis: Randomised controlled trial. *BMJ* 2002;325:752–7.
 174. Cedraschi C, Desmeules J, Rapiti E, et al. Fibromyalgia: A randomised, controlled trial of a treatment programme based on self management. *Ann Rheum Dis* 2004;63:290–6.
 175. Buckelew SP, Conway R, Parker J, et al. Biofeedback/relaxation training and exercise interventions for fibromyalgia: A prospective trial. *Arthritis Care Res* 1998;11:196–209.
 176. Thorstensson CA, Roos EM, Petersson IF, Ekdahl C. Six-week high-intensity exercise program for middle-aged patients with knee osteoarthritis: A randomized controlled trial. *BMC Musculoskel. Dis* 2005;6:27.
 177. Semble EL. Rheumatoid arthritis: New approaches for its evaluation and management. *Arch Phys Med Rehabil* 1995;76: 190–201.
 178. Messier SP, Mihalko S, Loeser RF, et al. Glucosamine/chondroitin combined with exercise for the treatment of knee osteoarthritis: A preliminary study. *Osteoarthritis Cartilage* 2007;15:1256–1266.
 179. Clegg DO, Reda DJ, Harris CL, et al. Glucosamine, chondroitin sulfate, and the two in combination for painful knee osteoarthritis. *N Engl J Med* 2006;354(8):795–808.
 180. Crofford LJ, Rowbotham MC, Mease PJ, et al.; Pregabalin 1008–105 Study Group. Pregabalin for the treatment of fibromyalgia syndrome: Results of a randomized, double-blind, placebo-controlled trial. *Arthritis Rheum* 2005;52:1264–73.
 181. Rossy LA, Buckelew SP, Dorr N, et al. A meta-analysis of fibromyalgia treatment interventions. *Ann Behav Med* 1999;21: 180–91.
 182. Physicians' Desk Reference (PDR). 62nd ed. Montvale, NJ: Thomson Healthcare; 2007.
 183. Arnett FC, Edworthy SM, Bloch DA, et al. The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis. *Arthritis Rheum* 1988;31:315–24.
 184. American College of Rheumatology Subcommittee on Rheumatoid Arthritis Guidelines. Guidelines for the management of rheumatoid arthritis. 2002 update. *Arthritis Rheum* 2002;46(2): 328–46.
 185. Bradley LA, Young LD, Anderson KO, et al. Effects of psychological therapy on pain behavior of rheumatoid arthritis patients. Treatment outcome and six-month follow-up. *Arthritis Rheum* 1990;30:1105–14.
 186. Keefe FJ, Caldwell DS, Williams DA, Gil KM, Mitchell D, Robertson C. Pain coping skills training in the management of osteoarthritis knee pain: A comparative study. *Behav Ther* 1990;21:49–62.
 187. Nielson WR, Walker C, McCain GA. Cognitive behavioral treatment of fibromyalgia syndrome: Preliminary findings. *J Rheumatol* 1992;19:98–103.
 188. Goldenberg DL, Kaplan KH, Nadeau MG, Brodeur C, Smith J, Schmid CH. A controlled study of a stress-reduction, cognitive behavioral treatment program in fibromyalgia. *J Musculoskel Pain* 1994;2:53–66.
 189. White KP, Nielson WR. Cognitive-behavioral treatment of fibromyalgia syndrome: A follow-up assessment. *J Rheumatol* 1995; 22:717–21.
 190. Fuchs HA, Sergent JS. Rheumatoid arthritis: The clinical picture. In: Koopman WJ, editor. *Arthritis and Allied Conditions: A Textbook of Rheumatology*. 13th ed. Baltimore: Williams & Wilkins; 1997: 1041–70.
 191. Meenan RF, Mason JH, Anderson JJ, Guccione AA, Kazis LE. AIMS2. The content and properties of a revised and expanded Arthritis Impact Measurement Scales health status questionnaire. *Arthritis Rheum* 1992;35:1–10.
 192. Bellamy N, Campbell J, Stevens J, Pilch L, Stewart C, Mahmood Z. Validation study of a computerized version of the Western Ontario and McMaster Universities v3.0 Osteoarthritis Index. *J Rheumatol* 1997;24:2413–5.
 193. Burckhardt CS, Clark CR, Bennett RM. The fibromyalgia impact questionnaire: Development and validation. *J Rheumatol* 1991;18: 728–33.
 194. King S, Wessel J, Bhamhani Y, Maikala R, Sholter D, Maksymowych W. Validity and reliability of the 6 minute walk in persons with fibromyalgia. *J Rheumatol* 1999;26:2233–7.
 195. Chamberlain MA, Care G, Harfield B. Physiotherapy in osteoarthritis of the knees. A controlled trial of hospital versus home exercises. *International Rehabilitation Medicine* 1982;4:101–6.
 196. Sullivan T, Allegrante JP, Peterson MGE, Kovar PA, MacKenzie CR. One-year follow-up of patients with osteoarthritis of the knee who participated in a program of supervised fitness walking and supportive patient education. *Arthritis Care Res* 1998;11: 228–33.
 197. Mikesky AE, Mazzuca SA, Brandt KD, Perkins SM, Damush T, Lane KA. Effects of strength training on the incidence and progression of knee osteoarthritis. *Arthritis Care Res* 2006;55:690–9.
 198. Buckelew SP, Murray SE, Hewett JE, Johnson J, Huyser B. Self-efficacy, pain, and physical activity among fibromyalgia patients. *Arthritis Care Res* 1995;8:43–50.
 199. Bennett RM, Burckhardt CS, Clark SR, O'Reilly CA, Wiens AN, Campbell SM. Group treatment of fibromyalgia: A 6 month out-patient program. *J Rheumatol* 1996;23:521–8.
 200. Lorig K, Mazonson PD, Holman HR. Evidence suggesting that health education for self-management in patients with chronic arthritis has sustained health benefits while reducing health care costs. *Arthritis Rheum* 1993;36:439–46.
 201. Kruger JMS, Helmick CG, Callahan LF, Haddix AC. Cost-effectiveness of the Arthritis Self-Help Course. *Arch Intern Med* 1998; 158:1245–9.

Chapter 11

1. Pither C, Nicholas M, eds. *The Identification of Iatrogenic Factors in the Development of Chronic Pain Syndromes: Abnormal Treatment Behaviour?* Proceedings of the Vith World Congress on Pain. Bond M, Charlton J, Woolf C, eds. Netherlands: Elsevier Science; 1991: 429–434.
2. Wahlgren D, Atkinson JH, Epping-Jordan JE, et al. One-year follow-up of first onset low back pain. *Pain* 1997;73:213–221.
3. Bigos S, et al. *Acute low back problems in adults. Clinical Practice Guideline No. 14.* Rockville, MD: 1994 Agency for Health Care Policy and Research, Public Health Service, U.S. Department of Health and Human Services.
4. *Clinical Standards Advisory Group C. Back Pain: Report of a CSAG Committee on Back Pain.* London: His Majesty's Stationary Office; 1994.
5. Chou R, Huffman LH. Nonpharmacologic therapies for acute and chronic low back pain: A review of the evidence for an American Pain Society/American College of Physicians clinical practice guideline. *Ann Intern Med* 2007; 147(7): 492–504.
6. Philadelphia Panel evidence-based clinical practice guidelines on selected rehabilitation interventions for low back pain. *Phys Ther* 2001;81(10):1641–1674.
7. Airaksinen O, et al. European guidelines for the management of chronic nonspecific low back pain. *Eur Spine J* 2006;15(Suppl2):S192–S300.
8. van Tulder M, et al. European guidelines for the management of acute nonspecific low back pain in primary care. *Eur Spine J* 2006;15(Suppl 2):S169–S191.
9. O'Brien K, et al. Progressive resistive exercise interventions for adults living with HIV/AIDS. *Cochrane Database Syst Rev* 2004(4): CD004248.
10. Nagi SZ. Disability concepts revisited: implications for prevention. In: AM Pope, AR Tarlov, eds. *Disability in America: Towards a National Agenda for*

- Prevention. Washington, DC: Division of Health Promotion and Disease Prevention. Institute of Medicine, National Academy Press; 1991:309–327.
11. WHO. International Classification of Functioning, Disability and Health (ICF) Online. 2006 (cited September 1, 2006); Available from: www.who.int/classifications/icf/en. Accessed 10/27/2008.
 12. WHO. International Classification of Functioning, Disability and Health (ICF). 2006 (cited 2006; Available from: <http://www3.who.int/icf/onlinebrowser/icf.cfm>. Accessed 10/27/2008.
 13. Papergeorgiou A, et al. Estimating the prevalence of low back pain in the general population. Evidence from the South Manchester back pain Survey. *Spine* 1995;20:1889–1894.
 14. Andersson G. The epidemiology of spinal disorders. In: Frymoyer J, ed. *The Adult Spine: Principles and Practice*, 2nd ed. Philadelphia: Lippincott-Raven: Philadelphia; 1997:93–142.
 - 14a. Frymoyer JW, Durrett CL. The economics of spinal disorders. In: JW Frymoyer, ed. *The Adult Spine: Principles and Practice*, 2nd ed. Philadelphia: Lippincott-Raven; 1997:143–150.
 15. Walsh K, Cruddas M, Coggon D. Low back pain in eight areas of Britain. *J Epidemiol Community Health*. 1992;46:227–230.
 16. Skovron M, et al. Sociocultural factors in and back pain. A population based study in Belgian adults. *Spine* 1994;19:129–137.
 17. Mason V. The prevalence of back pain in Great Britain. In: Office of Population Censuses and Surveys, 1994, Social Survey Division. London: HMSO; 1994:1–24.
 18. Anderson J, Felson D. Factors associated with osteoarthritis of the knee in the first national Health and Nutrition Examination Survey (HANES I). Evidence for an association with overweight, race, and physical demands of work. *Am J Epidemiol* 1988;128: 179–189.
 19. Croft P, ed. *Low Back Pain*. Oxford: Radcliffe Medical Press, 1997. 20. Von Korff M, et al. An epidemiologic comparison of pain complaints. *Pain* 1988;32:173–183.
 21. Waddell G. Low back pain: A twentieth century health care enigma. *Spine* 1996;21(24):2820–2825.
 22. Cherkin D, et al. An international comparison of back surgery rates. *Spine* 1994;19:1201–1206.
 23. Jette AM, et al. Physical therapy episodes of care for patients with low back pain. *Phys Ther* 1994;74(2):101–110; discussion 110–115.
 24. Wennburg J. Practice variations and the challenge to leadership. *Spine* 1996;21:910–916.
 25. Waddell G. *The Back Pain Revolution*. Edinburgh: Churchill Livingstone; 1998.
 26. Battie M, et al. Managing low back pain: Attitudes and treatment preferences of physical therapists. *Phys Ther* 1994;74:219–226.
 27. New Zealand acute low back pain guide, and Guide in assessing psychosocial yellow flags in acute low back pain. Wellington, NZ: Accident Rehabilitation and Compensation Insurance Corporation of New Zealand and the National Health Committee; 1997.
 28. Chou R. Evidence-based medicine and the challenge of low back pain: Where are we now? *Pain Pract* 2005;5(3):153–178.
 29. Van Tulder M. Evidence-based physical therapy for low back pain: A promising future. *Dutch Journal of Physical Therapy* 1999;109: 29–32.
 30. van Eijk FA, Chavannes AW, Gubbels JW. A randomized trial of exercise therapy in patients with acute low back pain. Efficacy on sickness absence. *Spine* 1995;15:941–947.
 31. Dahan R, et al. The challenge of using the low back pain guidelines: A qualitative research. *J Eval Clin Pract* 2007;13(4): 616–620.
 32. Fullen BM, et al. Adherence of Irish general practitioners to European guidelines for acute low back pain: A prospective pilot study. *Eur J Pain* 2007;11(6):614–623.
 33. Bekkering, G.E., et al. Implementation of clinical guidelines on physical therapy for patients with low back pain: Randomized trial comparing patient outcomes after a standard and active implementation strategy. *Phys Ther* 2005;85(6):544–555.
 34. Espeland A, Baerheim A. Factors affecting general practitioners' decisions about plain radiography for back pain: Implications for classification of guideline barriers-A qualitative study. *BMC Health Serv Res* 2003;3(1):8.
 35. Armstrong MP, McDonough S, Baxter GD. Clinical guidelines versus clinical practice in the management of low back pain. *Int J Clin Pract* 2003;57(1):9–13.
 36. Koes BW, et al. Clinical guidelines for the management of low back pain in primary care: An international comparison. *Spine* 2001;26(22):2504–2513; discussion 2513–2514.
 37. Foster NE, et al. Management of nonspecific low back pain by physiotherapists in Britain and Ireland. A descriptive questionnaire of current clinical practice. *Spine* 1999;24(13):1332–1342.
 38. Bekkering GE, et al. Effect on the process of care of an active strategy to implement clinical guidelines on physiotherapy for low back pain: A cluster randomised controlled trial. *Qual Saf Health Care* 2005;14(2):107–112.
 39. Burgers JS, et al. Characteristics of effective clinical guidelines for general practice. *Br J Gen Pract* 2003;53:15–19.
 40. Grol R. Successes and failures in the implementation of evidencebased guidelines for clinical practice. *Med Care* 2001;39(8 Suppl2):II-46–II-54.
 41. Fritz JM, Cleland JA, Brennan GP. Does adherence to the guideline recommendation for active treatments improve the quality of care for patients with acute low back pain delivered by physical therapists? *Med Care* 2007;45(10):973–980.
 42. Croft P et al. Short-term physical risk indicators for new episodes of low back pain. *Spine* 1999;24:1556–1561.
 43. Sullivan MJ, Rodgers WM, Kirsch I. Catastrophizing, depression and expectancies for pain and emotional distress. *Pain* 2001; 91(1–2):147–154.
 44. Turk D. The role of demographic and psychosocial factors in transition from acute to chronic pain. In: Jensen T, Turner J, Wiesenfeld-Hallin Z, eds. *Proceedings of the 8th World Congress on Pain, Progress in Pain Research and Management*. Seattle: IASP Press; 1997:185–213.
 45. Grazier KL, et al. The frequency of occurrence, impact, and cost of musculoskeletal conditions in the United States. Chicago: American Academy of Orthopedic Surgeons; 1984.
 46. Nachemson A, ed. *Exercise, fitness and back pain*. Scientific Proceedings of the International Conference on Exercise, Fitness and Health. Champaign, IL: Human Kinetics; 1998.
 47. Fritz JM, Cleland JA, Childs JD. Subgrouping patients with low back pain: Evolution of a classification approach to physical therapy. *J Orthop Sports Phys Ther* 2007;37(6):290–302.
 48. Cleland JA, et al. Development of a clinical prediction rule for guiding treatment of a subgroup of patients with neck pain: Use of thoracic spine manipulation, exercise, and patient education. *Phys Ther* 2007;87(1):9–23.
 49. Fritz JM, et al. An examination of the reliability of a classification algorithm for subgrouping patients with low back pain. *Spine* 2006;31(1):77–82.
 50. Spitzer W, et al. Scientific approach to the assessment and management of activity-related spinal disorders. *Spine* 1987;7S:S1–S55.
 51. Selim A, et al. The importance of radiating leg pain in assessing health outcomes among patients with low back pain: Results from the Veterans Health Study. *Spine* 1998;23:470–474.
 52. Simmonds M, Lee C, Jones S. Pain distribution and physical function in patients with low back pain. In: 13th International Congress of World Confederation for Physical Therapy. Yokohama, Japan; 1999.
 53. Bekkering GE, et al. Prognostic factors for low back pain in patients referred for physiotherapy: Comparing outcomes and varying modeling techniques. *Spine* 2005;30(16):1881–1886.
 54. Willis W, ed. Introduction to the basic science of pain and headache for the clinician: physiological concepts. In: Max M, ed. *Pain 1999-An Updated Review*. Seattle: IASP Press; 1999: 561–572.
 55. International Association for the Study of Pain: Subcommittee on Taxonomy. *Pain Terms: A list with definitions and notes on usage*. *Pain* 1980;8:249–252
 56. Tillman D. *Heat response properties of unmyelinated nociceptors*. Baltimore, MD: Johns Hopkins University; 1992.
 57. Vlayen J, Linton S. Fear-avoidance and its consequences in chronic musculoskeletal pain: A state of the art. *Pain* 2000;85:317–332.
 58. Mannion AF, et al. A randomized clinical trial of three active therapies for chronic low back pain. *Spine* 1999;24(23):2435–2448.
 59. Dettori J, et al. The effects of spinal flexion and extension exercises and their associated postures in patients with acute low back pain. *Spine* 1995;20:2303–2312.
 60. Jensen M, Karoly P. Self-report scales and procedures for assessing pain in adults. In: Turk D, Mleczak R, eds. *Handbook of Pain Assessment*. D. Turk and R.

- Melzack. New York: Guilford Press; 1992:135–151.
61. Simmonds MJ, Claveau Y. Measures of pain and physical function in patients with low back pain. *Physiother Theory Pract* 1997;13: 53–65.
 62. Jette A, et al. Physical therapy episodes of care for patients with low back pain. *Phys Ther* 1994;74:101–115.
 63. Lee CE, et al. Self-reports and clinician-measured physical function among patients with low back pain: A comparison. *Arch Phys Med Rehabil* 2001;82(2):227–231.
 64. Fairbank J, et al. The Oswestry low back pain disability questionnaire. *Physiotherapy* 1980;66:271–273.
 65. Roland M, Morris R. A study of the natural history of back pain. Part I: Development of a reliable and sensitive measure of disability in low-back pain. *Spine* 1983;8(2):141–144.
 66. Ware J, Sherbourne C. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 1992;30:473–483.
 67. Simmonds MJ, et al. Physical performance tests: are they psychometrically sound and clinically useful for patients with low back pain? *Spine* 1998;23(22):2412–2421.
 68. Marras W, Wongsamm P. Flexibility and velocity of normal and impaired lumbar spine. *Arch Phys Med Rehabil* 1986;67:213–217.
 69. Simmonds M, et al. Disability prediction in patients with back pain using performance based models. In: Joint Meeting North American Spine Society and American Pain Society, 1998; Charleston, South Carolina.
 70. Simmonds MJ, Kumar S, Lechelt E. Psychological factors in disabling low back pain: causes or consequences? *Disabil Rehabil* 1996;18(4):161–168.
 71. Smeets RJ, et al. The association of physical deconditioning and chronic low back pain: A hypothesis-oriented systematic review. *Disabil Rehabil* 2006;28(11):673–693.
 72. Smeets RJ, et al. Do patients with chronic low back pain have a lower level of aerobic fitness than healthy controls? Are pain, disability, fear of injury, working status, or level of leisure time activity associated with the difference in aerobic fitness level? *Spine* 2006;31(1):90–97; discussion 98.
 73. Wittink H, et al. The association of pain with aerobic fitness in patients with chronic low back pain. *Arch Phys Med Rehabil* 2002;83: 1467–1471.
 74. Wittink H, et al. Aerobic fitness testing in patients with chronic low back pain: Which test is best? *Spine* 2000;25(13):1704–1710.
 75. Protas E. Physical activity and low back pain. In: Max M, ed. *Pain 1999—An updated review*. Seattle: IASP Press 1999:145–152.
 76. U.S. Department of Health and Human Services. *Physical Activity and Health: A report of the Surgeon General*. Atlanta: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion; 1996.
 77. Helewa A, et al. Does strengthening the abdominal muscles prevent low back pain—A randomized controlled trial. *J Rheumatol* 1999;26:1808–1815.
 78. Battie M, et al. The role of spinal flexibility in back pain complaints within industry. *Spine* 1990;15:768–773.
 79. Nelson R. *NIOSH Low back atlas of standardized tests and measures*. Springfield, VA: National Technical Information Service; 1988.
 80. Waddell G. A new clinical model for the treatment of low back pain. *Spine* 1987;12:632–644.
 81. Newton M, et al. Trunk strength testing with Iso-machines. Part 2: Experimental evaluation of the Cybex II Back testing system in normal subjects and patients with chronic low back pain. *Spine* 1993;18(7):812–824.
 82. Mayer T, Gatchel R. *Functional Restoration for Spinal Disorders: The Sports Medicine Approach*. Philadelphia: Lea and Febiger; 1988.
 83. Protas E. Aerobic exercise in the rehabilitation of individuals with chronic low back pain: a review. *Critical Reviews in Physical and Rehabilitation Medicine* 1996;8:283–295.
 84. Bronfort G, et al. Efficacy of spinal manipulation and mobilization for low back pain and neck pain: A systematic review and best evidence synthesis. *The Spine Journal* 2004;4:335–356.
 85. Clare HA, Adams R, Maher CG. A systematic review of efficacy of McKenzie therapy for spinal pain. *Aust J Physiother* 2004;50: 209–216.
 86. van der Roer N, et al. What is the most cost-effective treatment for patients with low back pain? A systematic review. *Best Pract Res Clin Rheumatol* 2005;19(4):671–684.
 87. Ostelo RW, et al. Behavioural treatment for chronic low-back pain. *Cochrane Database Syst Rev* 2005(1):CD002014.
 88. Staal JB, et al. Return-to-work interventions for low back pain: A descriptive review of contents and concepts of working mechanisms. *Sports Med* 2002;32(4):251–267.
 89. van Tulder MW, et al. Behavioral treatment for chronic low back pain: A systematic review within the framework of the Cochrane Back Review Group. *Spine* 2001;26(3):270–281.
 90. van Tulder M, et al. Exercise therapy for low back pain: A systematic review within the framework of the cochrane collaboration back review group. *Spine* 2000;25(21):2784–2796.
 91. van Tulder MW, et al. (Chronic low back pain: Exercise therapy, multidisciplinary programs, NSAID's, back schools and behavioral therapy effective; traction not effective; results of systematic reviews). *Ned Tijdschr Geneesk* 2000;144(31):1489–1494.
 92. Hancock MJ, et al. Systematic review of tests to identify the disc, SIJ or facet joint as the source of low back pain. *Eur Spine J* 2007.
 93. Liddle SD, Gracey JH, Baxter GD. Advice for the management of low back pain: A systematic review of randomised controlled trials. *Man Ther* 2007;12(4):310–327.
 94. Gulich M, et al. (Development of a guideline for rehabilitation of patients with low back pain. Phase 2: Analysis of data of the classification of therapeutic procedures). *Rehabilitation (Stuttg)* 2003;42(2):109–117.
 95. Pengel HM, Maher CG, Refshauge KM. Systematic review of conservative interventions for subacute low back pain. *Clin Rehabil* 2002;16(8):811–820.
 96. Koes BW, et al. Spinal manipulation for low back pain. An updated systematic review of randomized clinical trials. *Spine* 1996;21(24): 2860–2871; discussion 2872–2873.
 97. Engers A, et al. Patient education for low-back pain. *Cochrane Database Syst Rev* 2003(1).
 98. Baker R, Lecouturier J, Bond S. Explaining variation in GP referral rates for x-rays for back pain. *Implementation Science* 2006;1:15.
 99. Negrini S, et al. General practitioners' management of low back pain: impact of clinical guidelines in a non-English-speaking country. *Spine* 2001;26(24):2727–2733; discussion 2734.
 100. Feuerstein M, et al. Evidence-based practice for acute low back pain in primary care: Patient outcomes and cost of care. *Pain* 2006; 124(1–2):140–149.
 101. Jackson D, Llewelyn-Philips H, Klaber-Moffett J. Categorization of back pain patients using an evidence based approach. *Musculoskeletal Management* 1996;2:39–46.
 102. Malmivaara A, et al. The treatment of acute low back pain—Bed rest, exercises, or ordinary activity? *N Engl J Med* 1995;332: 351–355.
 103. Frost H, et al. A fitness programme for patients with chronic low back pain: 2-year follow-up of a randomized controlled trial. *Pain* 1998;75:273–279.
 104. Tortensen T, et al. Efficiency and costs of medical exercise therapy, conventional physiotherapy, and self-exercise in patients with chronic low back pain. *Spine* 1998;23:2616–2624.
 105. Maher C, Latimer, J, Refshauge K. Prescription of activity for low back pain: What works? *Aust J Physiother* 1999;45:121–132.
 106. Faas A, Battie M, Malmivaara A. Exercises: Which ones are worth trying for which patients and when? *Spine* 1996;21:2874–2879.
 107. Abenhaim L, et al. The role of activity in the therapeutic management of back pain. Report of the International Paris Task Force on Back Pain. *Spine*, 2000;25(4 Suppl):1S–33S.
 108. Videman T, et al. The long-term effects of physical loading and exercise lifestyles on back-related symptoms, disability, and spinal pathology among men. *Spine* 1995;20:699–709.
 109. van Tulder M, et al. Exercise therapy for low back pain (Cochrane Review). *The Cochrane Library*. Oxford: Update Software, 2000(2).
 110. van Tulder M, et al. European Guidelines for the management of acute nonspecific low back pain in primary care. In *COST B13 Working Group on Guidelines for the Management of Acute Low Back Pain in Primary Care*. European Commission, Research Directorate-General, Department of Policy, Coordination and Strategy;2004. Available at www.backpainurope.org.
 111. Keen S, et al. Individuals with low back pain: How do they view physical activity? *Fam Pract* 1999;16:39–45.

112. Friedrich M, et al. Combined exercise and motivation program: Effect on the compliance and level of disability of patients with chronic low back pain: A randomized controlled trial. *Arch Phys Med Rehabil* 1998;79(5):475–487.
113. Simmonds MJ, et al. Physical therapy assessment: Expanding the model. In: *Proceedings of the 9th World Congress on Pain. Progress on Pain. Progress in Pain Research and Management*, vol 16. DeverM, Rowbotham MC, Iesenfield-Hallin. IASP Press, Seattle: 2000; 1013–1030.

Chapter 12

1. American College of Sports Medicine. *American College of Sports Medicine's Guidelines for Exercise Testing and Prescription*. 7th ed. Baltimore (MD): Lippincott Williams & Wilkins; 2006. 366 p.
2. Anderson GL, Limacher M, Assaf AR, et al. Effects of conjugated equine estrogen in postmenopausal women with hysterectomy: the Women's Health Initiative randomized controlled trial. *JAMA*. 2004;291:1701–12.
3. Bainbridge KE, Sowers M, Lin X, et al. Risk factors for low bone mineral density and the 6-year rate of bone loss among premenopausal and perimenopausal women. *Osteoporos Int*. 2004;15:439–46.
4. Bass SL. The prepubertal years: a uniquely opportune stage of growth when the skeleton is most responsive to exercise? *Sports Med*. 2000;30:73–8.
5. Bassey EJ, Ramsdale SJ. Weight-bearing exercise and ground reaction forces: a 12-month randomized controlled trial of effects on bone mineral density in healthy postmenopausal women. *Bone*. 1995;16:469–76.
6. Beck BR, Snow CM. Bone health across the lifespan—exercising our options. *Exerc Sport Sci Rev*. 2003;31:117–22.
7. Bischoff-Ferrari HA, Dawson-Hughes B, Willett WC, et al. Effect of vitamin D on falls: a meta-analysis. *JAMA*. 2004;291:1999–2006. Bonaiuti D, Shea B, Lovine R, et al. Exercise for preventing and treating osteoporosis in postmenopausal women. *Cochrane Data-base Syst Rev*. 2002(2), CD000333.
9. Bonnick SL. Osteoporosis in men and women. *Clin Cornerstone*. 2006;8:28–39.
10. Bonnick SL, Saag KG, Kiel DP, et al. Comparison of weekly treatment of postmenopausal osteoporosis with alendronate versus risedronate over two years. *J Clin Endocrinol Metab*. 2006;91:2631–7.
11. Borer KT, Fogleman K, Gross M, et al. Walking intensity for postmenopausal bone mineral preservation and accrual. *Bone*. 2007;41:713–21.
12. Bouxsein ML, Myers ER, Hayes WC. Biomechanics of age-related fractures. In: Marcus R, Feldman D, Kelsey JL, editors. *Osteoporosis*. San Diego (CA): Academic Press, Inc; 1996. p. 373–93.
13. Briggs RC, Gossman MR, Birch R, et al. Balance performance among noninstitutionalized elderly women. *Phys Ther*. 1989;69: 748–56.
14. Burge R, Dawson-Hughes B, Solomon DH, et al. Incidence and economic burden of osteoporosis-related fractures in the United States, 2005–2025. *J Bone Miner Res*. 2007;22:465–75.
15. Canalis E, Giustina A, Bilezikian JP. Mechanisms of anabolic therapies for osteoporosis. *N Engl J Med*. 2007;357:905–16.
16. Chavassieux P, Seeman E, Delmas PD. Insights into material and structural basis of bone fragility from diseases associated with fractures: how determinants of the biomechanical properties of bone are compromised by disease. *Endocr Rev*. 2007;28: 151–64.
17. Chen P, Miller PD, Delmas PD, et al. Change in lumbar spine BMD and vertebral fracture risk reduction in teriparatide-treated post-menopausal women with osteoporosis. *J Bone Miner Res*. 2006;21:1785–90.
18. Chesnut CH, Silverman S, Andriano K, et al. A randomized trial of nasal spray salmon calcitonin in postmenopausal women with established osteoporosis: the prevent recurrence of osteoporotic fractures study. *Am J Med*. 2000;109:267–76.
19. Chrischilles EA, Butler CD, Davis CS, et al. A model of lifetime osteoporosis impact. *Arch Intern Med*. 1991;151:2026–32.
20. Chrischilles EA, Shireman T, Wallace R. Costs and health effects of osteoporotic fractures. *Bone*. 1994;15:377–85.
21. Cimen OB, Ulubas B, Sahin G, et al. Pulmonary function tests, respiratory muscle strength, and endurance of patients with osteoporosis. *South Med J*. 2003;96:423–6.
22. Cowley A, Kerr K. A review of clinical balance tools for use with elderly populations. *Crit Rev Phys Rehabil Med*. 2003;15:167–206.
23. Cranney A, Tugwell P, Zytaruk N, et al. Meta-analyses of therapies for postmenopausal osteoporosis. VI. Meta-analysis of calcitonin for the treatment of postmenopausal osteoporosis. *Endocr Rev*. 2002;23:540–51.
24. Culham EG, Jimenez HA, King CE. Thoracic kyphosis, rib mobility, and lung volumes in normal women and women with osteoporosis. *Spine*. 1994;19:1250–5.
25. Cutler WB, Friedmann E, Genovese-Stone E. Prevalence of kyphosis in a healthy sample of pre- and postmenopausal women. *Am J Phys Med Rehabil*. 1993;72:219–25.
26. Dalsky GP, Stocke KS, Ehsani AA, et al. Weight-bearing exercise training and lumbar bone mineral content in postmenopausal women. *Ann Intern Med*. 1988;108:824–8.
27. Delmas PD, Ensrud KE, Adachi JD, et al. Efficacy of raloxifene on vertebral fracture risk reduction in postmenopausal women with osteoporosis: four-year results from a randomized clinical trial. *J Clin Endocrinol Metab*. 2002;87:3609–17.
28. Ensrud KE, Black DM, Palermo L, et al. Treatment with alendronate prevents fractures in women at highest risk: results from the Fracture Intervention Trial. *Arch Intern Med*. 1997;157: 2617–24.
29. Ensrud KE, Palermo L, Black DM, et al. Hip and calcaneal bone loss increase with advancing age: longitudinal results from the study of osteoporotic fractures. *J Bone Miner Res*. 1995;10:1778–87.
30. Fehling PC, Alekel L, Clasey J, et al. A comparison of bone mineral densities among female athletes in impact loading and active loading sports. *Bone*. 1995;17:205–10.
31. Feskanich D, Willett WC, Colditz GA. Calcium, vitamin D, milk consumption, and hip fractures: a prospective study among post-menopausal women. *Am J Clin Nutr*. 2003;77:504–11.
32. Fiatarone MA, Marks EC, Ryan ND, et al. High-intensity strength training in nonagenarians: effects on skeletal muscle. *JAMA*. 1990;263:3029–34.
33. Flemming PJ. Utilization of a screening tool to identify home-bound older adults at risk for falls: validity and reliability. *Home Health Care Serv Q*. 2006;25:1–26.
34. Forsmo S, Hvam HM, Rea ML, et al. Height loss, forearm bone density and bone loss in menopausal women: a 15-year prospective study. The Nord-Trøndelag Health Study, Norway. *Osteoporos Int*. 2007;18:1261–9.
35. Gehlsen GM, Whaley MH. Falls in the elderly. Part II: Balance, strength, and flexibility. *Arch Phys Med Rehabil*. 1990;71:739–41.
36. Gennari L, Bilezikian JP. Osteoporosis in men: pathophysiology and treatment. *Curr Rheumatol Rep*. 2007;9:71–7.
37. Glynn NW, Meilahn EN, Charron M, et al. Determinants of bone mineral density in older men. *J Bone Miner Res*. 1995;10:1769–77.
38. Grey A. Emerging pharmacologic therapies for osteoporosis. *Expert Opin Emerg Drugs*. 2007;12:493–508.
39. Guralnik JM, Ferrucci L, Simonsick EM, et al. Lower-extremity function in persons over the age of 70 years as a predictor of subsequent disability. *N Engl J Med*. 1995;332:556–61.
40. Guthrie JR, Leher P, Dennerstein L, et al. The relative effect of endogenous estradiol and androgens on menopausal bone loss: a longitudinal study. *Osteoporos Int*. 2004;15:881–6.
41. Hallstrom H, Wolk A, Glynn A, et al. Coffee, tea and caffeine consumption in relation to osteoporotic fracture risk in a cohort of Swedish women. *Osteoporos Int*. 2006;17:1055–64.
42. Harridge SD, Kryger A, Stensgaard A. Knee extensor strength, activation, and size in very elderly people following strength training. *Muscle Nerve*. 1999;22:831–9.
43. Hatori M, Hasegawa A, Adachi H, et al. The effects of walking at the anaerobic threshold on vertebral bone loss in postmenopausal women. *Calcif Tissue Int*. 1993;52:411–4.
44. Hawk C, Hyland J, Rupert R, et al. Assessment of balance and risk for falls in a sample of community-dwelling adults aged 65 and older. *Chiropractic & Osteopathy*. 2006;14:3.

45. Heaney RP. Pathophysiology of osteoporosis. *Am J Med Sci.* 1996; 312:251–6.
46. Heaney RP. Effects of caffeine on bone and the calcium economy. *Food Chem Toxicol.* 2002;40:1263–70.
47. Heitmann DK, Gossman MR, Shaddeau SA, et al. Balance performance and step width in noninstitutionalized, elderly, female fallers and nonfallers. *Phys Ther.* 1989;69:923–31.
48. Heller HJ, Stewart A, Haynes S, et al. Pharmacokinetics of calcium absorption from two commercial calcium supplements. *J Clin Pharmacol.* 1999;39:1151–4.
49. Hillard TC, Whitcroft SJ, Marsh MS, et al. Long-term effects of transdermal and oral hormone replacement therapy on postmenopausal bone loss. *Osteoporos Int.* 1994;4:341–8.
50. Hornbrook MC, Stevens VJ, Wingfield DJ. Seniors' program for injury control and education. *J Am Geriatr Soc.* 1993;41: 309–14.
51. Hundrup YA, Ekholm O, Hoidrup S, et al. Risk factors for hip fracture and a possible effect modification by hormone replacement therapy. The Danish nurse cohort study. *Eur J Epidemiol.* 2005;20:871–7.
52. Jones G, Nguyen T, Sambrook P, et al. Progressive loss of bone in the femoral neck in elderly people: longitudinal findings from the Dubbo osteoporosis epidemiology study. *BMJ.* 1994;309:691–5.
53. Kanis JA, Johansson H, Johnell O, et al. Alcohol intake as a risk factor for fracture. *Osteoporos Int.* 2005;16:737–42.
54. Kanis JA, Johnell O, Oden A, et al. Smoking and fracture risk: a meta-analysis. *Osteoporos Int.* 2005;16:155–62.
55. Kannus P, Haapasalo H, Sankelo M, et al. Effect of starting age of physical activity on bone mass in the dominant arm of tennis and squash players. *Ann Intern Med.* 1995;123:27–31.
56. Khovidhunkit W, Shoback DM. Clinical effects of raloxifene hydrochloride in women. *Ann Intern Med.* 1999;130:431–9.
57. Kiel DP, Felson DT, Hannan MT, et al. Caffeine and the risk of hip fracture: the Framingham Study. *Am J Epidemiol.* 1990;132:675–84.
58. King AC, Pruitt LA, Phillips W, et al. Comparative effects of two physical activity programs on measured and perceived physical functioning and other health-related quality of life outcomes in older adults. *J Gerontol A Biol Sci Med Sci.* 2000;55:M74–M83.
59. Kirwan LD, MacLusky NJ, Shapiro HM, et al. Acute and chronic effects of hormone replacement therapy on the cardiovascular system in healthy postmenopausal women. *J Clin Endocrinol Metab.* 2004;89:1618–29.
60. Knopp JA, Diner BM, Blitz M, et al. Calcitonin for treating acute pain of osteoporotic vertebral compression fractures: a systematic review of randomized, controlled trials. *Osteoporos Int.* 2005;16: 1281–90.
61. Kohrt WM. Aging and the osteogenic response to mechanical loading. *Int J Sport Nutr Exerc Metab.* 2001;11(Suppl):S137–S142.
62. Kohrt WM, Bloomfield SA, Little KD, et al. American College of Sports Medicine Position Stand: physical activity and bone health. *Med Sci Sports Exerc.* 2004;36:1985–96.
63. Kohrt WM, Snead DB, Slatopolsky E, et al. Additive effects of weight-bearing exercise and estrogen on bone mineral density in older women. *J Bone Miner Res.* 1995;10:1303–11.
64. Korpelainen R, Keinanen-Kiukkaanniemi S, Heikkinen J, et al. Effect of impact exercise on bone mineral density in elderly women with low BMD: a population-based randomized controlled 30-month intervention. *Osteoporos Int.* 2006;17: 109–18.
65. Kung AW, Pasion EG, Sofiyan M, et al. A comparison of teriparatide and calcitonin therapy in postmenopausal Asian women with osteoporosis: a 6-month study. *Curr Med Res Opin.* 2006;22:929–37.
66. LaPier TK. Chest wall expansion values in supine and standing across the adult lifespan. *Physical and Occupational Therapy in Geriatrics.* 2002;21:65–81.
67. Leslie WD, Adler RA, El Hajj FG, et al. Application of the 1994 WHO classification to populations other than postmenopausal Caucasian women: the 2005 ISCD Official Positions. *J Clin Densitom.* 2006;9:22–30.
68. Levine JP. Pharmacologic and nonpharmacologic management of osteoporosis. *Clin Cornerstone.* 2006;8:40–53.
69. Lombardi I, Oliveira LM, Monteiro CR, et al. Evaluation of physical capacity and quality of life in osteoporotic women. *Osteoporos Int.* 2004;15:80–5.
70. Looker AC, Orwoll ES, Johnston CC, Jr., et al. Prevalence of low femoral bone density in older U.S. adults from NHANES III. *J Bone Miner Res.* 1997;12:1761–8.
71. Lynch NA, Ryan AS, Evans J, et al. Older elite football players have reduced cardiac and osteoporosis risk factors. *Med Sci Sports Exerc.* 2007;39:1124–30.
72. McClung M, Clemmesen B, Daifotis A, et al. Alendronate prevents postmenopausal bone loss in women without osteoporosis. A double-blind, randomized, controlled trial. *Ann Intern Med.* 1998;128:253–61.
73. McClung MR, Geusens P, Miller PD, et al. Effect of risedronate on the risk of hip fracture in elderly women. Hip Intervention Program Study Group. *N Engl J Med.* 2001;344:333–40.
74. Mortensen L, Charles P, Bekker PJ, et al. Risedronate increases bone mass in an early postmenopausal population: two years of treatment plus one year of follow-up. *J Clin Endocrinol Metab.* 1998;83:396–402.
75. Mukamal KJ, Robbins JA, Cauley JA, et al. Alcohol consumption, bone density, and hip fracture among older adults: the cardiovascular health study. *Osteoporos Int.* 2007;18:593–602.
76. National Osteoporosis Foundation. *Boning up on osteoporosis: A guide to prevention and treatment.* 2000. Washington, DC: National Osteoporosis Foundation.
77. National Osteoporosis Foundation. *America's bone health: The state of osteoporosis and low bone mass.* 2002. Washington, DC: National Osteoporosis Foundation.
78. Nelson ME, Fiatarone MA, Morganti CM, et al. Effects of high-intensity strength training on multiple risk factors for osteoporotic fractures. A randomized controlled trial. *JAMA.* 1994;272: 1909–14.
79. Nelson ME, Rejeski WJ, Blair SN, et al. Physical activity and public health in older adults: recommendation from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc.* 2007;39:1435–45.
80. Nguyen ND, Eisman JA, Nguyen TV. Anti-hip fracture efficacy of bisphosphonates: a Bayesian analysis of clinical trials. *J Bone Miner Res.* 2006;21:340–9.
81. Nguyen TV, Kelly PJ, Sambrook PN, et al. Lifestyle factors and bone density in the elderly: implications for osteoporosis prevention. *J Bone Miner Res.* 1994;9:1339–46.
82. Nichols DL, Sanborn CF, Bonnick SL, et al. The effects of gymnastics training on bone mineral density. *Med Sci Sports Exerc.* 1994;26:1220–5.
83. NIH Consensus Development Panel on Osteoporosis Prevention DaT. *Osteoporosis Prevention, Diagnosis, and Therapy.* *JAMA.* 2001;285:785–95.
84. Nikander R, Sievanen H, Heinonen A, et al. Femoral neck structure in adult female athletes subjected to different loading modalities. *J Bone Miner Res.* 2005;20:520–8.
85. Nordstrom A, Karlsson C, Nyquist F, et al. Bone loss and fracture risk after reduced physical activity. *J Bone Miner Res.* 2005;20:202–7.
86. Nordstrom A, Olsson T, Nordstrom P. Sustained benefits from previous physical activity on bone mineral density in males. *J Clin Endocrinol Metab.* 2006;91:2600–4.
87. Pedersen BK, Saltin B. Evidence for prescribing exercise as therapy in chronic disease. *Scand J Med Sci Sports.* 2006;16(Suppl 1):3–63.
88. Perell KL, Nelson A, Goldman RL, et al. Fall risk assessment measures: an analytic review. *J Gerontol A Biol Sci Med Sci.* 2001;56: M761–M766.
89. Prince RL, Smith M, Dick IM, et al. Prevention of postmenopausal osteoporosis: a comparative study of exercise, calcium supplementation, and hormone-replacement therapy. *N Engl J Med.* 1991;325:1189–95.
90. Province MA, Hadley EC, Hornbrook MC, et al. The effects of exercise on falls in elderly patients. A preplanned meta-analysis of the FICSIT Trials. *Frailty and Injuries: Cooperative Studies of Intervention Techniques.* *JAMA.* 1995;273:1341–7.
91. Rapuri PB, Gallagher JC, Balhorn KE, et al. Smoking and bone metabolism in elderly women. *Bone.* 2000;27:429–36.
92. Reid IR, Wattie DJ, Evans MC, et al. Continuous therapy with pamidronate, a potent bisphosphonate, in postmenopausal osteoporosis. *J Clin Endocrinol Metab.* 1994;79:1595–9.
93. Rhodes EC, Martin AD, Taunton JE, et al. Effects of one year of resistance training on the relation between muscular strength and bone density in elderly women. *Br J Sports Med.* 2000;34:18–22.
94. Ringe JD, Dorst A, Faber H, Ibach K. Alendronate treatment of established primary osteoporosis in men: 3-year results of a prospective, comparative, two-arm

- study. *Rheumatol Int.* 2004;24:110–3.
95. Ringe JD, Faber H, Farahmand P, et al. Efficacy of risedronate in men with primary and secondary osteoporosis: results of a 1-year study. *Rheumatol Int.* 2006;26:427–31.
 96. Ringe JD, van der Geest SA, Moller G. Importance of calcium co-medication in bisphosphonate therapy of osteoporosis: an approach to improving correct intake and drug adherence. *Drugs Aging.* 2006;23:569–78.
 97. Rodan GA. Introduction to bone biology. *Bone.* 1992;13:S3–S6.
 98. Rollins D, Imrhan V, Czajka-Narins DM, et al. Lower bone mass detected at femoral neck and lumbar spine in lower-weight vs normal-weight small-boned women. *J Am Diet Assoc.* 2003;103:742–4.
 99. Rossouw JE, Anderson GL, Prentice RL, et al. Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results From the Women's Health Initiative randomized controlled trial. *JAMA.* 2002;288:321–33.
 100. Rubenstein LZ. Falls in older people: epidemiology, risk factors and strategies for prevention. *Age Ageing.* 2006;35(Suppl 2): ii37–ii41.
 101. Rubenstein LZ, Josephson KR. Falls and their prevention in elderly people: what does the evidence show? *Med Clin North Am.* 2006;90:807–24.
 102. Sawka AM, Papaioannou A, Adachi JD, et al. Does alendronate reduce the risk of fracture in men? A meta-analysis incorporating prior knowledge of anti-fracture efficacy in women. *BMC Musculoskelet Disord.* 2005;6:39.
 103. Schlaich C, Minne HW, Bruckner T, et al. Reduced pulmonary function in patients with spinal osteoporotic fractures. *Osteoporos Int.* 1998;8:261–7.
 104. Seeman E, Crans GG, Diez-Perez A, et al. Anti-vertebral fracture efficacy of raloxifene: a meta-analysis. *Osteoporos Int.* 2006;17: 313–6.
 105. Shea B, Wells G, Cranney A, et al. Calcium supplementation on bone loss in postmenopausal women. *Cochrane Database Syst Rev.* 2004CD004526.
 106. Sinaki M. Postmenopausal spinal osteoporosis: physical therapy and rehabilitation principles. *Mayo Clin Proc.* 1982;57:699–703.
 107. Sinaki M, Mikkelsen BA. Postmenopausal spinal osteoporosis: flexion versus extension exercises. *Arch Phys Med Rehab.* 1984; 65:593–6.
 108. Sinaki M, Wollan PC, Scott RW, et al. Can strong back extensors prevent vertebral fractures in women with osteoporosis? *Mayo Clin Proc.* 1996;71:951–6.
 109. Siris ES, Miller PD, Barrett-Connor E, et al. Identification and fracture outcomes of undiagnosed low bone mineral density in post-menopausal women: results from the National Osteoporosis Risk Assessment. *JAMA.* 2001;286:2815–22.
 110. Snow CM. Exercise and bone mass in young premenopausal women. *Bone.* 1996;18:51S–5S.
 111. Sowers M. Pregnancy and lactation as risk factors for subsequent bone loss and osteoporosis. *J Bone Miner Res.* 1996;11: 1052–60.
 112. Suominen H. Muscle training for bone strength. *Aging Clin Exp Res.* 2006;18:85–93.
 113. Tang BM, Eslick GD, Nowson C, et al. Use of calcium or calcium in combination with vitamin D supplementation to prevent fractures and bone loss in people aged 50 years and older: a meta-analysis. *Lancet.* 2007;370:657–66.
 114. Tansavatdi K, McClain B, Herrington DM. The effects of smoking on estradiol metabolism. *Minerva Ginecol.* 2004;56: 105–14.
 115. Thiebaud D, Burckhardt P, Kriegbaum H, et al. Three monthly in- travenous injections of ibandronate in the treatment of postmenopausal osteoporosis. *Am J Med.* 1997;103:298–307.
 116. Trivedi DP, Doll R, Khaw KT. Effect of four monthly oral vitamin D3 (cholecalciferol) supplementation on fractures and mortality in men and women living in the community: randomised doubleblind controlled trial. *BMJ.* 2003;326:469.
 117. Trudelle-Jackson EJ, Jackson AW, Morrow J. Muscle strength and postural stability in healthy, older women: implications for fall prevention. *J Phys Act Health.* 2006;3:
 118. Tucker KL, Morita K, Qiao N, et al. Colas, but not other carbon-ated beverages, are associated with low bone mineral density in older women: The Framingham Osteoporosis Study. *Am J Clin Nutr.* 2006;84:936–42.
 119. Tudor-Locke C, McColl RS. Factors related to variation in pre- menopausal bone mineral status: a health promotion approach. *Osteoporos Int.* 2000;11:1–24.
 120. Uusi-Rasi K, Kannus P, Cheng S, et al. Effect of alendronate and exercise on bone and physical performance of postmenopausal women: a randomized controlled trial. *Bone.* 2003;33:132–43.
 121. Valimaki MJ, Laitinen KA, Tahtela RK, et al. The effects of transdermal estrogen therapy on bone mass and turnover in early postmenopausal smokers: a prospective, controlled study. *Am J Obstet Gynecol.* 2003;189:1213–20.
 122. Vestergaard P, Rejnmark L, Mosekilde L. Fracture reducing poten- tial of hormone replacement therapy on a population level. *Maturitas.* 2006;54:285–93.
 123. Weir JP, Cramer JT. Principles of musculoskeletal exercise programming. In: American College of Sports Medicine, editors. *ACSM's Resource Manual for Guidelines for Exercise Testing and Prescription*, Baltimore (MD): Lippincott Williams & Wilkins; 2006. p. 350–365.
 124. Wells G, Tugwell P, Shea B, et al. Meta-analyses of therapies for postmenopausal osteoporosis. V. Meta-analysis of the efficacy of hormone replacement therapy in treating and preventing osteoporosis in postmenopausal women. *Endocr Rev.* 2002;23:529–39.
 125. Whipple RH, Wolfson LI, Amerman PM. The relationship of knee and ankle weakness to falls in nursing home residents: an isokinetic study. *J Am Geriatr Soc.* 1987;35:13–20.
 126. Wickham C, Cooper C, Margetts BM, et al. Muscle strength, activ- ity, housing and the risk of falls in elderly people. *Age Ageing.* 1989;18:47–51.
 127. Wimalawansa SJ. A four-year randomized controlled trial of hormone replacement and bisphosphonate, alone or in combination, in women with postmenopausal osteoporosis. *Am J Med.* 1998; 104:219–26.
 128. World Health Organization. *Assessment of Fracture Risk and Its Application to Screening for Postmenopausal Osteoporosis*. Geneva: World Health Organization; 1994.
 129. Wosje KS, Kalkwarf HJ. Bone density in relation to alcohol intake among men and women in the United States. *Osteoporos Int.* 2007;18:391–400.

Chapter 13

1. Manek NJ, MacGregor AJ. Epidemiology of back disorders: prevalence, risk factors, and prognosis. *Curr Opin Rheumatol* 2005;17: 134–40.
2. Andersson GB. Epidemiology of low back pain. *Acta Orthop Scand Suppl* 1998;281:28–31.
3. Frymoyer JW, Cats-Buril WL. An overview of the incidences and cost of low back pain. *Orthop Clin North Am* 1991;22:263–71.
4. Walker BF. The prevalence of low back pain: A systematic review of the literature 1966 to 1998. *J Spinal Disor* 2000;13:205–17.
5. Hestbaek L, Leboef-Yde C, Manniche C. Low back pain: what is the long term cause? A review of studies of general patient populations. *Eur Spine.* 2003;12:149–65.
6. Troup JD, Martin JW, Lloyd DC. Back pain in industry: A prospective survey. *Spine* 1981;6:61–9.
7. Anderson RA. A case study in integrating medicine: alternate theories and the language of biomedicine. *J Altern Complement Med* 1999;5:165–73.
8. Frymoyer JW. Can low back pain disability be prevented? *Baillieres Clin Rheumo* 1992;6:595–606.
9. National Institute of Neurological Disorders and Stroke. Web site [Internet]. NIH, Bethesda, MD: National Institute of Neurological Disorders and Stroke. Low back pain fact sheet; [cited 2007 Aug 12]. Available from: http://www.ninds.nih.gov/disorders/backpain/detail_backpain.htm
10. Center for Disease Control and Prevention. Prevalence of disabilities and associated health conditions among adults: United States, 1999. *MMWR* 2001;50:120–5.
11. World Health Organization Web site [Internet]. Geneva: Towards a Common language for Functioning, Disability and Health ICF; 2002. Available from: <http://www.who.int/classifications/icf/site/beginners/bg.pdf>.
12. Krismser M, van Tulder M, The Low Back Pain Group of the Bone and Joint Health Strategies for Europe Project. Low back pain (nonspecific). *Best Pract Res Clin Rheumatol* 2007;21:77–91.
13. Nelemans PJ, deBie RA, deVet HCW, Sturmans F. Injection therapy for subacute and chronic benign low back pain. *Cochrane Database Syst Rev* 2000(2):CD001824.
14. Bogduk N. *Clinical Anatomy of the Lumbar Spine and Sacrum*, 3rd ed. New York: Churchill Livingstone; 1997:187–213.
15. Kuslich SD, Ulstrom CL, Michael CJ. The tissue origin of low back pain and sciatica: a report of pain response to tissue stimulation during operations on the

- lumbar spine using local anesthesia. *Orthop Clin North Am* 1991;22:181–7.
16. Schwarzer, AC, Aprill CN, Derby R, et al. The relative contributions of the disc and zygapophyseal joint in chronic low back pain. *Spine* 1994;19:801–6.
 17. Jönsson B, Strömquist B. Clinical appearance of contained and noncontained lumbar disc herniation. *J Spinal Disord* 1996;9:32–8.
 18. Sahrman SA. *Diagnosis and Treatment of Movement Impairment Syndromes*. St. Louis: Mosby; 2001:6.
 19. Kirkaldy-Willis WH. Three phases of the spectrum of degenerative disease. In: Kirkaldy-Willis WH, Burton CV, eds. *Managing Low Back Pain*, 3rd ed. New York: Churchill-Livingstone; 1992: 105–19.
 20. Waddell G. *The Back Pain Revolution*. Edinburgh: Churchill Livingstone; 1998:9–25.
 21. Hurwitz EL, Morgenstern H. Correlates of back problems and back related disability in the United States. *J Clin Epidemiol* 1997;50: 669–81.
 22. Kopec JH, Sayre EC, Esclaile JM. Predictors of back pain in a general population cohort. *Spine* 2004;29:70–7.
 23. Manchikanti L. Epidemiology of low back pain. *Pain Physician* 2000;3:167–92.
 24. Waddell G, Newton M, Henderson I, et al. A Fear-Avoidance Beliefs Questionnaire (FABQ) and the role of fear-avoidance beliefs in chronic low back pain and disability. *Pain* 1993;52:157–68.
 25. Beck AT, Ward CH, Mendelson M, et al. An inventory for measuring depression. *Arch Gen Psychiatry* 1961;4:53–63.
 26. DeRosa CP, Porterfield JA. A physical therapy model for the treatment of low back pain. *Phys Ther* 1992;72:261–9.
 27. American Physical Therapy Association. *Guide to physical therapist practice*, 2nd ed. *Phys Ther* 2001;81:9–744.
 28. Cassidy JD, Carroll LJ, Cote P. The Saskatchewan health and back pain survey. The prevalence of low back pain and related disability in Saskatchewan adults. *Spine* 1998;23:1860–6.
 29. Abenhaim L, Rossignol M, Gobeille D, et al. The prognostic consequences of making the initial medical diagnosis of work-related back injuries. *Spine* 1995;20:791–5.
 30. Lawrence RC, Helmick CG, Arnett FC, et al. Estimates of the prevalence of arthritis and selected musculoskeletal disorders in the United States. *Arthritis Rheum* 1998;41:778–99.
 31. Chou R, Qaseem A, Snow V, et al. Diagnosis and treatment of low back pain: A joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Ann Intern Med* 2007;147:478–91.
 32. Eliasson K, Elfving B, Nordgren B, et al. Urinary incontinence in women with low back pain. *Man Ther* 2008;13:206–12.
 33. Fairbank JC, Couper J, Davies JB, et al. The Oswestry low back pain disability questionnaire. *Physiotherapy* 1980;66:271–7.
 34. Roland M, Morris R. A study of the natural history of back pain. Part I: development of a reliable and sensitive measure of disability in low back pain. *Spine* 1983;8:141–4.
 35. Stratford P, Gill C, Westaway M, et al. Assessing disability and change on individual patients: A report of a patient specific measure. *Physiother Can* 1995;47:258–63.
 36. Sahrman SA. *Diagnosis and Treatment of Movement Impairment Syndromes*. St. Louis: Mosby; 2001:9–50.
 37. Meadows JTS. *Orthopedic Differential Diagnosis in Physical Therapy: A Case Study Approach*. New York: McGraw-Hill; 1999:4.
 38. Panjabi MM. The stabilizing system of the spine. Part II: Neutral zone and instability hypothesis. *J Spinal Dis* 1992;5:390–7.
 39. Sahrman SA. *Diagnosis and Treatment of Movement Impairment Syndromes*. St. Louis: Mosby; 2001:51–119.
 40. Pye SR, Reid DM, Smith R, et al. Radiographic features of lumbar disc degeneration and self-reported back pain. *J Rheumatol* 2004; 31:753–8.
 41. Scientific approach to the assessment and management of activity-related spinal disorders. A monograph for clinicians. Report of the Quebec Task Force on Spinal Disorders. *Spine* 1987;12:S1–59.
 42. Fritz JM, Cleland JA, Childs JD. Subgrouping patients with low back pain: Evolution of a classification approach to physical therapy. *J Orthop Sports Phys Ther* 2007;37:290–302.
 43. Delitto A, Erhard RE, Bowling RW. A treatment-based classification approach to low back syndrome: Identifying and staging patients for conservative treatment. *Phys Ther* 1995;75:470–85.
 44. Childs JD, Fritz JM, Flynn TW, et al. A clinical prediction rule to identify patients with low back pain most likely to benefit from spinal manipulation: A validation study. *Ann Intern Med* 2004;141: 920–8.
 45. Flynn T, Fritz J, Whitman J, et al. A clinical prediction rule for classifying patients with low back pain who demonstrate short-term improvement with spinal manipulation. *Spine* 2002;27:2835–43.
 46. Hicks GE, Fritz JM, Delitto A, et al. Preliminary development of a clinical prediction rule for determining which patients with low back pain will respond to a stabilization exercise program. *Arch Phys Med Rehabil* 2005;86:1753–62.
 47. Kane RL. Looking for physical therapy outcomes. *Phys Ther* 1994;74: 425–9.
 48. van Tulder MV, Scholten RJ, Koes BW, et al. Nonsteroidal anti-inflammatory drugs for low back pain: A systematic review within the Cochrane Collaboration Back Review Group. *Spine* 2000;25:2501–13.
 49. Cherkin DC, Wheeler KJ, Barlow W, et al. Medication use for low back pain in primary care. *Spine* 1998;23:607–14.
 50. Philadelphia Panel evidenced-based clinical practice guidelines on selected rehabilitation interventions for low back pain. *Phys Ther* 2001;81:1641–74.
 51. Greenman PE. *Principles of Manual Medicine*. Baltimore: Williams & Wilkins; 1989:88–93.
 52. Klaber Moffett JA, Carr J, Howarth E. High fear-avoiders of physical activity benefit from an exercise program for patients with back pain. *Spine* 2004;29:1167–72.
 53. Bigos S, Bower O, Braen G, et al. *Acute Low Back Pain in Adults*. US Department of Health and Human Services. Public Health Service. Agency for Health Care Policy and Research Guideline No. 14. AHCPR Publication No. 97-N012. Rockville, MD, February 1997.
 54. Waddell G, Feder G, Lewis M. Systematic reviews of bed rest and advice to stay active for acute low back pain. *Br J Gen Pract* 1997; 47:647–52.
 55. Deyo RA, Diehl AK, Rosenthal M. How many days of bed rest for acute low back pain? A randomized clinical trial. *N Engl J Med* 1986;23:1064–70.
 56. Hodges PW, Richardson CA. Contraction of the abdominal muscles associated with movement of the lower limb. *Phys Ther* 1997;77: 132–42.
 57. Hodges PW, Richardson CA. Feedforward contraction of the transversus abdominis is not influenced by the direction of arm movement. *Exp Brain Res* 1997;114:362–70.
 58. Sihvonen T, Lindgren KA, Airaksinen O, et al. Movement disturbances of the lumbar spine and abnormal back muscle electromyographic findings in recurrent low back pain. *Spine* 1997;22:289–95.
 59. Hides JA, Richardson CA, Jull GA. Multifidus muscle recovery is not automatic after resolution of acute, first-episode low back pain. *Spine* 1996;21:2763–9.
 60. Neumann P, Gill V. Pelvic floor and abdominal muscle interaction: EMG activity and intra-abdominal pressure. *Int Urogynecol J* 2002; 13:125–32.
 61. Richardson C, Jull G, Hodges P, et al. *Therapeutic Exercise for Spinal Segmental Stabilization in Low Back Pain: Scientific Basis and Clinical Approach*. Edinburgh: Churchill Livingstone; 1999:17.
 62. Richardson C, Jull G, Hodges P, et al. *Therapeutic Exercise for Spinal Segmental Stabilization in Low Back Pain: Scientific Basis and Clinical Approach*. Edinburgh: Churchill Livingstone; 1999:105–23.
 63. The mainstreaming of alternative medicine. *Consumer Reports*. May 2000:17–25.
 64. van Tulder MV, Malmivara A, Esmail R, et al. Exercise therapy for low back pain (Cochrane Review). In: *The Cochrane Library*, Issue 2, 2001. Oxford: Update Software.
 65. McKenzie RA, May S. *The Lumbar Spine: Mechanical Diagnosis and Therapy*. Vol. 1. Waikanae, New Zealand: Spinal Publications New Zealand Ltd; 2003:139–80.
 66. Fritz JM, George S. The use of a classification approach to identify subgroups of patients with acute low back pain. Interrater reliability and short-term treatment outcomes. *Spine* 2000;25:106–14.
 67. Donelson R, Silva G, Murphy K. Centralization phenomenon. Its usefulness in evaluating and treating referred pain. *Spine* 1990;15:211–3.

68. Sahrman SA. *Diagnosis and Treatment of Movement Impairment Syndromes*. St. Louis: Mosby; 2001:4.
69. Richardson CA, Jull GA. Muscle control–pain control. What exercises would you prescribe? *Man Ther* 1995;1:2–10.
70. Gill KP, Callaghan MJ. The measurement of lumbar proprioception in individuals with and without low back pain. *Spine* 1998;23: 371–7.
71. McGill SM. Low back exercises: Evidence for improving exercise regimens. *Spine* 1998;78:754–65.
72. Cady LD, Bischoff DP, O’Connell ER, et al. Strength and fitness and subsequent back injuries in firefighters. *J Occup Med* 1979;21: 269–72.
73. van Tulder MV, Koes BW, Bouter LM. Conservative treatment of acute and chronic nonspecific low back pain. A systematic review of randomized controlled trials of the most common interventions. *Spine* 1997;22:2128–56.
74. Rubin DI. Epidemiology and risk factors for spine pain. *Neurol Clin* 2007;25:353–71.

Chapter 14

1. Dillingham TR, Pezzin LE, Mackenzie EJ. Limb amputations and limb deficiency: Epidemiology and recent trends in the United States. *South Med J* 2002;95:875–883.
2. Gonzalez EG, Edelstein JE. Energy expenditure during ambulation. In: Gonzalez EG, Myers SJ, Edelstein JE, et al., eds. *Downey & Darling’s Physiological Basis of Rehabilitation Medicine*. Boston: Butterworth Heinemann; 2001:413–446
3. Gailey RS, Wenger MA, Raya M et al. Energy expenditure of transtibial amputees during ambulation at self-selected pace. *Prosthet Orthot Int* 1994;18:84–91.
4. Jaegers SMHJ, Vos LD, Rispens P, et al. The relationship between comfortable and most metabolically efficient walking speed in person with unilateral above-knee amputation. *Arch Phys Med Rehabil* 1993;74:521–525.
5. Nowroozi F, Satonelli MI, Gerber LH. Energy expenditure in hip disarticulation and hemipelvectomy. *Arch Phys Med Rehabil* 1983; 64:300–303.
6. Waters RL, Mulroy SJ. Energy expenditure of walking in individuals with lower limb amputation. In: Smith DG, Michael JW, Bowker JH, eds. *Atlas of Amputations and Limb Deficiencies*. Rosemont, IL: American Academy of Orthopedic Surgeons; 2004:400–403.
7. Pitetti KH, Snell PG, Stray-Gundersen J. Aerobic training exercise for individuals who had amputation of the lower limb. *J Bone Joint Surg* 1987;69:914–921.
8. Hrubec Z, Ryder RA. Traumatic limb amputation and subsequent mortality from cardiovascular disease and other causes. *J Chronic Dis* 1978;33:239–250.
9. Rose HC, Schweitzer P, Charoenkul V, et al. Cardiovascular disease risk factors in combat veterans after traumatic leg amputation. *Arch Phys Med Rehabil* 1987;68:20–23.
10. Braune W, Fischer O. *On the Center of Gravity of the Human Body*. Berlin, Germany: Springer-Verlag; 1984.
11. Kusljagic A, Kapidzic-Durakovic S, Kudumovic Z, et al. Chronic low back pain in individuals with lower-limb amputation. *Bosn J Basic Med Sci* 2006;6(2):67–70.
12. Gailey RS, Gailey AM. *Stretching and Strengthening for Lower Extremity Amputees*. Miami, FL: Advanced Rehabilitation Therapy; 1994. (Correspondence: Advanced Rehabilitation Therapy, Inc., 7641 SW 126th Street, Miami, FL 33156.)
13. Gailey RS, Gailey AM. *Balance, Agility, Coordination and Endurance for Lower Extremity Amputees*. Miami, FL: Advanced Rehabilitation Therapy; 1994.
14. Gailey RS, Gailey AM, Sendelbach SJ. *Home Exercise Guide for Lower Extremity Amputees*. Miami, FL: Advanced Rehabilitation Therapy; 1995.
15. American College of Sports Medicine. *ACSM’s Exercise Management for Persons with Chronic Diseases and Disabilities*. Champaign, IL: Human Kinetics; 2003:374.
16. Meulenbelt HEJ, Geertzen JHB, Kijkstra PU, et al. Skin problems in lower limb amputees: An overview by case reports. *J Eur Acad Derm Venereol* 2007;21:147–155.
17. Edelstein JE. Amputations and prostheses. In: Cameron MH, Monroe LG, eds. *Physical Rehabilitation. Evidence-Based Examination, Evaluation, and Intervention*. Philadelphia: Saunders-Elsevier; 2007:267–299.
18. Kendall FP, McCreary EK, Provance PG, et al. *Muscles: Testing and Function*. 5th ed. Baltimore, MD: Lippincott Williams & Wilkins; 2005:4–47.
19. Bowler JH, Michael JW. *Atlas of Limb Prosthetics: Surgical, Prosthetic and Rehabilitation Principles*. American Academy of Orthopaedic Surgeons. St. Louis, MO: Mosby-Year Book; 1992:930.
20. American College of Sports Medicine. *ACSM’s Guidelines for Exercise Testing and Prescription*. 6th ed. Philadelphia: Lippincott Williams & Wilkins; 2000.
21. Priebe M, Davidoff G, Lampman RM. Exercise testing and training in patients with peripheral vascular disease and lower extremity amputation. *West J Med* 1991;154:598–601.
22. James U. Effect of physical training in healthy male unilateral above-knee amputees. *Scand J Rehabil Med* 1973;5:88–101.
23. Davidoff G, Lampman R, Westbury L, et al. Exercise testing and training of dysvascular amputees: Safety and efficacy of arm ergometry. *Arch Phys Med Rehabil* 1992; 73:334–338.
24. Burgess EM, Rappoport A. *Physical Fitness: A Guide for Individuals with Lower Limb Loss*. Washington, DC: Department of Veterans Affairs; 1991:245.
25. Kegel B. Physical fitness: Sports and recreation for those with lower limb amputation or impairment. *J Rehabil Res Dev Clin Suppl* 1985; 1:1–125.
26. Machin P, de C Williams AC. Stiff upper lip: Coping strategies of World War II veterans with phantom limb pain. *Clin J Pain* 1998; 14(4):290–294.
27. Wilkins KL, McGrath PJ, Finley GA, et al. Phantom limb sensations and phantom limb pain in child and adolescent amputees. *Pain* 1998;78(1):7–12.
28. Wartan SW, Hamann W, Wedley JR, et al. Phantom pain and sensation among British veterans amputees. *Br J Anaesth* 1997;78(6):652–659.
29. Schmid HJ. Phantom limb after amputation—Overview and new knowledge. *Schweiz Rundsch Med Prax* 2000;89(3):87–94.
30. Lindsay JE. Multiple pain complaints in amputees. *J R Soc Med* 1985;78(6):452–455.