

1. Boyle P, Lewin B. (2008) World Cancer Report 2008. International Agency for Research on Cancer, France.
2. National Cancer Institute, USA. (2009) <http://www.cancer.gov/cancertopics/wyntk/Overview/page2>. Accessed 25 January 2009.
3. Garcia M, Jemal A, Ward EM, C-eter MM, Hao Y, Siegal RL,, Thun MJ. (2007) Global Cancer Facts & Figures. American Cancer Society, Atlanta, GA.
4. Coleman MP, Quaresma M, Berrino F, et al. (2008) Cancer survival in five continents: a worldwide population-based study (CONCORD). *Lancet Oncol* 9:730–756.
5. Twombly R. (2004) What's in a name: who is a cancer survivor? *J Natl Cancer Inst* 96: 1414–1415.
6. Aziz NM. (2002) Cancer survivorship research: challenge and opportunity. *J Nutr* 132:3494S–3503S.
7. Hewitt M, Rowland JH, Yancik R. (2003) Cancer survivors in the united states: age, health, and disability. *J Gerontol A Biol Sci Med Sci* 58:82–91.
8. Demark-Wahnefried W, Aziz NM, Rowland JH, Pinto BM. (2005) Riding the crest of the teachable moment: promoting long-term health after the diagnosis of cancer. *J Clin Oncol* 23:5814–5830.
9. Capocaccia R, Colonna M, Corazziari I, et al. (2002) Measuring cancer prevalence in Europe: the EUROPREVAL project. *Ann Oncol* 13:831–839.
10. Brown BW, Brauner C, Minnotte MC. (1993) Noncancer deaths in white adult cancer patients. *J Natl Cancer Inst* 85:979–987.
11. McBride CM, Emmons KM, Lipkus IM. (2003) Understanding the potential of teachable moments: the case of smoking cessation. *Health Educ Res* 18:156–170.
12. McBride CM, Ostroff JS. (2003) Teachable moments for promoting smoking cessation: the context of cancer care and survivorship. *Cancer Control* 10:325–333.
13. Dishman RK, Washburn RA, Heath GW. (2004) Physical activity epidemiology. *Human Kinetics*, Champaign, IL, pp 4.
14. Pate RR, Pratt M, Blair SN, et al. (1995) Physical activity and public health. A recommendation from the centers for disease control and prevention and the American college of sports medicine. *J Am Med Assoc* 273:402–407.
15. Haskell WL, Lee IM, Pate RR, et al. (2007) Physical activity and public health: updated recommendation for adults from the American college of sports medicine and the American heart association. *Med Sci Sports Exerc* 39:1423–1434.
16. Department of Health. (2004) Evidence on the impact of physical activity and its relationship to health. Chief Medical Officer's Report. HSMO, London.
17. World Cancer Research Fund/American Institute for Cancer Research. (2007) Food, Nutrition, Physical Activity, and the prevention of cancer: a global perspective. AICR, Washington, DC.
18. Knols R, Aaronson NK, Uebelhart D, Franssen J, Aufdemkampe G. (2005) Physical exercise in cancer patients during and after medical treatment: a systematic review of randomized and controlled clinical trials. *J Clin Oncol* 23:3830–3842.
19. McNeely ML, Campbell KL, Rowe BH, Klassen TP, Mackey JR, Courneya KS. (2006) Effects of exercise on breast cancer patients and survivors: a systematic review and meta-analysis. *CMAJ* 175:34–41.
20. Holmes MD, Chen WY, Feskanich D, Kroenke CH, Colditz GA. (2005) Physical activity and survival after breast cancer diagnosis. *J Am Med Assoc* 293:2479–2486.
21. Holick CN, Newcomb PA, Trentham-Dietz A, et al. (2008) Physical activity and survival after diagnosis of invasive breast cancer. *Cancer Epidemiol Biomark Prev* 17:379–386.
22. Pierce JP, Stefanick ML, Flatt SW, et al. (2007) Greater survival after breast cancer in physically active women with high vegetable-fruit intake regardless of obesity. *J Clin Oncol* 25:2345–2351.
23. Irwin ML, Smith AW, McTiernan A, et al. (2008) Influence of pre- and post-diagnosis physical activity on mortality in breast cancer survivors: the health, eating, activity, and lifestyle study. *J Clin Oncol* 26:3958–3964.
24. Meyerhardt JA, Giovannucci EL, Holmes MD, et al. (2006) Physical activity and survival after colorectal cancer diagnosis. *J Clin Oncol* 24:3527–3534.
25. Meyerhardt JA, Heseltine D, Niedzwiecki D, et al. (2006) Impact of physical activity on cancer recurrence and survival in patients with stage III colon cancer: findings from CALGB 89803. *J Clin Oncol* 24:3535–3541.
26. Haydon AM, Macinnis RJ, English DR, Giles GG. (2006) Effect of physical activity and body size on survival after diagnosis with colorectal cancer. *Gut* 55:62–67.
27. Kroenke CH, Chen WY, Rosner B, Holmes MD. (2005) Weight, weight gain, and survival after breast cancer

- diagnosis. *J Clin Oncol* 23:1370–1378.
28. Bastarrachea J, Hortobagyi GN, Smith TL, Kau SW, Buzdar AU. (1994) Obesity as an adverse prognostic factor for patients receiving adjuvant chemotherapy for breast cancer. *Ann Intern Med* 120:18–25.
 29. Bassett WW, Cooperberg MR, Sadetsky N, et al. (2005) Impact of obesity on prostate cancer recurrence after radical prostatectomy: data from CaPSURE. *Urology* 66:1060–1065.
 30. Caspersen CJ, Powell KE, Christenson GM. (1985) Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Public Health Rep* 100: 126–131.
 31. Winter EM, Fowler N. (2009) Exercise defined and quantified according to the Systeme International d'Unites. *J Sports Sci* 27:447–460.
 32. Physical Activity, Fitness, and Health Consensus Statement. Human Kinetics Publishers 1993, Kingswood, Australia, p 15.
 33. Kushi LH, Byers T, Doyle C, et al. (2006) American cancer society guidelines on nutrition and physical activity for cancer prevention: reducing the risk of cancer with healthy food choices and physical activity. *CA Cancer J Clin* 56:254–281.
 34. Cancer Research UK. (2008) <http://info.cancerresearchuk.org/cancerstats/>, <http://info.cancerresearchuk.org/cancerstats/>. Accessed 30 January 2008.
 35. Doyle C, Kushi LH, Byers T, et al. (2006) Nutrition and physical activity during and after cancer treatment: an American cancer society guide for informed choices. *CA Cancer J Clin* 56:323–353.
 36. Sheikh A, Smeeth L, Ashcroft R. (2002) Randomised controlled trials in primary care: scope and application. *Br J Gen Pract* 52:746–751.
 37. Cella DF, Tulsky DS, Gray G, et al. (1993) The functional assessment of cancer therapy scale: development and validation of the general measure. *J Clin Oncol* 11:570–579.
 38. Craig CL, Marshall AL, Sjostrom M, et al. (2003) International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* 35:1381–1395.
 39. Godin G, Shephard RJ. (1985) A simple method to assess exercise behavior in the community. *Can J Appl Sport Sci* 10:141–146.
 40. Godin G, Jobin J, Bouillon J. (1986) Assessment of leisure time exercise behavior by self-report: a concurrent validity study. *Can J Public Health* 77:359–362.
 41. Sallis JF, Saelens BE. (2000) Assessment of physical activity by self-report: status, limitations, and future directions. *Res Q Exerc Sport* 71:S1–S14.
 42. Ainsworth BE, Haskell WL, Whitt MC, et al. (2000) Compendium of physical activities: an update of activity codes and MET intensities. *Med Sci Sports Exerc* 32:S498–S504.
-
1. Radbruch L, Strasser F, Elsner F, Goncalves JF, Loge J, Kaasa S, et al. (2008) Fatigue in palliative care patients – an EAPC approach. *Palliative Med* 22:13–32.
 2. Stone PC, Minton O. (2008) Cancer-related fatigue. *Eur J Cancer* 44:1097–1104.
 3. NCCN. (2008) National Comprehensive Cancer Network Clinical Practice Guidelines in Oncology: cancer related fatigue. Available from: <http://www.nccn.org> . Accessed 17 August 2008.
 4. Hofman M, Ryan JL, Figueroa-Moseley CD, Jean-Pierre P, Morrow GR. (2007) Cancerrelated fatigue: the scale of the problem. *Oncologist* 12:4–10.
 5. Stevinson C, Lawlor DA, Fox KA. (2004) Exercise interventions for cancer patients: systematic review of controlled trials. *Cancer Cause Control* 15:1035–1056.
 6. Knols R, Aaronson NK, Duebelhart D, Franssen J, Aufdemkampe G. (2005) Physical exercise in cancer patients during and after medical treatment: a systematic review of randomized and controlled trials. *J Clin Oncol* 23:3830–3841.
 7. Schmitz KH, Holtzman J, Courneya KS, Masse LC, Duval S, Kane R. (2005) Controlled physical activity trials in cancer survivors: a systematic review and meta-analysis. *Cancer Epidem Biomar Prev* 14:1588–1595.
 8. Hjermstad MJ, Fayers PM, Bjordal K, Kaasa S. (1998) Using reference data on quality of life – the importance of adjusting for age and gender, exemplified by the EORTC QLQ-C30 (+3). *Eur J Cancer* 34:1381–1389.
 9. Prue G, Rankin J, Allen J, Gracey J, Cramp F. (2006) Cancer-related fatigue: a critical appraisal. *Eur J Cancer*

42:846–863.

10. Meeske K, Smith AW, Alfano CM, McGregor BA, McTiernan A, Baumgartner KB, et al. (2007) Fatigue in breast cancer survivors two to five years post diagnosis: a HEAL study report. *Qual Life Res* 16:947–960.
11. Servaes P, Gielissen MF, Verhagen S, Bleijenberg G. (2007) The course of severe fatigue in disease-free breast cancer patients: a longitudinal study. *Psycho Oncol* 16:787–795.
12. Hjermland MJ, Fossa SD, Oldervoll L, Holte H, Jacobsen AB, Loge JH. (2005) Fatigue in long-term Hodgkin's disease survivors: a follow-up study. *J Clin Oncol* 23:6587–6595.
13. Mock V, St Ours C, Hall S, Bositis A, Tillery M, Belcher A, et al. (2007) Using a conceptual model in nursing research – mitigating fatigue in cancer patients. *J Adv Nurs* 58: 503–512.
14. Ruckdeschel JC. (2005) Fatigue is becoming an exhausting problem. *Cancer* 103:213–215.
15. Bruera E. (2007) Managing cancer-related symptoms: advances in supportive oncology. *Clin Adv Hematol Oncol* 5:29–31.
16. Jacobsen PB, Donovan KA, Vadaparampil ST, Small BJ. (2007) Systematic review and metaanalysis of psychological and activity-based interventions for cancer-related fatigue. *Health Psychol* 26:660–667.
- 34 M.L. McNeely and K.S. Courneya 17. Ahlberg K, Ekman T, Gaston-Johansson F, Mock V. (2003) Assessment and management of cancer-related fatigue in adults. *Lancet* 362:640–650.
18. Boohene JA, Bruera E. (2006) Treatment of fatigue in palliative care. In: Bruera E, et al. (eds) *Textbook of Palliative Medicine*. Hodder Arnold, New York, pp 639–672.
19. Evans WJ, Lambert CP. (2007) Physiological basis of fatigue. *Am J Phys Med Rehabil* 86:S29–S46.
20. Mock V, Atkinson A, Barsevick AM, Berger AM, Cimprich B, Eisenberger MA, et al. (2007) Cancer-related fatigue: clinical practice guidelines in oncology. *J Natl Compr Cancer Netw* 5:1054–1078.
21. Loge J. (2006) Assessment of fatigue in palliative care. In: Bruera E, et al. (eds) *Textbook of Palliative Med*. Hodder Arnold, New York, pp. 613–620.
22. Cella D, Peterman A, Passik S, Jacobsen P, Breitbart W. (1998) Progress toward guidelines for the management of fatigue. *Oncology (Williston Park)* 12:369–377.
23. Minton O, Stone P. (2009) A systematic review of scales used for the measurement of cancer-related fatigue. *Ann Oncol* 20:27–33.
24. Stone P. (2002) The measurement, causes and effective management of cancer-related fatigue. *Int J Palliative Nurs* 8:120–128.
25. Cella D, Davis K, Breitbart W, Curt G. (2001) Cancer-related fatigue: prevalence of proposed diagnostic criteria in a united states sample of cancer survivors. *J Clin Oncol* 19:3385–3391.
26. Minton O, Richardson A, Sharpe M, Hotopf M, Stone P. (2008) A systematic review and meta-analysis of the pharmacological treatment of cancer-related fatigue. *J Natl Cancer Inst* 100:1155–1166.
27. Bohlius J, Weingart O, Trelle S, Engert A. (2006) Cancer-related anemia and recombinant human erythropoietin – an updated Overview. *Nat Clin Pract Oncol* 3:152–164.
28. Bennett CL, Silver SM, Djulbegovic B, Samaras AT, Blau CA, Gleason KJ, et al. (2008) Venous thromboembolism and mortality associated with recombinant erythropoietin and darbepoetin administration for the treatment of cancer-associated anemia. *J Am Med Assoc* 299:914–924.
29. Jim HSL, Jacobsen PB. (2008) Assessment and management of cancer-related fatigue. In: Lyman GH, Crawford J (eds) *Cancer Supportive Care*. Informa Healthcare, New York, pp 13–29.
30. Luebbert K, Dahme B, Hasenbring M. (2001) The effectiveness of relaxation training in reducing treatment-related symptoms and improving emotional adjustment in acute nonsurgical cancer treatment: a meta-analytical review. *Psycho Oncol* 10:490–502.
31. McNeely ML, Campbell KL, Rowe BH, Klassen TP, Mackey JR, Courneya KS, et al. (2006) Effects of exercise on breast cancer patients and survivors: a systematic review and metaanalysis. *CMAJ* 175:34–41.
32. Oldervoll LM, Kaasa S, Hjermland MJ, Lund JA, Loge JH. (2004) Physical exercise results in the improved subjective well-being of a few or is effective rehabilitation for all cancer patients?. *Eur J Cancer* 40:951–962.
33. Cramp F, Daniel J. (2008). Exercise for the management of cancer-related fatigue in adults. *Cochrane Database Syst Rev* (2):CD006145.
34. Markes M, Brockow T, Resch KL. (2006). Exercise for women receiving adjuvant therapy for breast cancer. *Cochrane Database Syst Rev* (4):CD005001.
35. McNeely ML, Peddle CJ, Parliament M, Courneya KS. (2006) Cancer rehabilitation: recommendations for

- integrating exercise programming in the clinical practice setting. *Curr Cancer Therapy Rev* 2:351–360.
36. Thorsen L, Courneya KS, Stevinson C, Fossa SD. (2008) A systematic review of physical activity in prostate cancer survivors: outcomes, prevalence, and determinants. *Supp Care Cancer* 16:987–997.
 37. Segal RJ, Reid RD, Courneya KS, Malone SC, Parliament MB, Scott CG, et al. (2003) Resistance exercise in men receiving androgen deprivation therapy for prostate cancer. *J Clin Oncol* 21:1653–1659.
 38. Segal RJ, Reid RD, Courneya KS, Sigal RJ, Kenny GP, Prud'Homme DG, et al. (2009) Randomized controlled trial of resistance or aerobic exercise in men receiving radiation therapy for prostate cancer. *J Clin Oncol* 27:344–351.
 39. Wiskemann J, Huber G. (2008) Physical exercise as adjuvant therapy for patients undergoing hematopoietic stem cell transplantation. *Bone Marrow Transpl* 41:321–329.
 40. Wilson RW, Jacobsen PB, Fields KK. (2005) Pilot study of a home-based aerobic exercise program for sedentary cancer survivors treated with hematopoietic stem cell transplantation. *Bone Marrow Transpl* 35:721–727.
 41. Carlson LE, Smith D, Russell J, Fibich C, Whittaker T. (2006) Individualized exercise program for the treatment of severe fatigue in patients after allogeneic hematopoietic stem-cell transplant: a pilot study. *Bone Marrow Transpl* 37:945–954.
 42. Dimeo FC, Stieglitz RD, Novelli-Fischer U, Fetscher S, Keul J. (1999) Effects of physical activity on the fatigue and psychologic status of cancer patients during chemotherapy. *Cancer* 85:2273–2277.
 43. Coleman EA, Coon S, Hall-Barrow J, Richards K, Gaylor D, Stewart B. (2003) Feasibility of exercise during treatment for multiple myeloma. *Cancer Nurs* 26:410–419.
 44. Lowe SS, Watanabe SM, Courneya KS. (2009) Physical activity as a supportive care intervention in palliative cancer patients: a systematic review. *J Supp Oncol* 7: 27–34.
 45. Oldervoll LM, Loge JH, Paltiel H, Asp MB, Vidvei U, Wiken AN, et al. (2006) The effect of a physical exercise program in palliative care: a phase II study. *J Pain Symptom Manag* 31:421–430.
 46. Headley JA, Ownby KK, John LD. (2004) The effect of seated exercise on fatigue and quality of life in women with advanced breast cancer. *Oncol Nurs Forum* 31:977–983.
 47. Courneya KS, Friedenreich CM, Sela RA, Quinney HA, Rhodes RE, Handman M. (2003) The group psychotherapy and home-based physical exercise (group-hope) trial in cancer survivors: physical fitness and quality of life outcomes. *Psycho Oncol* 12:357–374.
 48. Heim ME, v d Malsburg ML, Niklas A. (2007) Randomized controlled trial of a structured training program in breast cancer patients with tumor-related chronic fatigue. *Oncologie* 30:429–434.
 49. Courneya KS, Jones LW, Peddle CJ, Sellar CM, Reiman T, Joy AA, et al. (2008) Effects of aerobic exercise training in anemic cancer patients receiving darbepoetin alfa: a randomized controlled trial. *Oncologist* 13:1012–1020.
 50. Courneya KS, Mackey JR, Rhodes RE. (2004) Cancer. In: Lemura LM, von Duvillard, SP (eds) *Clinical Exercise Physiology*. Lippincott Williams & Wilkins, Philadelphia, pp 387–404.
 51. Lane K, McKenzie DC. (2005) Cancer. In: Skinner JS (ed) *Exercise Testing and Exercise Prescription for Special Cases*. Lippincott Williams & Wilkins, Baltimore, pp 363–376.
 52. ACSM (2006) ACSM's guidelines for exercise testing and prescription, 7th ed. ed. Whaley MH, Brubaker PH (eds), Lippincott Williams & Wilkins, Baltimore.
 53. Finch E, Brooks D, Stratford PW, Mayo NE. (2002) *Physical Rehabilitation Outcome Measures*, 2nd ed. BC Decker, Hamilton, pp. 1–292.
 54. Rikli RE, Jones CJ. (2001) *Senior Fitness Test Manual*. Human Kinetics, Champaign, p 161.
 55. Cress ME, Petrella JK, Moore TL, Schenkman ML. (2005) Continuous-scale physical functional performance test: validity, reliability, and sensitivity of data for the short version. *Phys Ther* 85:323–335.
 56. McKelvie RS. (2005) Heart failure. In: Skinner JS (ed) *Exercise Testing and Exercise Prescription For Special Cases*. Lippincott Williams & Wilkins, Baltimore, pp 323–336.
 57. MacVicar MG, Winningham ML. (1986) Promoting the functional capacity of cancer patients. *Cancer Bull* 38:235–239.
 58. Mock V, Burke MB, Sheehan P, Creaton EM, Winningham ML, McKenney-Tedder S, et al. (1994) A nursing rehabilitation program for women with breast cancer receiving adjuvant chemotherapy. *Oncol Nurs Forum* 21:899–907.
- 36 M.L. McNeely and K.S. Courneya
59. McArdle WD, Katch FI, Katch VL. (2001) *Exercise Physiology*, 5th ed. Lippincott Williams & Wilkins, Baltimore, p 1158.

60. Bennett K. (2005) Therapeutic exercise for arthritis. In: Hall CM, Brody LT (eds) *Therapeutic Exercise: Moving Towards Function*. Lippincott Williams & Wilkins, Baltimore, pp 229–241.
1. Ferlay J, Autier P, Boniol M, et al. (2007) Estimates of the cancer incidence and mortality in Europe in 2006. *Ann Oncol* 18(3):581–592.
 2. American Cancer Society (2009) Statistics for 2008. <http://www.cancer.org/downloads/STT/2008CAFFfinalsecured.pdf>. Accessed 11 March 2009
 3. Rock CL, Demark Wahnefried W. (2002) Nutrition and survival after the diagnosis of breast cancer: a review of the evidence. *J Clin Oncol* 20(15):3302–3316.
 4. National Comprehensive Cancer Network (2009) Cancer-Related Fatigue. http://www.nccn.org/professionals/physician_gls/PDF/fatigue.pdf. Accessed 18 March 2009
 5. Hoff AO, Gagel RF. (2005) Osteoporosis in breast and prostate cancer survivors. *Oncology* 19(5):651–658.
 6. Browall M, Ahlberg K, Karlsson P, et al. (2008) Health-related quality of life during adjuvant treatment for breast cancer among postmenopausal women. *Eur J Oncol Nurs* 12(3):180–189.
 7. Burgess C, Cornelius V, Love S, et al. (2005) Depression and anxiety in women with early breast cancer: five year observational cohort study. *BMJ* 330(7493):702. Epub 2005 February 4.
 8. Groenvold M, Fayers PM, Petersen MA, et al. (2007) Breast cancer patients on adjuvant chemotherapy report a wide range of problems not identified by health-care staff. *Breast Cancer Res Treat* 103(2):185–195.
 9. Burstein HJ. (2007) Cognitive side-effects of adjuvant treatments. *Breast* 16(Suppl 2): S166–S168.
 10. Phillips KA, Bernhard J. (2003) Adjuvant breast cancer treatment and cognitive function: current knowledge and research directions. *J Natl Cancer Inst* 95(3):190–197.
 11. Rugo HS, Ahles T. (2003) The impact of adjuvant therapy for breast cancer on cognitive function: current evidence and directions for research. *Semin Oncol* 30(6):749–762.
 12. Cleeland CS. (2007) Symptom burden: multiple symptoms and their impact as patient-reported outcomes. *J Natl Cancer Inst Monogr* 37:16–21.
 13. Gapstur RL. (2007) Symptom burden: a concept analysis and implications for oncology nurses. *Oncol Nurs Forum* 34(3):673–680. 50
 14. Andrykowski MA, Beacham AO, Jacobsen PB. (2007) Prospective, longitudinal study of leisure-time exercise in women with early-stage breast cancer. *Cancer Epidemiol Biomarkers Prev* 16(3):430–438.
 15. Irwin ML, Crumley D, McTiernan A, et al. (2003) Physical activity levels before and after a diagnosis of breast carcinoma: the health, eating, activity, and lifestyle (HEAL) study. *Cancer* 97(7):1746–1757.
 16. Pinto BM, Trunzo JJ, Reiss P, et al. (2002) Exercise participation after diagnosis of breast cancer: trends and effects on mood and quality of life. *Psychooncology* 11(5):389–400.
 17. Sjöström M, Oja P, Hagströmer M, et al. (2006) Health-enhancing physical activity across European union countries: the eurobarometer study. *J Public Health* 14(5):291–300.
 18. Kortebein P, Ferrando A, Lombeida J, et al. (2007) Effect of 10 days of bed rest on skeletal muscle in healthy older adults. *Jama* 297(16):1772–1774.
 19. Demiot C, Dignat-George F, Fortrat JO, et al. (2007) WISE 2005: chronic bed rest impairs microcirculatory endothelium in women. *Am J Physiol Heart Circ Physiol* 293(5):159–164.
 20. Haskell WL, Lee IM, Pate RR, et al. (2007) Physical activity and public health: updated recommendation for adults from the American college of sports medicine and the American heart association. *Med Sci Sports Exerc* 39(8):1423–1434.
 21. Markes M, Brockow T, Resch KL. (2006) Exercise for women receiving adjuvant therapy for breast cancer. *Cochrane Database Syst Rev* 18(4):CD005001.
 22. Campbell A, Mutrie N, White F, et al. (2005) A pilot study of a supervised group exercise programme as a rehabilitation treatment for women with breast cancer receiving adjuvant treatment. *Eur J Oncol Nurs* 9(1):56–63.
 23. Mutrie N, Campbell AM, Whyte F, et al. (2007) Benefits of supervised group exercise programme for women being treated for early stage breast cancer: pragmatic randomized controlled trial. *BMJ* 334(7592):517. Epub 2007 February 16.
 24. Courneya KS, Segal RJ, Mackey JR, et al. (2007) Effects of aerobic and resistance exercise in breast cancer patients

- receiving adjuvant chemotherapy: a multicenter randomized controlled trial. *J Clin Oncol* 25(28):4396–4404.
25. Mock V, Frangakis C, Davidson NE, et al. (2005) Exercise manages fatigue during breast cancer treatment: A randomized controlled trial. *Psychooncology* 14(6):464–477.
 26. Segal R, Evans W, Johnson D, et al. (2001) Structured exercise improves physical functioning in women with stages I and II breast cancer: Results of a randomized controlled trial. *J Clin Oncol* 19(3):657–665.
 27. Schwartz AL, Winters-Stone K, Gallucci B. (2007) Exercise effects on bone mineral density in women with breast cancer receiving adjuvant chemotherapy. *Oncol Nurs Forum* 34(3): 627–633.
 28. Payne JK, Held J, Thorpe J, et al. (2008) Effect of exercise on biomarkers, fatigue, sleep disturbances, and depressive symptoms in older women with breast cancer receiving hormonal therapy. *Oncol Nurs Forum* 35(4):635–642.
 29. Winningham ML, MacVicar MG. (1988) The effect of aerobic exercise on patients reports of nausea. *Oncol Nurs Forum* 15(4):447–450.
 30. Crowley SA (2003) The effect of a structured exercise program on fatigue, strength, endurance, physical self-efficacy, and functional wellness in women with early stage breast cancer. *Digital Dissertations, University of Michigan*. p. 127
 31. Drouin J (2002) Aerobic exercise training effects on physical function, fatigue and mood, immune status, and oxidative stress in subjects undergoing radiation treatment for breast cancer. *Wayne State University*. p. 142
 32. Battaglini C, Bottaro M, Dennehy C, et al. (2006) The effects of resistance training on muscular strength and fatigue levels in breast cancer patients. *Rev Bras Med Esporte* 12(3):139e–144e.
 33. Mock V, Dow KH, Meares CJ, et al. (1997) Effects of exercise on fatigue, physical functioning, and emotional distress during radiation therapy for breast cancer. *Oncol Nurs Forum* 24(6):991–1000.
 34. MacVicar MG, Winningham ML. (1986) Promoting the functional capacity of cancer patients. *Cancer Bull* 28:235–239.
 35. US Department of Health and Human Services (1996) Physical Activity and Health: A Report of the Surgeon General. <http://www.cdc.gov/nccdphp/sgr/pdf/sgrfull.pdf>. Assessed 18 March 2009
 36. Barnett GC, Shah M, Redman K, et al. (2008) Risk factors for the incidence of breast cancer: do they affect survival from the disease? *J Clin Oncol* 26(20):3310–3316.
 37. Adler RA. (2007) Cancer treatment-induced bone loss. *Curr Opin Endocrinol Diabetes Obes* 14(6):442–445.
 38. Brown SA, Guise TA. (2009) Cancer treatment-related bone disease. *Crit Rev Eukaryot Gene Expr* 19(1):47–60.
 39. Ioannidis JP, Evans SJ, Gotzsche PC, et al. (2004) Better reporting of harms in randomized trials: an extension of the CONSORT statement. *Ann Intern Med* 141(10):781–788.
 40. Schmitz KH, Ahmed RL, Hannan PJ, et al. (2005) Safety and efficacy of weight training in recent breast cancer survivors to alter body composition, insulin, and insulin-like growth factor axis proteins. *Cancer Epidemiol Biomarkers Prev* 14(7):1672–1680.
 41. Kim CJ, Kang DH, Smith BA, et al. (2006) Cardiopulmonary responses and adherence to exercise in women newly diagnosed with breast cancer undergoing adjuvant therapy. *Cancer Nurs* 29:156–165.
 42. Mac Vicar MG, Winningham ML, Nickel JL. (1989) Effects of aerobic interval training on cancer patients' functional capacity. *Nurs Res* 38:348–351.
 43. Courneya KS, Segal RJ, Gelmon K, et al. (2007) Six-month follow-up of patient-rated outcomes in a randomized controlled trial of exercise training during breast cancer chemotherapy. *Cancer Epidemiol Biomarkers Prev* 16(12):2572–2578.
 44. Courneya KS, McKenzie DC, Reid RD, et al. (2008) Barriers to supervised exercise training in a randomized controlled trial of breast cancer patients receiving chemotherapy. *Ann Behav Med* 35(1):116–122.
 45. Emslie C, Whyte F, Campbell A, et al. (2007) 'I wouldn't have been interested in just sitting round a table talking about cancer'; exploring the experiences of women with breast cancer in a group exercise trial. *Health Educ Res* 22(6):827–838.
 46. Ries LAG, Melbert D, Krapcho M, et al. (2008) SEER Cancer Statistics Review, 1975–2005. http://seer.cancer.gov/csr/1975_2005/. Assessed 12 March 2009.
 47. Große-Wächter E, Ledvina I, Schüle K, et al. (2004) Bewegung und Sport in der Krebsnachsorge. http://www.wirimsport.de/templates/dokukategorien/dokumanagement/psdoc/file/77/2004_04_02434d15ad599bf.pdf. Assessed 18 March 2009.

1. Office of National Statistics. (2005) Cancer Statistics Registrations: Registrations of Cancer Diagnosed in 2003. Series MB1 No. 33, England.
2. Parkin DM, Bray F, Ferlay J, et al. (2005) Global cancer statistics 2002. *CA Cancer J Clin* 55:74–108.
3. Office of National Statistics. (2002) Cancer survival in the health authorities of England 1993– 2000. *Health Stats Quart* 13:95–103.
4. Hewitt M, Greenfield S, Stovall E. (2006) *From Cancer Patient to Cancer Survivor: Lost in Translation*. US National Academy of Sciences, Washington, DC.
5. Courneya KS, Friedenreich CM. (1999) Physical exercise and quality of life following cancer diagnosis: a literature review. *Ann Behav Med* 21:171–179.
6. Glanz K, Lerman C. (1992) Psychosocial impact of breast cancer: a critical review. *Ann Behav Med* 14:204–212.
7. Byar KL, Berger AM, Bakken SL, et al. (2006) Impact of adjuvant breast cancer chemotherapy on fatigue, other symptoms, and quality of life. *Oncol Nurs Forum* 33:E18–26.
8. Harmer V. (2003) *Breast Cancer: Nursing Care and Management*. Whurr, London.
9. Morrow GR, Shelke AR, Roscoe JA, et al. (2005) Management of cancer-related fatigue. *Cancer Invest* 23:229–239.
10. Fialka-Moser V, Crevenna R, Korpan M, et al. (2003) Cancer rehabilitation: particularly with aspects on physical impairments. *J Rehab Med* 35:153–162.
11. Fallowfield L, Clark A. (1992) *Breast Cancer*. Routledge, London.
12. Golant M, Altman T, Martin C. (2003) Managing cancer side effects to improve quality of life: a cancer psychoeducation program. *Cancer Nurs* 26:37–44.
13. Burgess C, Ramirez A, Cornelius V, et al. (2005) Depression and anxiety in women with early breast cancer: 5-year observational cohort data. *BMJ* 330:702–705.
14. Greeno CG, Wing RR. (1994) Stress-induced eating. *Psychol Bull* 115:444–464.
15. Wardle J, Steptoe A, Oliver G, et al. (2000) Stress, dietary restraint and food intake. *J Psychosomatic Res* 48:195–202.
16. Meyer T, Mark M. (1995) Effects of psychosocial interventions with adult cancer patients: a meta-analysis of randomised experiments. *Health Psychol* 14:101–108.
17. Courneya KS, Friedenreich CM. (1997a) The relationship between exercise pattern across the cancer experience and current quality of life in colorectal cancer survivors. *J Altern Complem Med* 3:215–226.
18. Department of Health. (2004) At least five a week: evidence on the impact of physical activity and its relationship to health. A Report from the Chief Medical Officer. (http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_4080994). Accessed 11 March 2007.
19. Taylor RS, Brown A, Ebrahim S, et al. (2004) Exercise-based rehabilitation for patients with coronary heart disease: systematic review and meta-analysis of randomized controlled trials. *Am J Med* 116:682–692.
20. Pedersen BK, Saltin B. (2006) Evidence for prescribing exercise as therapy in chronic disease. *Scand J Med Sci Sports* 1:3–63.
21. Lawlor DA, Hopker SW. (2001) The effectiveness of exercise as an intervention in the management of depression: systematic review and meta-regression analysis of randomized controlled trials. *BMJ* 322:1–8.
22. Etnier J, Salazar W, Landers D, et al. (1997) The influence of physical fitness and exercise upon cognitive functioning: a meta-analysis. *J Sport Exerc Psychol* 19: 249–277.
23. King AC, Oman RF, Brassington GS, et al. (1997) Moderate-intensity exercise and self-rated quality of sleep in older adults: a randomised controlled trial. *J Am Med Assoc* 277:32–37.
24. Blanchard CM, Courneya KS, Stein K. (2008) Cancer survivors' adherence to lifestyle behavior recommendations and associations with health-related quality of life: Results from the American Cancer Society's SCS-II. *J Clin Oncol* 26:2198–2220.
25. Irwin ML, McTiernan A, Bernstein L, et al. (2004) Physical activity levels among breast cancer survivors. *Med Sci Sports Exerc* 36:1484–1491.
26. Irwin M, Crumley D, McTiernan A, et al. (2003) Physical activity levels before and after a diagnosis of breast carcinoma. *Cancer* 97:1746–1757.
27. Blanchard CM, Cokkinides V, Courneya KS, et al. (2003) Comparison of physical activity of post-treatment breast cancer survivors and non-cancer controls. *Behav Med* 28:140–149.
28. Ness KK, Wall MM, Oakes JM, et al. (2006) Physical performance limitations and participation restrictions among cancer survivors: a population-based study. *Ann Epidemiol* 16:197–205.

29. MacVicar M, Winningham ML. (1986) Promoting the functional capacity of cancer patients. *Cancer Bull* 38:235–239.
30. Winningham ML, MacVicar MG. (1988) The effect of aerobic exercise on patient reports of nausea. *Oncol Nurs Forum* 15:447–450.
31. MacVicar M, Winningham ML, Nickel JL. (1989) Effects of aerobic interval training on cancer patients' functional capacity. *Nurs Res* 38:348–351.
32. Winningham ML, MacVicar MG, Bondoc M, et al. (1989) Effect of aerobic exercise on body weight and composition in patients with breast cancer on adjuvant chemotherapy. *Oncol Nurs Forum* 16:683–689.
33. Pinto BM, Clark MM, Maruyama NC, et al. (2003) Psychological and fitness changes associated with exercise participation among women with breast cancer. *Psychooncology* 12:118–126.
34. Courneya KS, Mackey JR, Bell GJ, et al. (2003) Randomized controlled trial of exercise training in postmenopausal breast cancer survivors: cardiopulmonary and quality of life outcomes. *J Clin Oncol* 21:1660–1668.
35. Knols R, Aaronson NK, Uebelhart D, et al. (2005) Physical exercise in cancer patients during and after medical treatment: a systematic review of randomized and controlled clinical trials. *J Clin Oncol* 23:3830–3842.
36. Daley AJ, Mutrie N, Crank H, et al. (2004) Exercise therapy in women who have had breast cancer: design of the Sheffield women's exercise and well-being project. *Health Educ Res* 19:686–697.
37. Daley AJ, Crank H, Saxton JM, et al. (2007) A Randomized trial of exercise therapy in women treated for breast cancer. *J Clin Oncol* 25:1713–1721.
38. Irwin ML, Cadmus L, Alvarez-Reeves M, et al. (2008) Recruiting and retaining breast cancer survivors into a randomized controlled exercise trial: the Yale Exercise and Survivorship Study. *Cancer* 112(11 Suppl):2593–2606.
39. Asikainen TM, Kukkonen-Harjula K, Miilunpalo S. (2004) Exercise for health for early postmenopausal women: a systematic review of randomised controlled trials. *Sports Med* 34:753–778.
40. Zehnacker CH, Bemis-Dougherty A. (2007) Effect of weighted exercises on bone mineral density in postmenopausal women: a systematic review. *J Geriatr Phys Ther* 30:79–88.
41. Chen Z, Maricic M, Aragaki AK, Mouton C, et al. (2009) Fracture risk increases after diagnosis of breast or other cancers in postmenopausal women: Results from the Women's Health Initiative. *Osteoporosis Int* 20:527–536.
42. Reid DM, Doughty J, Eastell R, et al. (2008) Guidance for the management of breast cancer treatment-induced bone loss: a consensus position statement from a UK Expert Group. *Cancer Treat Rev* 34(Suppl 1):S3–S18.
43. Ohira T, Schmitz KH, Ahmed RL, et al. (2006) Effects of weight training on quality of life in recent breast cancer survivors: the Weight Training for Breast Cancer Survivors (WTBS) study. *Cancer* 106:2076–2083.
44. Ozaslan C, Kuru B. (2004) Lymphedema after treatment for breast cancer. *Am J Surg* 187: 69–72.
45. Mirolo BR, Bunce IH, Chapman M, et al. (1995) Psychosocial benefits of post-mastectomy lymphedema therapy. *Cancer Nurs* 18:197–205.
46. Ahmed RL, Thomas W, Yee D, et al. (2006) Randomized controlled trial of weight training and lymphedema in breast cancer survivors. *J Clin Oncol* 24:2765–2772.
47. Schmitz KH, Ahmed RL, Hannan PJ, et al. (2005) Safety and efficacy of weight training in recent breast cancer survivors to alter body composition, insulin, and insulin-like growth factor axis proteins. *Cancer Epidemiol Biomark Prev* 14:1672–1680.
48. Nieman DC, Cook VD, Henson DA, et al. (1995) Moderate exercise training and natural killer cell cytotoxic activity in breast cancer patients. *Int J Sports Med* 16:334–337.
49. McKenzie DC, Kalda AL. (2003) Effect of upper extremity exercise on secondary lymphedema in breast cancer patients: a pilot study. *J Clin Oncol* 21:463–466.
50. Herrero F, San Juan AF, Fleck SJ, et al. (2006) Combined aerobic and resistance training in breast cancer survivors: a randomized, controlled pilot trial. *Int J Sports Med* 27: 573–580.
51. Milne HM, Wallman KE, Gordon S, et al. (2008) Effects of a combined aerobic and resistance exercise program in breast cancer survivors: a randomized controlled trial. *Breast Cancer Res Treat* 108:279–288.
52. Cella D, Eton DT, Lai JS, et al. (2002) Combining anchor and distribution-based methods to derive minimal clinically important differences on the Functional Assessment of Cancer Therapy (FACT) anemia and fatigue scales. *J Pain Symptom Manag* 24:547–561.
53. Milne HM, Wallman KE, Gordon S, et al. (2008) Impact of a combination resistance and aerobic exercise program on motivational variables in breast cancer survivors: a randomised controlled trial. *Ann Behav Med* 36:158–166.
54. Mutrie N, Campbell AM, Whyte F, et al. (2007) Benefits of supervised group exercise programme for

- women being treated for early stage breast cancer: pragmatic randomised controlled trial. *BMJ* 334:517.
55. Pinto BM, Frierson GM, Rabin C, et al. (2005) Home-based physical activity intervention for breast cancer patients. *J Clin Oncol* 23:3577–3587.
 56. Demark-Wahnefried W, Clipp EC, Morey MC, et al. (2006) Lifestyle intervention development study to improve physical function in older adults with cancer: outcomes from project LEAD. *J Clin Oncol* 24:3465–3473.
 57. Vallance JK, Courneya KS, Plotnikoff RC, et al. (2007) Randomized controlled trial of the effects of print materials and step pedometers on physical activity and quality of life in breast cancer survivors. *J Clin Oncol* 25:2352–2359.
 58. Vallance JK, Courneya KS, Plotnikoff RC, et al. (2008) Maintenance of physical activity in breast cancer survivors after a randomized trial. *Med Sci Sports Exerc* 40: 173–180.
 59. Jones LW, Courneya KS, Fairey AS, et al. (2004) Effects of an oncologist's recommendation to exercise on self-reported exercise behavior in newly diagnosed breast cancer survivors: a single-blind, randomized controlled trial. *Ann Behav Med* 28:105–113.
 60. Holmes MD, Chen WY, Feskanich D, et al. (2005) Physical activity and survival after breast cancer diagnosis. *J Am Med Assoc* 293:2479–2486.
 61. Pinto BM, Rabin C, Abdow S, et al. (2008) A pilot study on disseminating physical activity promotion among cancer survivors: a brief report. *Psychooncology* 17:517–521.
 62. Mustian KM, Katula JA, Gill DL, et al. (2004) Tai Chi Chuan, health-related quality of life and self-esteem: a randomized trial with breast cancer survivors. *Supp Care Cancer* 12: 871–876.
 63. Culos-Reed SN, Carlson LE, Daroux LM, et al. (2006) A pilot study of yoga for breast cancer survivors: physical and psychological benefits. *Psychooncology* 15:891–897.
 64. Moadel AB, Shah C, Wylie-Rosett J, et al. (2007) Randomized controlled trial of yoga among a multiethnic sample of breast cancer patients: effects on quality of life. *J Clin Oncol* 25: 4387–4395.
 65. Blake-Mortimer J, Gore-Felton C, Kimerling R, et al. (1999) Improving the quality and quantity of life among patients with cancer: a review of the effectiveness of group psychotherapy. *Eur J Cancer* 35:1581–1586.
 66. Courneya KS, Friedenreich CM, Sela RA, et al. (2003) The group psychotherapy and homebased physical exercise (group-hope) trial in cancer survivors: physical fitness and quality of life outcomes. *Psychooncology* 12:357–374.
 67. McNeely ML, Campbell KL, Rowe BH, et al. (2006) Effects of exercise on breast cancer patients and survivors: a systematic review and meta-analysis. *CMAJ* 175:34–41.
 68. Rehse B, Pukrop R. (2003) Effects of psychosocial interventions on quality of life in adult cancer patients: meta analysis of 37 published controlled outcome studies. *Patient Educ Counsel* 50:179–186.
 69. Stevinson C, Lawlor D, Fox KR. (2004) Exercise interventions for cancer patients: systematic review of controlled trials. *Cancer Cause Contr* 15:1035–1056.
 70. Oldervoll LM, Kaasa S, Hjermland MJ, et al. (2004) Physical exercise Results in the improved subjective well-being of a few or is effective rehabilitation for all cancer patients? *Eur J Cancer* 40:951–962.
 71. Schmitz KH, Holtzman J, Courneya KS, et al. (2005) Controlled physical activity trials in cancer survivors: a systematic review and meta-analysis. *Cancer Epidemiol Biomark Prev* 14:1588–1595.
 72. Midtgaard J, Stelter R, Rorth M, et al. (2007) Regaining a sense of agency and shared self-reliance: the experience of advanced disease cancer patients participating in a multidimensional exercise intervention while undergoing chemotherapy – analysis of patient diaries. *Scand J Psychol* 48:181–190.
 73. Midtgaard J, Rorth M, Stelter R, et al. (2006) The group matters: an explorative study of group cohesion and quality of life in cancer patients participating in physical exercise intervention during treatment. *Eur J Cancer Care* 15:25–33.
 74. Adamsen L, Rasmussen JM, Pedersen LS. (2001) Brothers in arms: how men with cancer experience a sense of comradeship through group intervention which combines physical activity with information relay. *J Clin Nurs* 10:528–537.
 75. Coon SK, Coleman EA. (2004) Keep moving: patients with myeloma talk about exercise and fatigue. *Oncol Nurs Forum* 31:1127.
 76. Stevinson C, Fox KR. (2006) Feasibility of an exercise rehabilitation programme for cancer patients. *Eur J Cancer Care* 15:386–396.
 77. Hennessy EM, Stevinson C, Fox KR. (2005) Preliminary study of the lived experience of exercise for cancer survivors. *Eur Onco Nurs* 9:155–166.
 78. Unruh AM, Elvin N. (2004) In the eye of the dragon: women's experience of breast cancer and the occupation of

- dragon boat racing. *Can J Occup Ther* 71:138–149.
79. Mitchell TL, Yakiwchuk CV, Griffin KL, et al. (2007) Survivor dragon boating: a vehicle to reclaim and enhance life after treatment for breast cancer. *Health Care Women Int* 28: 122–140.
 80. Sabiston CM, McDonough MH, Crocker PR. (2007) Psychosocial experiences of breast cancer survivors involved in a dragon boat program: exploring links to positive psychological growth. *J Sport Exerc Psychol* 29:419–438.
 81. Courneya KS, Mackay JR, Jones LW. (2000) Coping with cancer: can exercise help? *Phys Sports Med* 28:49–73.
 82. Doyle C, Kushi LH, Byers T, et al. (2006) Nutrition and physical activity during and after cancer treatment: an American Cancer Society guide for informed choices. *CA Cancer J Clin* 56:323–353.
 83. Humpel N, Iverson DC. (2005) Review and critique of the quality of exercise recommendations for cancer patients and survivors. *Supp Care Cancer* 13:493–502.
 84. Rogers LQ, Courneya KS, Shah P, et al. (2007) Exercise stage of change, barriers, expectations, values and preferences among breast cancer patients during treatment: a pilot study. *Eur J Cancer Care* 16:55–66.
 85. Eversley R, Estrin D, Dibble S, et al. (2005) Post-treatment symptoms among ethnic minority breast cancer survivors. *Oncol Nurs Forum* 32:250–256.
 86. Northouse LL, Caffey M, Deichelbohrer L, et al. (1999) The quality of life of African American women with breast cancer. *Res Nurs Health* 22:449–460.
 87. Roux L, Pratt M, Tengs TO, et al. (2008) Cost effectiveness of community-based physical activity interventions. *Am J Prev Med* 35:578–588.
-
1. IARC (International Agency of Cancer Research). (2009) <http://www-dep.iarc.fr/>. Accessed 21-1-2009.
 2. Bray F, McCarron P, Parkin DM. (2004) The changing global patterns of female breast cancer incidence and mortality. *Breast Cancer Res* 6:229–239.
 3. Kerlikowske K, Miglioretti DL, Buist DS, Walker R, Carney PA. (2007) Declines in invasive breast cancer and use of postmenopausal hormone therapy in a screening mammography population. *J Natl Cancer Inst* 99:1335–1339.
 4. Cancer Research UK. (2009) UK cancer mortality statistics. <http://info.cancerresearchuk.org/cancerstats/types/breast/mortality/>; Accessed 21-1-2009.
 5. Surveillance Epidemiology and End Results (SEER) Breast Cancer Statistics. (2009) <http://seer.cancer.gov/faststats/selections.php#Output>. Accessed 21-1-2009.
 6. Fentiman IS, Fourquet A, Hortobagyi GN. (2006) Male breast cancer. *Lancet* 367:595–604.
 7. Department of Health Statistics on Obesity Physical Activity and Diet: England. (2008) <http://www.ic.nhs.uk/webfiles/publications/opan08/OPAD%20Jan%202008%20final%20v7%20with%20links%20and%20buttons%20-%20NS%20logo%20removed%2020112008.pdf>. Accessed 13-11-2008.
 8. Dunn SE, Kari FW, French J, et al. (1997) Dietary restriction reduces insulin-like growth factor I levels, which modulates apoptosis, cell proliferation, and tumor progression in p53- deficient mice. *Cancer Res* 57:4667–4672.
 9. ATAC Trialists' Group. (2009) Personal Communication.
 10. Caan BJ, Emond JA, Natarajan L, et al. (2006) Post-diagnosis weight gain and breast cancer recurrence in women with early stage breast cancer. *Breast Cancer Res Treat* 99:47–57.
 11. Goodwin PJ, Panzarella T, Boyd NF. (1988) Weight gain in women with localized breast cancer – a descriptive study. *Breast Cancer Res Treat* 11:59–66.
 12. Harvie MN, Campbell IT, Baidam A, Howell A. (2004) Energy balance in early breast cancer patients receiving adjuvant chemotherapy. *Breast Cancer Res Treat* 83:201–210.
 13. Saquib N, Flatt SW, Natarajan L, et al. (2007) Weight gain and recovery of pre-cancer weight after breast cancer treatments: evidence from the women's healthy eating and living (WHEL) study. *Breast Cancer Res Treat* 105:177–186.
 14. Heideman WH, Russell NS, Gundy C, Rookus MA, Voskuil DW. (2009) The frequency, magnitude and timing of post-diagnosis body weight gain in Dutch breast cancer survivors. *Eur J Cancer* 45:119–126.
 15. Han HS, Lee KW, Kim JH, et al. (2009) Weight changes after adjuvant treatment in Korean women with early breast cancer. *Breast Cancer Res Treat* 114:147–153
 16. Harvie M, Howell A. (2006) Incorporating weight control into the management of patients with early breast cancer

- in the UK. In: Mctiernan A (ed) *Cancer Prevention and Management Through Exercise and Weight Control*. Marcel Dekker Inc., New York.
17. Malinowsky KM, Cameron D, Douglas S, et al. (2004) Breast cancer patients' experiences on endocrine therapy: monitoring with a checklist for patients on endocrine therapy (C-PET). *Breast* 13:363–368.
 18. Baum M, Budzar AU, Cuzick J, et al. (2002) Anastrozole alone or in combination with tamoxifen versus tamoxifen alone for adjuvant treatment of postmenopausal women with early breast cancer: first Results of the ATAC randomised trial. *Lancet* 359:2131–2139.
 19. Espeland MA, Stefanick ML, Kritz-Silverstein D, et al. (1997) Effect of postmenopausal hormone therapy on body weight and waist and hip girths. Postmenopausal Estrogen- Progestin Interventions Study Investigators. *J Clin Endocrinol Metabol* 82:1549–1556.
 20. Dawson-Hughes B, Harris S. (1992) Regional changes in body composition by time of year in healthy postmenopausal women. *Am J Clin Nutr* 56:307–313.
 21. Larosche I, Letteron P, Fromenty B, et al. (2007) Tamoxifen inhibits topoisomerases, depletes mitochondrial DNA, and triggers steatosis in mouse liver. *J Pharmacol Exp Ther* 321:526–535.
 22. Machinal-Quelin F, Dieudonne MN, Pecquery R, Leneuve MC, Giudicelli Y. (2002) Direct in vitro effects of androgens and estrogens on ob gene expression and leptin secretion in human adipose tissue. *Endocrine* 18:179–184.
 23. Francini G, Petrioli R, Montagnani A, et al. (2006) Exemestane after tamoxifen as adjuvant hormonal therapy in postmenopausal women with breast cancer: effects on body composition and lipids. *Br J Cancer* 95:153–158.
 24. Harvie MN, Howell A, Thatcher N, Baildam A, Campbell I. (2005) Energy balance in patients with advanced NSCLC, metastatic melanoma and metastatic breast cancer receiving chemotherapy – a longitudinal study. *Br J Cancer* 92:673–680.
 25. Carmichael AR, Bendall S, Lockerbie L, Prescott RJ, Bates T. (2004) Does obesity compromise survival in women with breast cancer? *Breast* 13:93–96.
 26. Kerlikowske K, Walker R, Miglioretti DL, Desai A, Ballard-Barbash R, Buist DS. (2008) Obesity, mammography use and accuracy, and advanced breast cancer risk. *J Natl Cancer Inst* 100:1724–1733.
 27. Daniell HW. (2009) Weight loss after breast cancer diagnosis may not improve prognosis. *J Clin Oncol* 27:829–830.
 28. Camoriano JK, Loprinzi CL, Ingle JN, Therneau TM, Krook JE, Veeder MH. (1990) Weight change in women treated with adjuvant therapy or observed following mastectomy for nodepositive breast cancer. *J Clin Oncol* 8:1327–1334.
 29. Kroenke CH, Chen WY, Rosner B, Holmes MD. (2005) Weight, weight gain, and survival after breast cancer diagnosis. *J Clin Oncol* 23:1370–1378.
 30. Hede K. (2008) Fat may fuel breast cancer growth. *J Natl Cancer Inst* 100:298–299.
 31. Chlebowski RT, Blackburn GL, Thomson CA, et al. (2006) Dietary fat reduction and breast cancer outcome: interim efficacy Results from the Women's Intervention Nutrition Study. *J Natl Cancer Inst* 98:1767–1776.
 32. McTiernan A, Rajan KB, Tworoger SS, et al. (2003) Adiposity and sex hormones in postmenopausal breast cancer survivors. *J Clin Oncol* 21:1961–1966.
 33. Dignam JJ, Wieand K, Johnson KA, Fisher B, Xu L, Mamounas EP. (2003) Obesity, tamoxifen use, and outcomes in women with estrogen receptor-positive early-stage breast cancer. *J Natl Cancer Inst* 95:1467–1476.
 34. Berclaz G, Li S, Price KN, et al. (2004) Body mass index as a prognostic feature in operable breast cancer: the International Breast Cancer Study Group experience. *Ann Oncol* 15: 875–884.
 35. Lethaby AE, Mason BH, Harvey VJ, Holdaway IM. (1996) Survival of women with node negative breast cancer in the Auckland region. *NZ Med J* 109:330–333.
 36. Lee CW, Norris B, Moravan V, Olivotto IA. (2003) Does body mass index influence the effectiveness of adjuvant tamoxifen in post menopausal women with early stage breast cancer? *Breast Cancer Res Treat* 82(Suppl 1):(Abstr 309).
 37. Lippman ME, Krueger KA, Eckert S, et al. (2001) Indicators of lifetime estrogen exposure: effect on breast cancer incidence and interaction with raloxifene therapy in the multiple outcomes of raloxifene evaluation study participants. *J Clin Oncol* 19:3111–3116.
 38. Fukuda M, Watanabe H, Suda T, Shimizu S. (1999) Sequential changes in hormone levels in postmenopausal breast cancer patients under long-term treatment with an aromatase inhibitor. Kanagawa AI Study Group. *Gan To Kagaku Ryoho* 26:2201–2208.
 39. Holmberg L, Tamini R. (2005) Reducing cancer drug doses in obese patients: dogma disputed. *Lancet* 366:1056–

40. Gladieff L, Fumoleau P, Kerbrat P, Bonneterre J, Mayer F, Namer M, Monnier A, Goudier MJ, Luporsi E, Roche H. (2004) Which are the interactions between baseline body mass index, toxicity of chemotherapy and efficacy of adjuvant treatment? 9-year follow up Results of French Adjuvant Study Group (FASG) trials. Proceedings for ASCO 40th Annual Meeting, New Orleans, Louisiana, 2004.
41. Dawood S, Broglio K, Gonzalez-Angulo AM, et al. (2008) Prognostic value of body mass index in locally advanced breast cancer. *Clin Cancer Res* 14:1718–1725.
42. Vona-Davis L, Rose DP. (2007) Adipokines as endocrine, paracrine, and autocrine factors in breast cancer risk and progression. *Endocr Relat Cancer* 14:189–206.
43. Daling JR, Malone KE, Doody DR, Johnson LG, Gralow JR, Porter PL. (2001) Relation of body mass index to tumor markers and survival among young women with invasive ductal breast carcinoma. *Cancer* 92:720–729.
44. Powis G, Reece P, Ahmann DL, Ingle JN. (1987) Effect of body weight on the pharmacokinetics of cyclophosphamide in breast cancer patients. *Cancer Chemother Pharmacol* 20:219–222.
45. Rodvold KA, Rushing DA, Tewksbury DA. (1988) Doxorubicin clearance in the obese. *J Clin Oncol* 6:1321–1327.
46. Williams G, Howell A, Jones M. (1988) The relationship of body weight to response to endocrine therapy, steroid hormone receptors and survival of patients with advanced cancer of the breast. *Br J Cancer* 58:631–634.
47. Artac M, Bozcuk H, Afacan B, Ozdogan M, Samur M. (2008) The impact of waist-to-hip ratio on clinical outcomes in metastatic breast cancer patients treated with aromatase inhibitors. *Breast* 17:418–422.
48. Borugian MJ, Sheps SB, Kim-Sing C, et al. (2003) Waist-to-hip ratio and breast cancer mortality. *Am J Epidemiol* 158:963–968.
49. Abrahamson PE, Gammon MD, Lund MJ, et al. (2006) General and abdominal obesity and survival among young women with breast cancer. *Cancer Epidemiol Biomark Prev* 15: 1871–1877.
50. Goodwin PJ. (2008) Insulin in the adjuvant breast cancer setting: a novel therapeutic target for lifestyle and pharmacologic interventions? *J Clin Oncol* 26:833–834.
51. Eccles D, Gerty S, Simmonds P, Hammond V, Ennis S, Altman DG. (2007) Prospective study of Outcomes in Sporadic versus Hereditary breast cancer (POSH): study protocol. *BMC Cancer* 7:160.
52. Hede K. (2008) Doctors seek to prevent breast cancer recurrence by lowering insulin levels. *J Natl Cancer Inst* 100:530–532.
53. Harvie MN, Malik R, Adams J, Frystyk J, Wale C, Howell A. (2004). The effects of weight loss and weight gain on IGF and IGF binding proteins in premenopausal women. *J Nutr* 134(12 suppl):(Conference abstract).
54. Irwin ML, McTiernan A, Bernstein L, et al. (2005) Relationship of obesity and physical activity with C-peptide, leptin, and insulin-like growth factors in breast cancer survivors. *Cancer Epidemiol Biomark Prev* 14:2881–2888.
55. Werner RS, McCormick B, Petrek J, et al. (1991) Arm edema in conservatively managed breast cancer: obesity is a major predictive factor. *Radiology* 180:177–184.
56. Shaw C, Mortimer P, Judd PA. (2007) A randomized controlled trial of weight reduction as a treatment for breast cancer-related lymphedema. *Cancer* 110:1868–1874.
57. Yancik R, Wesley MN, Ries LA, Havlik RJ, Edwards BK, Yates JW. (2001) Effect of age and comorbidity in postmenopausal breast cancer patients aged 55 years and older. *J Am Med Assoc* 285:885–892.
58. Manjer J, Kaaks R, Riboli E, Berglund G. (2001) Risk of breast cancer in relation to anthropometry, blood pressure, blood lipids and glucose metabolism: a prospective study within the Malmo Preventive Project. *Eur J Cancer Prev* 10:33–42.
59. Schairer C, Mink PJ, Carroll L, Devesa SS. (2004) Probabilities of death from breast cancer and other causes among female breast cancer patients. *J Natl Cancer Inst* 96: 1311–1321.
60. Barone BB, Yeh HC, Snyder CF, et al. (2008) Long-term all-cause mortality in cancer patients with preexisting diabetes mellitus: a systematic review and meta-analysis. *J Am Med Assoc* 300:2754–2764.
61. Tammemagi CM, Nerenz D, Neslund-Dudas C, Feldkamp C, Nathanson D. (2005) Comorbidity and survival disparities among black and white patients with breast cancer. *J Am Med Assoc* 294:1765–1772.
62. Hryniuk W. M, Banerji M. A, Heppner G, DuW, Severson R, Song EY. Adjuvant tamoxifen may be relatively ineffective in women with diabetes and breast cancer. *Proc Am Soc Clin Oncol* 20(36a):(Conference abstract).
63. de Waard F, Ramlaui R, Mulders Y, de Vries T, van Waveren S. (1993) A feasibility study on weight reduction in obese postmenopausal breast cancer patients. *Eur J Cancer Prev* 2: 233–238.
64. Garrow JS, Summerbell CD. (1995) Meta-analysis: effect of exercise, with or without dieting, on the body

- composition of overweight subjects. *Eur J Clin Nutr* 49:1–10.
65. Goodwin P, Espen MJ, Butler K, et al. (1998) Multidisciplinary weight management in locoregional breast cancer: Results of a phase II study. *Breast Cancer Res Treat* 48: 53–64.
 66. Loprinzi CL, Athmann LM, Kardinal CG, et al. (1996) Randomized trial of dietician counseling to try to prevent weight gain associated with breast cancer adjuvant chemotherapy. *Oncology* 53:228–232.
 67. Markes M, Brockow T, Resch KL. (2006) Exercise for women receiving adjuvant therapy for breast cancer. *Cochrane Database Syst Rev* CD005001.
 68. Ingram C, Courneya KS, Kingston D. (2006) The effects of exercise on body weight and composition in breast cancer survivors: an integrative systematic review. *Oncol Nurs Forum* 33:937–947.
 69. Demark-Wahnefried W, Case LD, Blackwell K, et al. (2008) Results of a diet/exercise feasibility trial to prevent adverse body composition change in breast cancer patients on adjuvant chemotherapy. *Clin Breast Cancer* 8:70–79.
 70. Demark-Wahnefried W, Clipp EC, Morey MC, et al. (2006) Lifestyle intervention development study to improve physical function in older adults with cancer: outcomes from Project LEAD. *J Clin Oncol* 24:3465–3473.
 71. Pirozzo S, Summerbell C, Cameron C, Glasziou P. (2003) Should we recommend low-fat diets for obesity? *Obes Rev* 4:83–90.
 72. Pierce JP, Natarajan L, Caan B, Parker B. A, Greenberg ER, Flatt S. W, Rock CL. (18 July 2007) Influence of a diet very high in vegetables, fruit and fiber and low in fat on prognosis following treatment for breast cancer. *J Am Med Assoc* 298(3):289–298.
 73. Bingham SA, Luben R, Welch A, Wareham N, Khaw KT, Day N. (2003) Are imprecise methods obscuring a relation between fat and breast cancer? *Lancet* 362:212–214.
 74. Foster GD, Wyatt HR, Hill JO, et al. (2003) A randomized trial of a low-carbohydrate diet for obesity. *N Engl J Med* 348:2082–2090.
 75. Shai I, Schwarzfuchs D, Henkin Y, et al. (2008) Weight loss with a low-carbohydrate Mediterranean, or low-fat diet. *N Engl J Med* 359:229–241.
 76. Pelkman CL, Fishell VK, Maddox DH, Pearson TA, Mauger DT, Kris-Etherton PM. (2004) Effects of moderate-fat (from monounsaturated fat) and low-fat weight-loss diets on the serum lipid profile in overweight and obese men and women. *Am J Clin Nutr* 79: 204–212.
 77. Prakash P, Krinsky NI, Russell RM. (2000) Retinoids, carotenoids, and human breast cancer cell cultures: a review of differential effects. *Nutr Rev* 58:170–176.
 78. Fowke JH, Longcope C, Hebert JR. (2000) Brassica vegetable consumption shifts estrogen metabolism in healthy postmenopausal women. *Cancer Epidemiol Biomark Prev* 9: 773–779.
 79. Rolls BJ, Roe LS, Meengs JS. (2004) Salad and satiety: energy density and portion size of a first-course salad affect energy intake at lunch. *J Am Diet Assoc* 104:1570–1576.
 80. Hamajima N, Hirose K, Tajima K, et al. (2002) Alcohol, tobacco and breast cancer – collaborative reanalysis of individual data from 53 epidemiological studies, including 58,515 women with breast cancer and 95,067 women without the disease. *Br J Cancer* 87:1234–1245.
 81. Rock CL, Demark-Wahnefried W. (2002) Nutrition and survival after the diagnosis of breast cancer: a review of the evidence. *J Clin Oncol* 20:3302–3316.
 82. Lindstrom J, Peltonen M, Eriksson JG, et al. (2006) High-fibre, low-fat diet predicts longterm weight loss and decreased type 2 diabetes risk: the Finnish Diabetes Prevention Study. *Diabetologia* 49:912–920.
 83. Holmes MD, Chen WY, Feskanich D, Kroenke CH, Colditz GA. (2005) Physical activity and survival after breast cancer diagnosis. *J Am Med Assoc* 293:2479–2486.
 84. Schwartz AL, Winters-Stone K, Gallucci B. (2007) Exercise effects on bone mineral density in women with breast cancer receiving adjuvant chemotherapy. *Oncol Nurs Forum* 34: 627–633.
 85. Saris WH, Blair SN van Baak MA, et al. (2003) How much physical activity is enough to prevent unhealthy weight gain? Outcome of the IASO 1st Stock Conference and consensus statement. *Obes Rev* 4:101–114.
 86. Laws R. (2004) Current approaches to obesity management in UK Primary Care: the Counterweight Programme. *J Hum Nutr Diet* 17:183–190.
 87. Renjilian DA, Perri MG, Nezu AM, McKelvey WF, Shermer RL, Anton SD. (2001) Individual versus group therapy for obesity: effects of matching participants to their treatment preferences. *J Consult Clin Psychol* 69:717–721.
 88. Djuric Z, DiLaura NM, Jenkins I, et al. (2002) Combining weight-loss counseling with the weight watchers plan for obese breast cancer survivors. *Obes Res* 10:657–665.

89. McBride CM, Clipp E, Peterson BL, Lipkus IM, Demark-Wahnefried W. (2000) Psychological impact of diagnosis and risk reduction among cancer survivors. *Psychooncology* 9:418–427.
90. Straw S, Hopwood P, Mitchell H, Harvie M. (2006) The need for lifestyle advice amongst breast cancer patients – A patient and health care professional survey. Fourth Year Option Medicine Project, University of Manchester, Manchester.
91. Avenell A, Broom J, Brown TJ, et al. (2004) Systematic review of the long-term effects and economic consequences of treatments for obesity and implications for health improvement. *Health Technol Assess* 8:iii–182.
92. Jebb SA, Siervo M, Murgatroyd PR, Evans S, Fruhbeck G, Prentice AM. (2007) Validity of the leg-to-leg bioimpedance to estimate changes in body fat during weight loss and regain in overweight women: a comparison with multi-compartment models. *Int J Obes (Lond)* 31:756–762.
93. Janssen I, Heymsfield SB, Allison DB, Kotler DP, Ross R. (2002) Body mass index and waist circumference independently contribute to the prediction of nonabdominal, abdominal subcutaneous, and visceral fat. *Am J Clin Nutr* 75:683–688.
94. Alberti KG, Zimmet P, Shaw J. (2006) Metabolic syndrome – a new world-wide definition. A Consensus Statement from the International Diabetes Federation. *Diabet Med* 23: 469–480.
95. World Health Organisation (WHO), National Union of Teachers NUT & National Council on Disability NCD. (1997) Obesity: preventing and managing the global epidemic. Report of WHO Consultation on Obesity, Geneva Switzerland.
96. Reid DM, Doughty J, Eastell R, et al. (2008) Guidance for the management of breast cancer treatment-induced bone loss: a consensus position statement from a UK Expert Group. *Cancer Treat Rev* 34:S3–S18.
97. Shapses SA, Riedt CS. (2006) Bone, body weight, and weight reduction: what are the concerns? *J Nutr* 136, 1453–1456.
98. Figueiredo MI, Cullen J, Hwang YT, Rowland JH, Mandelblatt JS. (2004) Breast cancer treatment in older women: does getting what you want improve your long-term body image and mental health? *J Clin Oncol* 22:4002–4009.
99. Guthrie JR, Taffe JR, Lehert P, Burger HG, Dennerstein L. (2004) Association between hormonal changes at menopause and the risk of a coronary event: a longitudinal study. *Menopause* 11:315–322.
100. Monnin S, Schiller MR, Sachs L, Smith AM. (1993) Nutritional concerns of women with breast cancer. *J Cancer Educ* 8:63–69.
101. McInnes JA, Knobf MT. (2001) Weight gain and quality of life in women treated with adjuvant chemotherapy for early-stage breast cancer. *Oncol Nurs Forum* 28:675–684.
102. Maciejewski ML, Patrick DL, Williamson DF. (2005) A structured review of randomized controlled trials of weight loss showed little improvement in health-related quality of life. *J Clin Epidemiol* 58:568–578.
103. Salminen E, Bishop M, Poussa T, Drummond R, Salminen S. (2004) Dietary attitudes and changes as well as use of supplements and complementary therapies by Australian and Finnish women following the diagnosis of breast cancer. *Eur J Clin Nutr* 58:137–144.
104. Demark-Wahnefried W, Morey MC, Clipp EC, et al. (2003) Leading the Way in Exercise and Diet (Project LEAD): intervening to improve function among older breast and prostate cancer survivors. *Control Clin Trials* 24:206–223.
105. Daley AJ, Bowden SJ, Rea DW, Billingham L, Carmicheal AR. (2008) What advice are oncologists and surgeons in the United Kingdom giving to breast cancer patients about physical activity? *Int J Behav Nutr Phys Act* 5:46.
106. Jones LW, Courneya KS, Peddle C, Mackey JR. (2005) Oncologists opinions towards recommending exercise to patients with cancer: a Canadian national survey. *Supp Care Cancer* 13:929–937.
107. Kreuter MW, Chheda SG, Bull FC. (2000) How does physician advice influence patient behavior? Evidence for a priming effect. *Arch Fam Med* 9:426–433.

1. ACS. (2006) Breast Cancer Facts and Figures: 2006–2007. American Cancer Society I, Atlanta.
2. Friedenreich CM. (2001) Physical activity and cancer prevention: from observational to intervention research. *Cancer Epidemiol Biomark Prev* 10(4):287–301.
3. Holmes MD, Chen WY, Feskanich D, Kroenke CH, Colditz GA. (2005) Physical activity and survival after breast cancer diagnosis. *J Am Med Assoc* 293(20):2479–2486.
4. Holick CN, Newcomb PA, Trentham-Dietz A, et al. (2008) Physical activity and survival after diagnosis of invasive breast cancer. *Cancer Epidemiol Biomark Prev* 17(2):379–386.
5. Irwin ML, Smith A, McTiernan A, et al. (2008) Influence of pre- and post-diagnosis physical activity on survival in breast cancer survivors: the health, eating, activity, and lifestyle (HEAL) study. *J Clin Oncol* 26(24):1–7.
6. Pierce JP, Stefanick ML, Flatt SW, et al. (2007) Greater survival after breast cancer in physically active women with high vegetable-fruit intake regardless of obesity. *J Clin Oncol* 25(17):2345–2351.
7. Calle EE, Rodriguez C, Walker-Thurmond K, Thun MJ. (2003) Overweight, obesity, and mortality from cancer in a prospectively studied cohort of US adults. *NEJM* 348(17):1625–1638.
8. Daling JR, Malone KE, Doody DR, Johnson LG, Gralow JR, Porter PL. (2001) Relation of body mass index to tumor markers and survival among young women with invasive ductal breast carcinoma. *Cancer* 92(4):720–729.
9. Kroenke CH, Chen WY, Rosner B, Holmes MD. (2005) Weight, weight gain, and survival after breast cancer diagnosis. *J Clin Oncol* 23(7):1370–1378.
10. Irwin ML, McTiernan A, Baumgartner RN, et al. (2005) Changes in body fat and weight after a breast cancer diagnosis: influence of demographic, prognostic, and lifestyle factors. *J Clin Oncol* 23(4):774–782.
11. Caan BJ, Emond JA, Natarajan L, et al. (2006) Post-diagnosis weight gain and breast cancer recurrence in women with early stage breast cancer. *Breast Cancer Res Treat* 99(1):47–57.
12. Irwin ML, Yasui Y, Ulrich CM, et al. (2003) Effect of exercise on total and intra-abdominal body fat in postmenopausal women: a randomized controlled trial. *J Am Med Assoc* 289(3):323–330.
13. Irwin ML, Alvarez-Reeves M, Cadmus L, et al. 2009 Randomized controlled trial of exercise on body fat, lean mass, and bone mineral density in breast cancer survivors (Under review)
14. Schmitz KH, Ahmed RL, Hannan PJ, Yee D. (2005) Safety and efficacy of weight training in recent breast cancer survivors to alter body composition, insulin, and insulin-like growth factor axis proteins. *Cancer Epidemiol Biomark Prev* 14(7):1672–1680.
15. Goodwin PJ, Ennis M, Pritchard KI, et al. (2002) Fasting insulin and outcome in early-stage breast cancer: Results of a prospective cohort study. *J Clin Oncol* 20(1):42–51.
16. Irwin ML, McTiernan A, et al. (2007) Associations between fasting c-peptide and breast cancer death: the Health, Eating, Activity, and Lifestyle (HEAL) Study. AACR Frontiers in Cancer Prevention Meeting, December 2007, Abstract B99.
17. Pollak MNCJ, Shepherd L, et al. (2006) Insulin resistance, estimated by serum C-peptide level, is associated with reduced event-free survival for postmenopausal women in NCIC CTG MA.14 adjuvant breast cancer trial. *J Clin Oncol* 24:(suppl 18S, abstr 524).
18. Gray SG, Stenfeldt Mathiasen I, De Meyts P. (2003) The insulin-like growth factors and insulin-signalling systems: an appealing target for breast cancer therapy?. *Horm Metab Res* 35(11-12):857–871.
19. Yu H, Rohan T. (2000) Role of the insulin-like growth factor family in cancer development and progression. *J Natl Cancer Inst* 92(18):1472–1489.
20. Hankinson S, Willett WC, Colditz GA, Hunter DJ, Michaud DS, Deroo B, et al. (1998) Circulating concentrations of insulin-like growth factor-1 and risk of breast cancer. *Lancet* 351:1393–1396.
21. Goodwin PJ, Ennis M, Pritchard KI, et al. (2002) Insulin-like growth factor binding proteins 1 and 3 and breast cancer outcomes. *Breast Cancer Res Treat* 74(1):65–76.
22. Ligibel JA, Campbell N, Partridge A, et al. (2008) Impact of a mixed strength and endurance exercise intervention on insulin levels in breast cancer survivors. *J Clin Oncol* 26(6):907–912.
23. Irwin ML, Varma K, Cadmus L, et al. (2009) Randomized controlled trial of exercise on insulin and IGFs in breast cancer survivors. *Cancer Epidemiol Biomark Prev* 18(1):306–313.
24. Fairey AS, Courneya KS, Field CJ, Bell GJ, Jones LW, Mackey JR. (2003) Effects of exercise training on fasting insulin, insulin resistance, insulin-like growth factors, and insulin-like growth factor binding proteins in postmenopausal breast cancer survivors: a randomized controlled trial. *Cancer Epidemiol Biomark Prev* 12(8):721–727.

25. Frank LL, Sorensen BE, Yasui Y, et al. (2005) Effects of exercise on metabolic risk variables in overweight postmenopausal women: a randomized clinical trial. *Obes Res* 13(3):615–.
26. Goodyear L. (1998) KBE, glucose transport, and insulin sensitivity. *Ann Rev Med* 49: 235–261.
27. Key T, Appleby P, Barnes I, Reeves G. (2002) Endogenous sex hormones and breast cancer in postmenopausal women: reanalysis of nine prospective studies. *J Natl Cancer Inst* 94(8): 606–616.
28. Howell A, Cuzick J, Baum M, et al. (2005) Results of the ATAC (arimidex, tamoxifen, alone or in combination) trial after completion of 5 years of adjuvant treatment for breast cancer. *Lancet* 365(9453):60–62.
29. Thurlimann B, Keshaviah A, Coates AS, et al. (2005) A comparison of letrozole and tamoxifen in postmenopausal women with early breast cancer. *N Engl J Med* 353(26):2747–2757.
30. McTiernan A, Tworoger SS, Ulrich CM, et al. (2004) Effect of exercise on serum estrogens in postmenopausal women: a 12-month randomized clinical trial. *Cancer Res* 64(8):2923–2928.
31. McTiernan A, Wu L, Chen C, et al. (2006) Relation of BMI and physical activity to sex hormones in postmenopausal women. *Obesity* 14(9):1662–1677.
32. McTiernan A, Rajan KB, Tworoger SS, et al. (2003) Adiposity and sex hormones in postmenopausal breast cancer survivors. *J Clin Oncol* 21(10):1961–1966.
33. Irwin ML, Alvarez-Reeves M, Cadmus L, et al. (2008) RCT of exercise on sex hormones in breast cancer survivors. AACR Frontiers in Cancer Prevention Meeting, December, 2008.
34. Vona-Davis L. (2007) RDA, paracrine, and autocrine factors in breast cancer risk and progression. *Endocrine-Related Cancer* 14:189–206.
35. Jarde T, Caldefie-Chezet F, Damez M, et al. (2008) Leptin and leptin receptor involvement in cancer development: a study on human primary breast carcinoma. *Oncol Rep* 19(4):905–911.
36. Housa DHJ, Vernerova Z, Haluzik M. (2006) Adipocytokines and cancer. *Physiol Res* 55: 233–244.
37. Nicolini A, Carpi A, Rossi G. (2006) Cytokines in breast cancer. *Cytokine Growth Factor Rev* 17:325–337.
38. Irwin ML, McTiernan A, Bernstein L, et al. (2005) Relationship of obesity and physical activity with C-peptide, leptin, and insulin-like growth factors in breast cancer survivors. *Cancer Epidemiol Biomark Prev* 14(12):2881–2888.
39. Miyoshi YFT, Kihara S, Taguchi T, Tamaki Y, Matsuzawa Y, Noguchi S. (2003) Association of serum adiponectin levels with breast cancer risk. *Clin Cancer Res* 9:5699–5704.
40. Arditi JD, Venihaki M, Karalis KP, Chrousos GP. (2007) AnTtiproliferative effect of adiponectin on MCF 7 breast cancer cells: a potential hormonal link between obesity and cancer. *Horm Metabol Res* 39(11):9–13.
41. Miyoshi Y, Funahashi T, Kihara S, et al. (2003) Association of serum adiponectin levels with breast cancer risk. *Clin Cancer Res* 9(15):5699–5704.
42. Irwin ML, Alvarez-Reeves M, Cadmus L, et al. (2009) RCT of exercise on leptin and adiponectin in breast cancer survivors. *Obesity* (Under review).
43. Ardizzioia ALP, Brivio F, et al. (1992) Tumor necrosis factor in solid tumors: increased blood levels in metastatic disease. *J Biol Regul Homeost Agents* 6:103.
44. Pierce BL, Neuhouser ML, Wener MH, et al. (10 April 2008) Correlates of circulating C-reactive protein and serum amyloid A concentrations in breast cancer survivors. *Breast Cancer Res Treat* 114:155–167.
45. Zhang SM, Lin J, Cook NR, et al. (2007) C-reactive protein and risk of breast cancer. *J Natl Cancer Inst* 99(11):890–894.
46. DeNardo DG, Coussens LM. (2007) Inflammation and breast cancer. Balancing immune response: crosstalk between adaptive and innate immune cells during breast cancer progression. *Breast Cancer Res* 9(4):212.
47. McMillan DCEM, Sattar N, Angerson WJ, Johnstone J, McArdle CS. (2001) Measurement of the systematic inflammatory response predicts cancer-specific and non-cancer survival in patients with cancer. *Nutr Cancer* 41(1&2):64–69.
48. Garcia-Tunon IRM, Ruiz A, Fraile B, Paniagua R, Royuela M. (2006) Role of tumor necrosis factor- α and its receptors in human benign breast lesions and tumors (in situ and infiltrative). *Cancer Sci* 97:1044–1049.
49. Knupfer H, Preib R. (2007) Significance of interleukin-6 (IL-6) in breast cancer (review). *Breast Cancer Res Treat* 102:129–135.
50. Mohamed-Ali VGS, Rawesh A, et al. (1997) Subcutaneous adipose tissue releases interleukin-6, but not tumor necrosis factor- α in vivo. *J Clin Endocrinol Metabol* 82:4196–4200.
51. Fairey AS, Courneya KS, Field CJ, et al. (2005) Effect of exercise training on C-reactive protein in postmenopausal

- breast cancer survivors: a randomized controlled trial. *Brain Behav Immun* 19(5):381–388.
52. Jones S, Hesselsweet S, Yu H, Irwin ML. (2009) Randomized controlled trial of exercise on inflammation in breast cancer survivors. *J Cancer Survivorship* (Under review).
 53. Fischer CPBA, Perstrup LB, Eskildsen P, Pedersen BK. (2007) Plasma levels of interleukin-6 and c-reactive protein are associated with physical inactivity independent of obesity. *Scand J Med Sci Sports* 17:580–587.
 54. Olson TPDD, Leon AS, Schmitz KH. (2007) Changes in inflammatory biomarkers following one-year of moderate resistance training in overweight women. *Int J Obesity* 31:996–1003.
 55. Dekker MJLS, Hudson R, et al. (2007) An exercise intervention without weight loss decreases circulating interleukin-6 in lean and obese men with and without type 2 diabetes mellitus. *Metabol Clin Exp* 56:332–338.
 56. Kaspis C, Thompson PD. (2005) The effects of physical activity on serum c-reactive protein and inflammatory markers. *J Am College Cardiol* 45(10):1563–1569.
 57. Hursting SD, Lashinger LM, Colbert LH, et al. (2007) Energy balance and carcinogenesis: underlying pathways and targets for intervention. *Curr Cancer Drug Targets* 7(5):484–491.
 58. Fahey AS, Courneya KS, Field CJ, Mackey JR. (2002) Physical exercise and immune system function in cancer survivors: a comprehensive review and future directions. *Cancer* 94(2): 539–551.
 59. Fahey AS, Courneya KS, Field CJ, Bell GJ, Jones LW, Mackey JR. (2005) Randomized controlled trial of exercise and blood immune function in postmenopausal breast cancer survivors. *J Appl Physiol* 98(4):1534–1540.

1. American Cancer Society, *Cancer Facts & Figures 2008*, 1–72, 2008.
2. Canadian Cancer Society/National Cancer Institute of Canada. (2008) *Canadian Cancer Statistics*, pp. 1–72.
3. Quinn M, Babb P. (2002) Patterns and trends in prostate cancer incidence, survival, prevalence and mortality. Part I: international comparisons. *Br J Urol Int* 90:162–173.
4. Hsing A, Tsao WL, Devesa SS. (1999) International trends and patterns of prostate cancer incidence and mortality. *Int J Cancer* 85:60–67.
5. National Comprehensive Cancer Network, *NCCN Clinical Practice Guidelines in Oncology: Prostate Cancer V.I. 2008*, 1–44, 2008.
6. Fowler FJ, Barry MJ, Lu-Yao G, Wasson J, Roman A, Wennberg J. (1995) Effect of radical prostatectomy for prostate cancer on patient quality of life: Results from a medicare survey. *Urology* 45:1007–1115.
7. Moore KN, Estey A. (1999) The early post-operative concerns of men after radical prostatectomy. *J Adv Nurs* 29:1121–1129.
8. Haythornthwaite JA, Raja SN, Fisher B, Frank SM, Brendler CB, Shir Y. (1998) Pain and quality of life following radical retropubic prostatectomy. *J Urol* 160: 1761–1764.
9. Meuleman E. (2003) Erectile function after radical prostatectomy: a review. *Eur Urol* 43: 95–102.
10. Peschel RE, Colberg JW. (2003) Surgery, brachytherapy, and external-beam radiotherapy for early prostate cancer. *Lancet Oncol* 4:233–241.
11. Litwin MS, Hays RD, Fink A, Ganz PA, Leake B, Leach GE, Brook RH. (1995) Quality-of-life outcomes in men treated for localized prostate cancer. *JAMA* 273:129–135.
12. Helgason AR, Adolfsson J, Dickman P, Fredrikson M, Arver S, Steineck G. (1996) Waning sexual function – the most important disease-specific distress for patients with prostate cancer. *Br J Cancer* 73:1417–1421.
13. Klotz L, Kiberl AS, Sanda M, Thompson I, Clarke N (2008) Surveillance Therapy Against Radical Treatment Trial (START), September 5, 2008.
14. Kennedy TJ, Sonneland AM, Marlett MM, Troup RH. (1992) Luteinizing hormone-releasing hormone downstaging of clinical stage C prostate cancer. *J Urol* 147:891–893.
15. Macfarlane MT, Abi-Aad A, Stein A, Danella J, Beldegrum A, deKernion JB. (1993) Neoadjuvant hormonal deprivation in patients with locally advanced prostate cancer. *J Urol* 150:132–134.
16. Oesterling JE, Andrews PE, Suman VJ, Zincke H, Myers RP. (1993) Preoperative androgen deprivation therapy: artificial lowering of serum prostate specific antigen without downstaging the tumor. *J Urol* 149:779–782.
17. Strum SE, McDermed JE, Scholz MC, Johnson H, Tisman G. (1997) Anaemia associated with androgen deprivation in patients with prostate cancer receiving combined hormone blockade. *Br J Urol* 79:933–941.
18. Daniell HW, Dunn SR, Ferguson DW, Lomas G, Niazi Z, Stratte PT. (2000) Progressive osteoporosis during androgen deprivation therapy for prostate cancer. *J Urol* 163: 181–186.

19. Smith MR, Finkelstein JS, McGovern FJ, Zietman AL, Fallon MA, Schoenfeld DA, Kantoff PW. (2002) Changes in body composition during androgen deprivation therapy for prostate cancer. *J Clin Endocrinol Metab* 87:599–603.
20. Mittan D, Lee S, Miller E, Perez RC, Basler JW, Bruder JM. (2002) Bone loss following hypogonadism in men with prostate cancer treated with GnRH analogs. *J Clin Endocrinol Metab* 87:3656–3661.
21. Alibhai SMH, Gogov S, Alibhai Z. (2006) Long-term side effects of androgen deprivation therapy in men with non-metastatic prostate cancer: a systematic literature review. *Crit Rev Oncol Hematol* 60:201–215.
22. Chander S, Choo R, Danjoux C, Morton G, Pearse A, Deboer G, Szumacher E, Loblaw A, Cheung P, Woo T. (2005) Effect of androgen suppression on hemoglobin in prostate cancer patients undergoing salvage radiotherapy plus 2-year buserelin acetate for rising PSA after surgery. *Int J Radiat Oncol Biol Phys* 62:719–724.
23. Stoch SA, Parker RA, Chen L, Bublely G, Ko Y, Vincelle A, Greenspan SL. (2001) Bone loss in men with prostate cancer treated with gonadotropin-releasing hormone agonists. *J Clin Endocrinol Metab* 86:2787–2791.
24. Preston DM, Torrens JI, Harding P, Howard RS, Duncan WE, Mcleod DG. (2002) Androgen deprivation in men with prostate cancer is associated with an increased rate of bone loss. *Prost Cancer Prost Dis* 5:304–310.
25. Pickett M, Bruner DW, Joseph A, Burgraf V. (2000) Prostate cancer elder alert: living with treatment choices and outcomes. *J Gerontol Nurs* 26:22–34.
26. Cassileth BR, Soloway MS, Vogelzang NJ, Chou JM, Schellhammer PD, Seidmon EJ, Kennealey GT. (1992) Quality of life and psychosocial status in stage D prostate cancer. *Qual Life Res* 1:323–330.
27. Cleary PD, Morrissey G, Oster G. (1995) Health-related quality of life in patients with advanced prostate cancer. *Qual Life Res* 4:207–220.
28. Alibhai SMH, Rahman S, Warde PR, Jewett MA, Jaffer T, Cheung AM. (2006) Prevention and management of osteoporosis in men receiving androgen deprivation therapy: a survey of urologists and radiation oncologists. *Urology* 68:126–131.
29. Basaria S, Lieb JI, Tang AM, DeWeese T, Carducci M, Eisenberger M, Dobs AS. (2002) Long-term effects of androgen deprivation therapy in prostate cancer patients. *Clin Endocrin* 56:779–786.
30. Soyupek F, Soyupek H, Ozorak A. (2008) Androgen deprivation therapy for prostate cancer: effects on hand function. *Urol Oncol* 26:141–146.
31. Mottet N, Prayer-Galetti T, Hammerer P, Kattan MW, Tunn U. (2006) Optimizing outcomes and quality of life in the hormonal treatment of prostate cancer. *Br J Urol Int* 98:20–27.
32. Tannock IF, de Wit R, Berry WR, Horti J, Pluzanska A, Chi KN, Oudard S, Theodore C, James ND, Turesson I, Rosenthal MA, Eisenberger MA. (2004) Docetaxel plus prednisone or mitoxantrone plus prednisone for advanced prostate cancer. *N Engl J Med* 351:1502–1512.
33. Berthold DR, Pond GR, Roessner M, deWit R, Eisenberger M, Tannock IF. (2008) Treatment of hormone-refractory prostate cancer with docetaxel or mitoxantrone: relationships between prostate-specific antigen, pain, and quality of life response and survival in the TAX-327 study. *Clin Cancer Res* 14:2763–2767.
34. Picus J, Schultz M. (1999) Docetaxel (taxotere) as monotherapy in the treatment of hormonerefractory prostate cancer: preliminary Results. *Semin Oncol* 26:14–18.
35. Casperson CJ, Powell KE, Christenson GM. (1985) Physical activity, exercise, and physical fitness: definition and distinctions for health-related research. *Public Health Rep* 100: 126–131.
36. Bouchard C, Shephard RJ, Physical Activity, Fitness, and Health: International Proceedings and Consensus Statement. Physical activity, fitness, and health: the model and key concepts, Windsor, ON: Human Kinetics, 1994, 77–88.
37. Paffenbarger RS, Hyde RT, Wing AL, Hsieh CC. (1986) Physical activity, all-cause mortality, and longevity of college alumni. *N Engl J Med* 314:605–613.
38. Warburton DER, Nicol CW, Bredin SSD. (2006) Health benefits of PA: the evidence. *Can Med Assoc J* 174:801–809.
39. Kampert JB, Blair SN, Barlow CE, Kohl HW III, Harold W. (1996) Physical activity, physical fitness, and all-cause and cancer mortality: a prospective study of men and women. *Ann Epidemiol* 6:452–457.
40. Friedenreich CM, Thune I. (2001) A review of PA and prostate cancer risk. *Cancer Causes Control* 12:461–475.
41. Min LI-. (2003) Physical activity and cancer prevention-data from epidemiologic studies. *Med Sci Sports Exerc* 35:1823–1827.
42. Giovannucci EL, Liu Y, Leitzmann MF, Stampfer MJ, Willett WC. (2005) A prospective study of PA and incident and fatal prostate cancer. *Arch Intern Med* 165:1005–1010.

43. Tymchuk CN, Barnard RJ, Heber D, Aronson WJ. (2001) Evidence of an inhibitory effect of diet and exercise on prostate cancer cell growth. *J Urol* 166:1185–1189.
44. Ngo TH, Barnard RJ, Tymchuk CN, Cohen P, Aronson WJ. (2002) Effect of diet and exercise on serum insulin, IGF-I, and IGFBP-1 levels and growth of LNCaP cells in vitro. *Cancer Causes Control* 13:929–935.
45. Galvão DA, Taaffe DR, Spry N, Newton RU. (2007) Exercise can prevent and even reverse adverse effects of androgen suppression treatment in men with prostate cancer. *Prost Cancer Prost Dis* 8:1–7.
46. Ngo TH, Barnard RJ, Leung P, Cohen P, Aronson WJ. (2003) Insulin-like growth factor (IGFI) and IGF binding protein-1 modulate prostate cancer cell growth and apoptosis: possible mediators for the effects of diet and exercise on cancer cell survival. *Endocrinology* 144: 2319–2324.
47. Barnard RJ, Ngo TH, Leung PS, Aronson WJ, Golding LA. (2003) A low-fat diet and/or strenuous exercise alters the IFG axis in vivo and reduces prostate tumor cell growth in vitro. *Prostate* 56:201–206.
48. Leung PS, Aronson WJ, Ngo TH, Golding LA, Barnard RJ. (2004) Exercise alters the IGF axis in vivo and increases p53 protein in prostate tumor cells in vitro. *J Appl Physiol* 96:450–454.
49. Tymchuk CN, Tessier SB, Aronson WJ, Barnard RJ. (1998) Effects of diet and exercise on insulin, sex hormone-binding globulin, and prostate-specific antigen. *Nutr Cancer* 31: 127–131.
50. Barnard RJ, Leung PS, Aronson WJ, Cohen P, Golding LA. (2007) A mechanism to explain how regular exercise might reduce the risk for clinical prostate cancer. *Eur J Cancer Prev* 16:415–421.
51. Segal RJ, Reid RD, Courneya KS, Malone SC, Parliament MB, Scott CG, Venner PM, Quinney HA, Jones LW, D'Angelo ME, Wells GA. (May 1. 2003) Resistance exercise in men receiving androgen deprivation therapy for prostate cancer. *J Clin Oncol* 21: 1653–1659.
52. Segal RJ, Reid R, Proulx C, Courneya K, Sigal R, Kenny G, Prud'homme S, Wells MG, (2009) Randomized controlled trial of resistance or aerobic exercise in men receiving radiation for prostate cancer. *Journal of Clinical Oncology*. 27: 344–351.
53. Galvao DA, Nosaka K, Taaffe DR, Spry N, Kristjanson LJ, McGuigan MR, Suzuki K, Kanemitsu Y, Newton RU. (2006) Resistance training and reduction of treatment side effects in prostate cancer patients. *Med Sci Sports Exerc* 38:2045–2052.
54. Duke University Medical Center. (2008) Exercise May Lead to Faster Prostate Tumor Growth, April 14, 2008.
55. Thorsen L, Courneya KS, Stevinson C, Fossa SD. (2008) A systematic review of PA in prostate cancer survivors: outcomes, prevalence, and determinants. *Supportive Care Cancer* 16:987–997.
56. Durak EP, Harris J, Ceriale SM. (2001) The effects of exercise on quality of life improvement in cancer survivors: the Results of a national survey. *J Exerc Online* 4:21–28.
57. Stevinson C, Lawlor D, Fox K. (2004) Exercise interventions for cancer patients: systematic review of controlled trials. *Cancer Causes Control* 15:1035–1056.
58. Freedland SJ, Krupski TL, Moul JW. (2006) Update on outcomes research databases in prostate cancer 2006. *Curr Opin Urol* 16:168–172.
59. Freedland SJ, Aronson WJ, Kane CJ, Presti JC, Amling CL, Elashoff D, Terris MK. (2004) Impact of obesity on biochemical control after radical prostatectomy for clinically localized prostate cancer: a report by the shared equal access regional cancer hospital database study group. *J Clin Oncol* 22:446–453.
60. Chang SS, Duong DT, Wells N, Cole EE, Smith JA, Cookson MS. (2004) Predicting blood loss and transfusion requirements during radical prostatectomy: the significant negative impact of increasing body mass index. *J Urol* 171:1861–1865.
61. Windsor PM, Nicol KF, Potter J. (2004) A randomized, controlled trial of aerobic exercise for treatment-related fatigue in men receiving radical external beam radiotherapy for localized prostate carcinoma. *Cancer* 101:550–557.
62. Monga U, Garber SL, Thornby J, Vallbona C, Kerrigan A, Monga TN, Zimmerman KP. (2007) Exercise prevents fatigue and improves quality of life in prostate cancer patients undergoing radiotherapy. *Arch Phys Med Rehabil* 88:1416–1422.
63. Smith JA, Jr. Soloway MS, Young MJ. (1999) Complications of advanced prostate cancer. *Urology* 54:8–14.
64. Williams MB, Hernandez J, Thompson I. (2005) Luteinizing hormone-releasing hormone agonist effects on skeletal muscle: how hormonal therapy in prostate cancer affects muscular strength. *J Urol* 173:1067–1071.
65. Clay CA, Perera S, Wagner JM, Miller ME, Nelson JB, Greenspan SL. (2007) Physical function in men with prostate cancer on androgen deprivation therapy. *Phys Ther* 87:1325–1333.
66. Breunis H, Leach M, Naglie G, Tannock IF, Fleshner NE, Krahn M, Duff-Canning S, Tomlinson G, Warde P,

- Alibhai SM. (2008) Androgen deprivation therapy (ADT) and physical function in men with nonmetastatic prostate cancer. *J Clin Oncol* 27:4895.
67. Potosky AL, Reeve BB, Clegg LX, Hoffman RM, Stephenson RA, Albertsen PC, Gilliland FD, Stanford JL. (2002) Quality of life following localized prostate cancer treated initially with androgen deprivation therapy or no therapy. *J Nat Cancer Inst* 94:430–437.
 68. Daniell HW. (1997) Osteoporosis after orchiectomy for prostate cancer. *J Urol* 157:439–444.
 69. Lee H, McGovern K, Finkelstein JS, Smith MR. (2005) Changes in bone mineral density and body composition during initial and long-term gonadotropin-releasing hormone agonist treatment for prostate carcinoma. *Cancer* 104:1633–1637.
 70. Shahinian VB, Yong-Fang K, Freeman JL, Goodwin JS. (2005) Risk of fracture after androgen deprivation for prostate cancer. *N Engl J Med* 352:154–164.
 71. Oeffelein MG, Ricchuiti V, Conrad W, Resnick M. (2002) Skeletal fractures negatively correlate with overall survival in men with prostate cancer. *J Urol* 168:1005–1007.
 72. Carmack Taylor CL, Demoor C, Smith MA, Dunn AL, Basen-Engquist K, Nielsen I, Pettaway C, Sellin R, Massey P, Gritz ER. (2006) Active for life after cancer: A randomized trial examining a lifestyle PA program for prostate cancer patients. *Psycho-Oncology* 15:847–862.
 73. Prochaska JO, Velicer WF. (1997) The transtheoretical model of health behavior change. *Am J Health Prom* 12:38–48.
 74. Prochaska JO, DiClemente CC. (1983) Stages and processes of self-change of smoking: toward an integrative model of change. *J Consult Clin Psychol* 51:390–395.
 75. Bandura A. (1986) *Social foundations of thought and action: a social cognitive theory*. Prentice Hall, Englewood Cliffs, NJ.
 76. Bandura A. (1997) *Self-efficacy: the exercise of control*. W.H. Freeman and Company, New York.
 77. Culos-Reed SN, Robinson JL, Lau H, O'Connor K, Keats MR. (2007) Benefits of a PA intervention for men with prostate cancer. *J Sport Exerc Psych* 29:118–127.
 78. Courneya KS, Friedenreich CM. (Fall 1997) Relationship between exercise pattern across the cancer experience and current quality of life in colorectal cancer survivors. *J Altern Complement Med* 3:215–226.
 79. Culos-Reed SN, Robinson JW, Lau H, Stephenson L, Keats M, Norris S, Kline G. (2007) Physical activity for men receiving androgen deprivation therapy for prostate cancer: benefits from a 16-week intervention *Support Caner Cancer*. 17 (online).
 80. Berglund G, Bolund C, Gustafsson UL, Sjoden PO. (1994) A randomized study of a rehabilitation program for cancer patients: the 'starting again' group. *Psycho-Oncology* 3:109–120.
 81. Berglund G, Bolund C, Gustafsson UL, Sjoden PO. (1994) One-year follow-up of the 'starting again' group rehabilitation programme for cancer patients. *Eur J Cancer* 30A:1744–1751.
 82. Knols R, Aaronson NK, Uebelhart D, Franssen J, Aufdemkampe G. (2005) Physical exercise in cancer patients during and after medical treatment: A systematic review of randomized and controlled clinical trials. *J Clin Oncol* 23:3830–3842.
 83. Courneya KS, Segal RJ, Reid RD, Jones LW, Malone SC, Venner PM, Parliament MB, Scott CG, Quinney HA, Wells GA. (2004) Three independent factors predicted adherence in a randomized controlled trial of exercise training among prostate cancer survivors. *J Clin Epidem* 57:571–579.
1. Thune I, Furberg AS. (2001) Physical activity and cancer risk: dose-response and cancer, all sites and site-specific. *Med Sci Sports Exerc* 33:S530–S550.
 2. Lee IM. (2003) Physical activity and cancer prevention – data from epidemiologic studies. *Med Sci Sports Exerc* 35:1823–1827.
 3. Segal RJ, Reid RD, Courneya KS, et al. (2003) Resistance exercise in men receiving androgen deprivation therapy for prostate cancer. *J Clin Oncol* 21:1653–1659.
 4. Galvao DA, Taaffe DR, Spry N, Newton RU. (2007) Exercise can prevent and even reverse adverse effects of androgen suppression treatment in men with prostate cancer. *Prostate Cancer Prostatic Dis* 10:340–346.
 5. Windsor PM, Nicol KF, Potter J. (2004) A randomized, controlled trial of aerobic exercise for treatment-related

- fatigue in men receiving radical external beam radiotherapy for localized prostate carcinoma. *Cancer* 101:550–557.
6. Giovannucci E, Leitzmann M, Spiegelman D, et al. (1998) A prospective study of physical activity and prostate cancer in male health professionals. *Cancer Res* 58:5117–5122.
 7. Giovannucci EL, Liu Y, Leitzmann MF, Stampfer MJ, Willett WC. (2005) A prospective study of physical activity and incident and fatal prostate cancer. *Arch Intern Med* 165:1005–1010.
 8. Hu G, Tuomilehto J, Silventoinen K, Barengo NC, Peltonen M, Jousilahti P. (2005) The effects of physical activity and body mass index on cardiovascular, cancer and all-cause mortality among 47 212 middle-aged Finnish men and women. *Int J Obes (Lond)* 29:894–902.
 9. Leitzmann MF, Park Y, Blair A, et al. (2007) Physical activity recommendations and decreased risk of mortality. *Arch Intern Med* 167:2453–2460.
 10. Orsini N, Mantzoros CS, Wolk A. (2008) Association of physical activity with cancer incidence, mortality, and survival: a population-based study of men. *Br J Cancer* 98:1864–1869.
 11. Ngo TH, Barnard RJ, Tymchuk CN, Cohen P, Aronson WJ. (2002) Effect of diet and exercise on serum insulin, IGF-I, and IGFBP-1 levels and growth of LNCaP cells in vitro (United States). *Cancer Causes Control* 13:929–935.
 12. Tymchuk CN, Barnard RJ, Heber D, Aronson WJ. (2001) Evidence of an inhibitory effect of diet and exercise on prostate cancer cell growth. *J Urol* 166:1185–1189.
 13. Tymchuk CN, Tessler SB, Aronson WJ, Barnard RJ. (1998) Effects of diet and exercise on insulin, sex hormone-binding globulin, and prostate-specific antigen. *Nutr Cancer* 31: 127–131.
 14. Tymchuk CN, Barnard RJ, Ngo TH, Aronson WJ. (2002) Role of testosterone, estradiol, and insulin in diet- and exercise-induced reductions in serum-stimulated prostate cancer cell growth in vitro. *Nutr Cancer* 42:112–116.
 15. Mantzoros CS, Tzonou A, Signorello LB, Stampfer M, Trichopoulos D, Adami HO. (1997) Insulin-like growth factor 1 in relation to prostate cancer and benign prostatic hyperplasia. *Br J Cancer* 76:1115–1118.
 16. Wolk A, Mantzoros CS, Andersson SO, et al. (1998) Insulin-like growth factor 1 and prostate cancer risk: a population-based, case-control study. *J Natl Cancer Inst* 90:911–915.
 17. Chan JM, Stampfer MJ, Giovannucci E, et al. (1998) Plasma insulin-like growth factor-I and prostate cancer risk: a prospective study. *Science* 279:563–566.
 18. Stattin P, Bylund A, Rinaldi S, et al. (2000) Plasma insulin-like growth factor-I, insulin-like growth factor-binding proteins, and prostate cancer risk: a prospective study. *J Natl Cancer Inst* 92:1910–1917.
 19. Ngo TH, Barnard RJ, Leung PS, Cohen P, Aronson WJ. (2003) Insulin-like growth factor I (IGF-I) and IGF binding protein-1 modulate prostate cancer cell growth and apoptosis: possible mediators for the effects of diet and exercise on cancer cell survival. *Endocrinology* 144:2319–2324.
 20. Gurumurthy S, Vasudevan KM, Rangnekar VM. (2001) Regulation of apoptosis in prostate cancer. *Cancer Metastasis Rev* 20:225–243.
 21. Barnard RJ, Ngo TH, Leung PS, Aronson WJ, Golding LA. (2003) A low-fat diet and/or strenuous exercise alters the IGF axis in vivo and reduces prostate tumor cell growth in vitro. *Prostate* 56:201–206.
 22. Barnard RJ, Leung PS, Aronson WJ, Cohen P, Golding LA. (2007) A mechanism to explain how regular exercise might reduce the risk for clinical prostate cancer. *Eur J Cancer Prev* 16:415–421.
 23. Leung PS, Aronson WJ, Ngo TH, Golding LA, Barnard RJ. (2004) Exercise alters the IGF axis in vivo and increases p53 protein in prostate tumor cells in vitro. *J Appl Physiol* 96:450–454.
 24. Bond VJ, Sridhar R, Kassa A, Blakely R, Austin R, Millis R. (2009) Aerobic fitness levels in African-American men, serum IGF-I, and growth of prostate cancer cells. *Med Sci Sports Exerc* 40:S289.
 25. Heron-Milhavet L, LeRoith D. (2002) Insulin-like growth factor I induces MDM2-dependent degradation of p53 via the p38 MAPK pathway in response to DNA damage. *J Biol Chem* 277:15600–15606.
 26. Heron-Milhavet L, Karas M, Goldsmith CM, Baum BJ, LeRoith D. (2001) Insulin-like growth factor-I (IGF-I) receptor activation rescues UV-damaged cells through a p38 signaling pathway. Potential role of the IGF-I receptor in DNA repair. *J Biol Chem* 276:18185–18192.
 27. Yu H, Rohan T. (2000) Role of the insulin-like growth factor family in cancer development and progression. *J Natl Cancer Inst* 92:1472–1489.
 28. Lehman TA, Bennett WP, Metcalf RA, et al. (1991) P53 mutations, ras mutations, and p53-heat shock 70 protein complexes in human lung carcinoma cell lines. *Cancer Res* 51: 4090–4096.
 29. Zhang Y, Fujita N, Tsuruo T. (1999) Caspase-mediated cleavage of p21waf1/cip1 converts cancer cells from growth arrest to undergoing apoptosis. *Oncogene* 18:1131–1138.

30. Tsujimoto Y. (2003) Cell death regulation by the bcl-2 protein family in the mitochondria. *J Cell Physiol* 195:158–167.
31. Nelson WG, De Marzo AM, Isaacs WB. (2003) Prostate cancer. *N Engl J Med* 349:366–381.
32. Huang B, Zhao J, Unkeless JC, Feng ZH, Xiong H. (2008) TLR signaling by tumor and immune cells: a double-edged sword. *Oncogene* 27:218–224.
33. Ornish D, Weidner G, Fair WR, et al. (2005) Intensive lifestyle changes May affect the progression of prostate cancer. *J Urol* 174:1065–1069.
34. Dewell A, Weidner G, Sumner MD, et al. (2007) Relationship of dietary protein and soy isoflavones to serum IGF-1 and IGF binding proteins in the prostate cancer lifestyle trial. *Nutr Cancer* 58:35–42.
35. Ornish D, Magbanua MJ, Weidner G, et al. (2008) Changes in prostate gene expression in men undergoing an intensive nutrition and lifestyle intervention. *Proc Natl Acad Sci USA* 105:8369–8374.
36. Feldser D, Agani F, Iyer NV, Pak B, Ferreira G, Semenza GL. (1999) Reciprocal positive regulation of hypoxia-inducible factor 1alpha and insulin-like growth factor 2. *Cancer Res* 59:3915–3918.
37. Goodyear LJ, Kahn BB. (1998) Exercise, glucose transport, and insulin sensitivity. *Annu Rev Med* 49:235–261.
38. Samani AA, Yakar S, LeRoith D, Brodt P. (2007) The role of the IGF system in cancer growth and metastasis: Overview and recent insights. *Endocr Rev* 28:20–47.
39. Manes S, Llorente M, Lacalle RA, et al. (1999) The matrix metalloproteinase-9 regulates the insulin-like growth factor-triggered autocrine response in DU-145 carcinoma cells. *J Biol Chem* 274:6935–6945.
40. Roberts CK, Won D, Pruthi S, et al. (2006) Effect of a short-term diet and exercise intervention on oxidative stress, inflammation, MMP-9, and monocyte chemotactic activity in men with metabolic syndrome factors. *J Appl Physiol* 100:1657–1665.
41. Brodt P, Fallavollita L, Khatib AM, Samani AA, Zhang D. (2001) Cooperative regulation of the invasive and metastatic phenotypes by different domains of the type I insulin-like growth factor receptor beta subunit. *J Biol Chem* 276:33608–33615.
42. Wu Y, Yakar S, Zhao L, Hennighausen L, LeRoith D. (2002) Circulating insulin-like growth factor-I levels regulate colon cancer growth and metastasis. *Cancer Res* 62:1030–1035.
43. Kingsley LA, Fournier PG, Chirgwin JM, Guise TA. (2007) Molecular biology of bone metastasis. *Mol Cancer Ther* 6:2609–2617.
44. Hellawell GO, Turner GD, Davies DR, Poulson R, Brewster SF, Macaulay VM. (2002) Expression of the type 1 insulin-like growth factor receptor is up-regulated in primary prostate cancer and commonly persists in metastatic disease. *Cancer Res* 62:2942–2950.
45. Belfiore A. (2007) The role of insulin receptor isoforms and hybrid insulin/IGF-I receptors in human cancer. *Curr Pharm Des* 13:671–686.

1. Mathers CD, Loncar D. (2006) Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med* 3(11):e442.
2. Flood DM, Weiss NS, Cook LS, et al. (2000) Colorectal cancer incidence in Asian migrants to the united states and their descendants. *Cancer Causes Control* 11(5):403–411.
3. Boyle P, Langman JS. (2000) ABC of colorectal cancer – epidemiology. *Br Med J* 321(7264):805–808.
4. Slaterry ML. (2004) Physical activity and colorectal cancer. *Sports Med* 34(4): 239–252.
5. Slaterry ML, Levin TR, Ma K, et al. (2003) Family history and colorectal cancer: predictors of risk. *Cancer Causes Control* 14(9):879–887.
6. Harriss DJ, Atkinson G, Batterham A, et al. (2009) Lifestyle factors and colorectal cancer risk (2): a systematic review and meta-analysis of associations with leisure-time physical activity. *Colorectal Dis* 11(7):689–701.
7. Neugut AI, Jacobson JS, De Vivo I. (1993) Epidemiology of colorectal adenomatous polyps. *Cancer Epidemiol Biomarkers Prev* 2(2):159–176.
8. Harriss DJ, Cable NT, George K, et al. (2007) Physical activity before and after diagnosis of colorectal cancer: disease risk, clinical outcomes, response pathways and biomarkers. *Sports Med* 37(11):947–960.
9. Wallace K, Baron JA, Karagas MR, et al. (2005) The association of physical activity and body mass index with the risk of large bowel polyps. *Cancer Epidemiol Biomarkers Prev* 14(9):2082–2086.

10. Rosenberg L, Boggs D, Wise LA, et al. (2006) A follow-up study of physical activity and incidence of colorectal polyps in African-American women. *Cancer Epidemiol Biomarkers Prev* 15(8):1438–1442.
11. Larsen IK, Grotmol T, Almendingen K, et al. (2006) Lifestyle as a predictor for colonic neoplasia in asymptomatic individuals. *BMC Gastroenterol* 6:5.
12. Abrahamson PE, King IB, Ulrich CM, et al. (2007) No effect of exercise on colon mucosal prostaglandin concentrations: a 12-month randomized controlled trial. *Cancer Epidemiol Biomarkers Prev* 16(11):2351–2356.
13. McTiernan A, Yasui Y, Sorensen B, et al. (2006) Effect of a 12-month exercise intervention on patterns of cellular proliferation in colonic crypts: a randomized controlled trial. *Cancer Epidemiol Biomarkers Prev* 15(9):1588–1597.
14. Basterfield L, Reul J, Mathers JC. (2005) Impact of physical activity on intestinal cancer development in mice. *J Nutr* 135(12):3002S–3008S.
15. Banerjee AK, Mandal A, Chanda D, et al. (2003) Oxidant, antioxidant and physical exercise. *Mol Cell Biochem* 253(1–2):307–312.
16. Shephard RJ, Shek PN. (1999) Effects of exercise and training on natural killer cell counts and cytolytic activity – a meta-analysis. *Sports Med* 28(3):177–195.
17. Dreher D, Junod AF. (1996) Role of oxygen free radicals in cancer development. *Eur J Cancer* 32A(1):30–38.
18. Demarzo MMP, Garcia SB. (2004) Exhaustive physical exercise increases the number of colonic preneoplastic lesions in untrained rats treated with a chemical carcinogen. *Cancer Lett* 216(1):31–34.
19. Westerlind KC. (2003) Physical activity and cancer prevention-mechanisms. *Med Sci Sports Exerc* 35(11):1834–1840.
20. Haydon AMM, MacInnis RJ, English DR, et al. (2006) Effect of physical activity and body size on survival after diagnosis with colorectal cancer. *Gut* 55(1):62–67.
21. Nickelsen TN, Jorgensen T, Kronborg O. (2005) Lifestyle and 30-day complications to surgery for colorectal cancer. *Acta Oncol* 44(3):218–223.
22. Meyerhardt JA, Giovannucci EL, Holmes MD, et al. (2006) Physical activity and survival after colorectal cancer diagnosis. *J Clin Oncol* 24(22):3527–3534.
23. Irwin ML, Crumley D, McTiernan A, et al. (2003) Physical activity levels before and after a diagnosis of breast carcinoma – the health, eating, activity, and lifestyle (HEAL) study. *Cancer* 97(7):1746–1757.
24. Courneya KS, Friedenreich CM, Quinney HA, et al. (2003) A randomized trial of exercise and quality of life in colorectal cancer survivors. *Eur J Cancer Care* 12(4):347–357.
25. Courneya KS, Friedenreich CM, Quinney HA, et al. (2004) Predictors of adherence and contamination in a randomized trial of exercise in colorectal cancer survivors. *Psycho-Oncology* 13(12):857–866.
26. Courneya KS, Friedenreich CM, Quinney HA, et al. (2005) A longitudinal study of exercise barriers in colorectal cancer survivors participating in a randomized controlled trial. *Ann Behav Med* 29(2):147–153.
27. Allgayer H, Owen RW, Nair J, et al. (2008) Short-term moderate exercise programs reduce oxidative DNA damage as determined by high-performance liquid chromatography-electrospray ionization-mass spectrometry in patients with colorectal carcinoma following primary treatment. *Scandinavian J Gastroenterol* 43(8):971–978.
28. Meyerhardt JA, Heseltine D, Niedzwiecki D, et al. (2006) Impact of physical activity on cancer recurrence and survival in patients with stage III colon cancer: findings from CALGB 89803. *J Clin Oncol* 24(22):3535–3541.
29. Courneya KS, Booth CM, Gill S, et al. (2008) The colon health and life-long exercise change trial: a randomized trial of the national cancer institute of Canada clinical trials group. *Curr Oncol* 15(6):262–270.
30. Spence RR, Heesch KC, Eakin EG et al. (2007) Randomised controlled trial of a supervised exercise rehabilitation program for colorectal cancer survivors immediately after chemotherapy: study protocol. *BMC Cancer* 7:154.
31. de Lima C, Alves L, Iagher F, et al. (2008) Anaerobic exercise reduces tumor growth, cancer cachexia and increases macrophage and lymphocyte response in walker 256 tumor-bearing rats. *Eur J Appl Physiol* 104(6):957–964.
32. Bacurau AVN, Belmonte MA, Navarro F, et al. (2007) Effect of a high-intensity exercise training on the metabolism and function of macrophages and lymphocytes of walker 256 tumor-bearing rats. *Exp Biol Med* 232:1289–1299.
33. Courneya KS, Friedenreich CM. (2001) Framework PEACE: an organizational model for examining physical exercise across the cancer experience. *Ann Behav Med* 23(4): 263–272.
34. Courneya KS, Friedenreich CM. (1997) Relationship between exercise during treatment and current quality of life among survivors of breast cancer. *J Psychosoc Oncol* 15(3–4): 35–57.
35. Schmitz KH, Holtzman J, Courneya KS, et al. (2005) Controlled physical activity trials in cancer survivors: a systematic review and meta-analysis. *Cancer Epidemiol Biomarkers Prev* 14(7):1588–1595.

36. Blair SN, Brodney S. (1999) Effects of physical inactivity and obesity on morbidity and mortality: current evidence and research issues. *Med Sci Sports Exerc* 31(11):S646–S662.
37. Coebergh JWW, Janssen-Heijnen MLG, Post PN, et al. (1999) Serious co-morbidity among unselected cancer patients newly diagnosed in the southeastern part of the Netherlands in 1993–1996. *J Clin Epidemiol* 52(12):1131–1136.
38. Baade PD, Fritschi L, Eakin EG. (2006) Non-cancer mortality among people diagnosed with cancer (Australia). *Cancer Causes Control* 17(3):287–297.
39. Thune I, Lund E. (1996) Physical activity and risk of colorectal cancer in men and women. *Br J Cancer* 73(9):1134–1140.
40. Peters HPF, De Vries WR, Vanberge-Henegouwen GP, et al. (2001) Potential benefits and hazards of physical activity and exercise on the gastrointestinal tract. *Gut* 48(3):435–439.
41. Holdstoc DJ, Misiewicz JJ. (1970) Factors controlling colonic motility – colonic pressures and transit after meals in patients with total gastrectomy, pernicious anaemia or duodenal ulcer. *Gut* 11(2):100–110.
42. De Schryver AM, Keulemans YC, Peters HP, et al. (2005) Effects of regular physical activity on defecation pattern in middle-aged patients complaining of chronic constipation. *Scandinavian J Gastroenterol* 40(4):422–429.
43. Cordain L, Latin RW, Behnke JJ. (1986) The effects of an aerobic running program on bowel transit-time. *J Sports Med Phys Fitness* 26(1):101–104.
44. Oettle GJ. (1991) Effect of moderate exercise on bowel habit. *Gut* 32(8):941–944.
45. Kofler KH, Menkes A, Redmond RA, et al. (1992) Strength training accelerates gastrointestinal transit in middle-aged and older men. *Med Sci Sports Exerc* 24(4): 415–419.
46. Liu F, Kondo T, Toda Y. (1993) Brief physical inactivity prolongs colonic transit-time in elderly active men. *Int J Sports Med* 14(8):465–467.
47. Bingham SA, Cummings JH. (1989) Effect of exercise and physical-fitness on large intestinal function. *Gastroenterology* 97(6):1389–1399.
48. Coenen C, Wegener M, Wedmann B, et al. (1992) Does physical exercise influence bowel transit-time in healthy young men. *Am J Gastroenterol* 87(3):292–295.
49. Robertson G, Meshkinpour H, Vandenberg K, et al. (1993) Effects of exercise on total and segmental colon transit. *J Clin Gastroenterol* 16(4):300–303.
50. Schatzkin A, Lanza E, Corle D, et al. (2000) Lack of effect of a low-fat, high-fiber diet on the recurrence of colorectal adenomas. *N Engl J Med* 342(16):1149–1155.
51. Asano T, McLeod RS. (2002) Dietary fibre for the prevention of colorectal adenomas and carcinomas. *Cochrane Database of Syst Rev* 2:CD003430.
52. Kuper H, Adami HO, Trichopoulos D. (2000) Infections as a major preventable cause of human cancer. *J Intern Med* 248(3):171–183.
53. Coussens LM, Werb Z. (2002) Inflammation and cancer. *Nature* 420(6917):860–867.
54. Demarzo MMP, Martins LV, Fernandes CR, et al. (2008) Exercise reduces inflammation and cell proliferation in rat colon carcinogenesis. *Med Sci Sports Exerc* 40(4):618–621.
55. Baniyash M. (2006) Chronic inflammation, immunosuppression and cancer: new insights and outlook. *Semin Cancer Biol* 16(1):80–88.
56. Boland CR, Luciani MG, Gasche C, et al. (2005) Infection, inflammation, and gastrointestinal cancer. *Gut* 54(9):1321–1331.
57. Johne B, Fagerhol MK, Lyberg T, et al. (1997) Functional and clinical aspects of the myelomonocyte protein calprotectin. *J Clin Pathol Mol Pathol* 50(3):113–123.
58. Kronborg O, Ugstad M, Fuglerud P, et al. (2000) Faecal calprotectin levels in a high risk population for colorectal neoplasia. *Gut* 46(6):795–800.
59. Poullis A, Foster R, Shetty A, et al. (2004) Bowel inflammation as measured by fecal calprotectin: a link between lifestyle factors and colorectal cancer risk. *Cancer Epidemiol Biomarkers Prev* 13(2):279–284.
60. Hauret KG, Bostick RM, Matthews CE, et al. (2004) Physical activity and reduced risk of incident sporadic colorectal adenomas: observational support for mechanisms involving energy balance and inflammation modulation. *Am J Epidemiol* 159(10): 983–992.
61. Eisinger AL, Prescott SM, Jones DA, et al. (2007) The role of cyclooxygenase-2 and prostaglandins in colon cancer. *Prostaglandins Other Lipid Mediat* 82(1–4):147–154.

62. Martinez ME, Heddens D, Earnest DL, et al. (1999) Physical activity, body mass index, and prostaglandin E-2 levels in rectal mucosa. *J Natl Cancer Inst* 91(11):950–953.
63. Corpet DE, Pierre F. (2003) Point: from animal models to prevention of colon cancer. Systematic review of chemoprevention in Min mice and choice of the model system. *Cancer Epidemiol Biomarkers Prev* 12(5):391–400.
64. Pischon T, Hankinson SE, Hotamisligil GS, et al. (2003) Leisure-time physical activity and reduced plasma levels of obesity-related inflammatory markers. *Obesity Res* 11(9):1055–1064.
65. Jankord R, Jemiolo B. (2004) Influence of physical activity on serum IL-6 and IL-10 levels in healthy older men. *Med Sci Sports Exerc* 36(6):960–964.
66. Stewart LK, Flynn MG, Campbell WW, et al. (2005) Influence of exercise training and age on CD14⁺ cell-surface expression of toll-like receptor 2 and 4. *Brain Behav Immun* 19(5):389–397.
67. Smith JK, Dykes R, Douglas JE, et al. (1999) Long-term exercise and atherogenic activity of blood mononuclear cells in persons at risk of developing ischemic heart disease. *JAMA* 281(18):1722–1727.
68. You TJ, Berman DM, Ryan AS, et al. (2004) Effects of hypocaloric diet and exercise training on inflammation and adipocyte lipolysis in obese postmenopausal women. *J Clin Endocrinol Metab* 89(4):1739–1746.
69. Triantafyllou M, Triantafyllou K. (2002) Lipopolysaccharide recognition: CD14, TLRs and the LPS-activation cluster. *Trends Immunol* 23(6):301–304.
70. Flynn MG, McFarlin BK, Phillips MD, et al. (2003) Toll-like receptor 4 and CD14 mRNA expression are lower in resistance exercise-trained elderly women. *J Appl Physiol* 95(5): 1833–1842.
71. McFarlin BK, Flynn MG, Campbell WW, et al. (2004) TLR4 is lower in resistance-trained older women and related to inflammatory cytokines. *Med Sci Sports Exerc* 36(11): 1876–1883.
72. Armstrong F, Mathers JC. (2000) Kill and cure: dietary augmentation of immune defences against colon cancer. *Proc Nutr Soc* 59(2):215–220.
73. Shevde LA, Joshi NN, Dudhat SB, et al. (1999) Immune functions, clinical parameters and hormone receptor status in breast cancer patients. *J Cancer Res Clin Oncol* 125(5): 313–320.
74. Liljefors M, Nilsson B, Skog ALH, et al. (2003) Natural killer (NK) cell function is a strong prognostic factor in colorectal carcinoma patients treated with the monoclonal antibody 17-1a. *Int J Cancer* 105(5):717–723.
75. Newsholme E, Parry-Billings M (1994) Effects of exercise on the immune system. In: Bouchard C, Shephard R, Stephens T (eds) *Physical Activity, Fitness, and Health: International Proceedings Consensus Statement*. Human Kinetics, Champaign (IL), p. 451.
76. Dishman RK, Hong S, Soares J, et al. (2000) Activity-wheel running blunts suppression of splenic natural killer cell cytotoxicity after sympathectomy and footshock. *Physiol Behav* 71(3–4):297–304.
77. Dishman RK, Warren JM, Hong S, et al. (2000) Treadmill exercise training blunts suppression of splenic natural killer cell cytotoxicity after footshock. *J Appl Physiol* 88(6): 2176–2182.
78. Fairey AS, Courneya KS, Field CJ, et al. (2002) Physical exercise and immune system function in cancer survivors – a comprehensive review and future directions. *Cancer* 94(2):539–551.
79. Nieman DC, Cook VD, Henson DA, et al. (1995) Moderate exercise training and natural-killer-cell cytotoxic activity in breast-cancer patients. *Int J Sports Med* 16(5): 334–337.
80. Hayes SC, Rowbottom D, Davies PSW, et al. (2003) Immunological changes after cancer treatment and participation in an exercise program. *Med Sci Sports Exerc* 35(1):2–9.
81. Allgayer H, Nicolaus S, Schreiber S. (2004) Decreased interleukin-1 receptor antagonist response following moderate exercise in patients with colorectal carcinoma after primary treatment. *Cancer Detect Prev* 28(3):208–213.
82. Nieman DC. (1998) Exercise and resistance to infection. *Can J Physiol Pharmacol* 76(5):573–580.
83. Fehrenbach E, Northoff H. (2001) Free radicals, exercise, apoptosis, and heat shock proteins. *Exerc Immunol Rev* 7:66–89.
84. Hu FB, Manson JE, Liu SM, et al. (1999) Prospective study of adult onset diabetes mellitus (type 2) and risk of colorectal cancer in women. *J Natl Cancer Inst* 91(6):542–547.
85. La Vecchia C, Negri E, Decarli A, et al. (1997) Diabetes mellitus and colorectal cancer risk. *Cancer Epidemiol Biomarkers Prev* 6(12):1007–1010.
86. Keku TO, Lund PK, Galanko J, et al. (2005) Insulin resistance, apoptosis, and colorectal adenoma risk. *Cancer Epidemiol Biomarkers Prev* 14(9):2076–2081.
87. Schoen RE, Weissfeld JL, Kuller LH, et al. (2005) Insulin-like growth factor-I and insulin are associated with the presence and advancement of adenomatous polyps. *Gastroenterology* 129(2):464–475.

88. Kaaks R, Toniolo P, Akhmedkhanov A, et al. (2000) Serum C-peptide, insulin-like growth factor (IGF)-I, IGF-binding proteins, and colorectal cancer risk in women. *J Natl Cancer Inst* 92(19):1592–1600.
89. Ma J, Giovannucci E, Pollak M, et al. (2004) A prospective study of plasma C-peptide and colorectal cancer risk in men. *J Natl Cancer Inst* 96(7):546–553.
90. Yam D, Fink A, Mashiah A, et al. (1996) Hyperinsulinemia in colon, stomach and breast cancer patients. *Cancer Lett* 104(2):129–132.
91. LaMonte MJ, Blair SN, Church TS. (2005) Physical activity and diabetes prevention. *J Appl Physiol* 99(3):1205–1213.
92. Platz EA, Hankinson SE, Rifai N, et al. (1999) Glycosylated hemoglobin and risk of colorectal cancer and adenoma (United States). *Cancer Causes Control* 10(5):379–386.
93. Irwin ML, McTiernan A, Bernstein L, et al. (2005) Relationship of obesity and physical activity with C-peptide, leptin, and insulin-like growth factors in breast cancer survivors. *Cancer Epidemiol Biomarkers Prev* 14(12):2881–2888.
94. Schmitz KH, Ahmed RL, Hannan PJ, et al. (2005) Safety and efficacy of weight training in recent breast cancer survivors to alter body composition, insulin, and insulin-like growth factor axis proteins. *Cancer Epidemiol Biomarkers Prev* 14(7):1672–1680.
95. Fairey AS, Courneya KS, Field CJ, et al. (2003) Effects of exercise training on fasting insulin, insulin resistance, insulin-like growth factors, and insulin-like growth factor binding proteins in postmenopausal breast cancer survivors: a randomized controlled trial. *Cancer Epidemiol Biomarkers Prev* 12(8):721–727.
96. Davies M, Gupta S, Goldspink G, et al. (2006) The insulin-like growth factor system and colorectal cancer: clinical and experimental evidence. *Int J Colorectal Dis* 21(3):201–208.
97. Ma J, Pollak MN, Giovannucci E, et al. (1999) Prospective study of colorectal cancer risk in men and plasma levels of insulin-like growth factor (IGF)-I and IGF-binding protein-3. *J Natl Cancer Inst* 91(7):620–625.
98. Renehan AG, Painter JE, Atkin WS, et al. (2001) High-risk colorectal adenomas and serum insulin-like growth factors. *Br J Surg* 88(1):107–113.
99. Renehan AG, Zwahlen M, Minder C, et al. (2004) Insulin-like growth factor (IGF)-I, IGF binding protein-3, and cancer risk: systematic review and meta-regression analysis. *Lancet* 363(9418):1346–1353.
100. Goodman GD, Barrett CE. (1997) Epidemiology of insulin-like growth factor-I in elderly men and women - the Rancho Bernardo study. *Am J Epidemiol* 145(11):970–976.
101. Rudman D, Mattson DE. (1994) Serum insulin-like growth factor-I in healthy older men in relation to physical activity. *J Am Geriatrics Soc* 42(1):71–76.
102. Chang S, Wu XF, Yu H, et al. (2002) Plasma concentrations of insulin-like growth factors among healthy adult men and postmenopausal women: associations with body composition, lifestyle, and reproductive factors. *Cancer Epidemiol Biomarkers Prev* 11(8):758–766.
103. Morimoto LM, Newcomb PA, White E, et al. (2005) Variation in plasma insulin-like growth factor-I and insulin-like growth factor binding protein-3: personal and lifestyle factors (United States). *Cancer Causes Control* 16(8):917–927.
104. Landin-Wilhelmsen K, Wilhelmsen L, Lappas G, et al. (1994) Serum insulin-like growth factor I in a random population sample of men and women: relation to age, sex, smoking habits, coffee consumption and physical activity, blood pressure and concentrations of plasma lipids, fibrinogen, parathyroid hormone and osteocalcin. *Clin Endocrinol* 41(3):351–357.
105. Haydon AMM, MacInnis RJ, English DR, et al. (2006) Physical activity, insulin-like growth factor 1, insulin-like growth factor binding protein 3, and survival from colorectal cancer. *Gut* 55(5):689–694.
106. Manetta J, Brun JF, Fedou C, et al. (2003) Serum levels of insulin-like growth factor-I (IGFI), and IGF-binding proteins-1 and-3 in middle-aged and young athletes versus sedentary men: relationship with glucose disposal. *Metab Clin Exp* 52(7):821–826.
107. Rosendal L, Langberg H, Flyvbjerg A, et al. (2002) Physical capacity influences the response of insulin-like growth factor and its binding proteins to training. *J Appl Physiol* 93(5): 1669–1675.
108. Poehlman ET, Rosen CJ, Copeland KC. (1994) The influence of endurance training on insulin-like growth-factor-I in older individuals. *Metab Clin Exp* 43(11):1401–1405.
109. Filaire E, Jouanel P, Colombier M, et al. (2003) Effects of 16 weeks of training prior to a major competition on hormonal and biochemical parameters in young elite gymnasts. *J Pediatr Endocrinol Metab* 16(5):741–750.

110. Gomez-Merino D, Chennaoui M, Drogou C, et al. (2004) Influence of energy deficiency on the insulin-like growth factor I axis in a military training program. *Horm Metab Res* 36(7):506–511.
111. Nemet D, Connolly PH, Pontello-Pescatello AM, et al. (2004) Negative energy balance plays a major role in the IGF-I response to exercise training. *J Appl Physiol* 96(1):276–282.
112. Cho KR, Vogelstein B. (1992) Suppressor gene alterations in the colorectal adenomacarcinoma sequence. *J Cell Biochem* 16G:137–141.
113. Bos JL, Fearon ER, Hamilton SR, et al. (1987) Prevalence of ras gene-mutations in human colorectal cancers. *Nature* 327(6120):293–297.
114. Rashid A, Zahurak M, Goodman SN, et al. (1999) Genetic epidemiology of mutated Kras proto-oncogene, altered suppressor genes, and microsatellite instability in colorectal adenomas. *Gut* 44(6):826–833.
115. Tortola S, Marcuello E, Gonzalez I, et al. (1999) P53 and K-ras gene mutations correlate with tumor aggressiveness but are not of routine prognostic value in colorectal cancer. *J Clin Oncol* 17(5):1375–1381.
116. Leung PS, Aronson WJ, Ngo TH, et al. (2004) Exercise alters the IGF axis in vivo and increases p53 protein in prostate tumor cells in vitro. *J Appl Physiol* 96(2):450–454.
117. Slattery ML, Anderson K, Curtin K, et al. (2001) Lifestyle factors and ki-ras mutations in colon cancer tumors. *Mut Res Fund Mol Mech Mutagen* 483(1–2):73–81.
118. Heavey PM, McKenna D, Rowland IR. (2004) Colorectal cancer and the relationship between genes and the environment. *Nutr Cancer International Journal* 48(2):124–141.
119. Buchmeyer K, Doering F, Daniel H, et al. (2008) Alteration of gene expression in rat colon mucosa after exercise. *Ann Anat-Anat Anz* 190(1):71–80.
120. Harriss DJ, Atkinson G, George K, et al. (2009) Lifestyle factors and colorectal cancer risk (1): systematic review and meta-analysis of associations with body mass index. *Colorectal Dis* 11(6):547–563.
121. MacInnis RJ, English DR, Hopper JL, et al. (2004) Body size and composition and colon cancer risk in men. *Cancer Epidemiol Biomarkers Prev* 13(4):553–559.
122. MacInnis RJ, English DR, Hopper JL, et al. (2006) Body size and composition and colon cancer risk in women. *Int J Cancer* 118(6):1496–1500.
123. Slattery ML, Ballard-Barbash R, Edwards S, et al. (2003) Body mass index and colon cancer: an evaluation of the modifying effects of estrogen (United States). *Cancer Causes Control* 14(1):75–84.
124. Mao Y, Pan SY, Wen SW, et al. (2003) Physical inactivity, energy intake, obesity and the risk of rectal cancer in Canada. *Int J Cancer* 105(6):831–837.
125. Grundy SM, Blackburn G, Higgins M, et al. (1999) Physical activity in the prevention and treatment of obesity and its comorbidities. *Med Sci Sports Exerc* 31(11):S502–S508.
126. Bensimhon DR, Kraus WE, Donahue MP. (2006) Obesity and physical activity: a review. *Am Heart J* 151(3):598–603.
127. Bouchard C, Tremblay A, Nadeau A, et al. (1990) Long-term exercise training with constant energy-intake. I. Effect on body-composition and selected metabolic variables. *Int J Obesity* 14(1):57–73.
128. Thompson HJ, Zhu ZJ, Jiang WQ. (2004) Weight control and breast cancer prevention: are the effects of reduced energy intake equivalent to those of increased energy expenditure? *J Nutr* 134(12):3407S–3411S.
129. Parkin DM, Bray F, Ferlay J, et al. (2005) Global cancer statistics, 2002. *CA Cancer J Clin* 55(2):74–108.

1. Bosetti C, Levi F, Lucchini F, Negri E, La VC. (2005) Lung cancer mortality in European women: recent trends and perspectives. *Ann Oncol* 16:1597–1604.
2. Samet JM. (1989) Radon and lung cancer. *J Natl Cancer Inst* 81(10):745–757.
3. American Cancer Society. *Cancer Facts and Figures*. 2006. Atlanta, GA, USA.
4. Jemal A, Thun MJ, Ries LA, et al. (2008) Annual report to the nation on the status of cancer, 1975–2005, featuring trends in lung cancer, tobacco use, and tobacco control. *J Natl Cancer Inst* 100:1672–1694.
5. Govindan R, Page N, Morgensztern D, et al. (2006) Changing epidemiology of small-cell lung cancer in the United States over the last 30 years: analysis of the surveillance, epidemiologic, and end Results database. *J Clin Oncol*

24:4539–4544.

6. Brambilla E, Travis WD, Colby TV, Corrin B, Shimosato Y. (2001) The new world health organization classification of lung tumours. *Eur Respir J* 18:1059–1068.
7. Yoshimi I, Ohshima A, Ajiki W, Tsukuma H, Sobue T. (2003) A comparison of trends in the incidence rate of lung cancer by histological type in the Osaka cancer registry, Japan and in the surveillance, epidemiology and end Results program, USA. *Jpn J Clin Oncol* 33:98–104.
8. Spiro SG, Gould MK, Colice GL. (2007) Initial evaluation of the patient with lung cancer: symptoms, signs, laboratory tests, and paraneoplastic syndromes: ACCP evidenced-based clinical practice guidelines (2nd edition). *Chest* 132:149S–160S.
9. (1997) Pretreatment evaluation of non-small-cell lung cancer. The American Thoracic Society and the European Respiratory Society. *Am J Respir Crit Care Med* 156:320–332.
10. Mountain CF. (1997) Revisions in the international system for staging lung cancer. *Chest* 111:1710–1717.
11. Boyle P, Ferlay J. (2005) Mortality and survival in breast and colorectal cancer. *Nat Clin Pract Oncol* 2:424–425.
12. Firat S, Bousamra M, Gore E, Byhardt RW. (2002) Comorbidity and KPS are independent prognostic factors in stage I non-small-cell lung cancer. *Int J Radiat Oncol Biol Phys* 52: 1047–1057.
13. Scott WJ, Howington J, Feigenberg S, Movsas B, Pisters K. (2007) Treatment of non-small cell lung cancer stage I and stage II: ACCP evidence-based clinical practice guidelines (2nd edition). *Chest* 132:234S–242S.
14. Ginsberg RJ, Rubinstein LV. (1995) Randomized trial of lobectomy versus limited resection for T1 N0 non-small cell lung cancer. Lung cancer study group. *Ann Thorac Surg* 60:615–622.
15. Daniels LJ, Balderson SS, Onaitis MW, D'Amico TA. (2002) Thoracoscopic lobectomy: a safe and effective strategy for patients with stage I lung cancer. *Ann Thorac Surg* 74:860–864.
16. Strauss GM, Herndon JE, Maddaus MA, Johnstone DW, Johnson EA, Watson DM, Sugarbaker DJ, Schilsky RA, Vokes EE, Green MR. (2006) Adjuvant chemotherapy in stage IB non-small cell lung cancer (NSCLC): Update of Cancer and Leukemia Group B (CALGB) protocol 9633. ASCO Annual Meeting. *J Clin Oncol*, ASCO Meeting Proceedings Part I 24:18S (June 20 Supplement).
17. Auperin A, Le PC, Pignon JP, et al. (2006) Concomitant radio-chemotherapy based on platin compounds in patients with locally advanced non-small cell lung cancer (NSCLC): a metaanalysis of individual data from 1764 patients. *Ann Oncol* 17:473–483.
18. Detterbeck F, Rivera M. (2001) Clinical presentation and diagnosis. In: Detterbeck, FC, Rivera, MP, Socinski, MA, Rosenman, JG (eds) *Diagnosis and Treatment of Lung Cancer: An Evidence-Based Guide for the Practicing Clinician*. W.B. Saunders Co., Philadelphia, pp. 45–72.
19. Socinski MA, Crowell R, Hensing TE. (2007) Treatment of non-small cell lung cancer, stage IV: ACCP evidence-based clinical practice guidelines (2nd edition). *Chest* 132: 277S–289S.
20. Makitaro R, Paakko P, Huhti E, Bloigu R, Kinnula VL. (2002) Prospective population-based study on the survival of patients with lung cancer. *Eur Respir J* 19:1087–1092.
21. Socinski MA, Morris DE, Masters GA, Lilenbaum R. (2003) Chemotherapeutic management of stage IV non-small cell lung cancer. *Chest* 123:226S–243S.
22. Socinski MA, Baggstrom MQ, Hensing TA. (2003) Duration of therapy in advanced, metastatic non-small-cell lung cancer. *Clin Adv Hematol Oncol* 1:33–38.
23. De RD, Pijls-Johannesma M, Vansteenkiste J, Kester A, Rutten I, Lambin P. (2006) Systematic review and meta-analysis of randomised, controlled trials of the timing of chest radiotherapy in patients with limited-stage, small-cell lung cancer. *Ann Oncol* 17:543–552.
24. Arriagada R, Le Chevalier T, Borie F. (1994) Randomized trial of prophylactic cranial irradiation (PCI) for patients with small cell lung cancer (SCLC) in complete remission. ASCO Annual Conference, 13.
25. The Prophylactic Cranial Irradiation Overview Collaborative Group (2000) Cranial irradiation for preventing brain metastases of small cell lung cancer in patients in complete remission. *Cochrane Database Systematic Review* CD002805.
26. Nici L, Donner C, Wouters E, et al. (2006) American thoracic society/european respiratory society statement on pulmonary rehabilitation. *Am J Respir Crit Care Med* 173:1390–1413.
27. Lacasse Y, Goldstein R, Lasserson TJ, Martin S. (2006) Pulmonary rehabilitation for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev* CD003793
28. Jaeschke R, Singer J, Guyatt GH. (1989) Measurement of health status. Ascertaining the minimal clinically

- important difference. *Control Clin Trials* 10:407–415.
29. Spruit MA, Vanderhoven-Augustin I, Janssen PP, Wouters EF. (2008) Integration of pulmonary rehabilitation in COPD. *Lancet* 371:12–13.
 30. Holland AE, Hill CJ, Conron M, Munro P, McDonald CF. (2008) Short term improvement in exercise capacity and symptoms following exercise training in interstitial lung disease. *Thorax* 63:549–554.
 31. Spruit MA, Wouters EF. (2007) New modalities of pulmonary rehabilitation in patients with chronic obstructive pulmonary disease. *Sports Med* 37:501–518.
 32. Jones LW, Eves ND, Mackey JR, et al. (2007) Safety and feasibility of cardiopulmonary exercise testing in patients with advanced cancer. *Lung Cancer* 55:225–232.
 33. Wilcock A, Maddocks M, Lewis M, England R, Manderson C. (2008) Symptoms limiting activity in cancer patients with breathlessness on exertion: ask about muscle fatigue. *Thorax* 63:91–92.
 34. Weinstein H, Bates AT, Spaltro BE, Thaler HT, Steingart RM. (2007) Influence of preoperative exercise capacity on length of stay after thoracic cancer surgery. *Ann Thorac Surg* 84: 197–202.
 35. Epstein SK, Faling LJ, Daly BD, Celli BR. (1995) Inability to perform bicycle ergometry predicts increased morbidity and mortality after lung resection. *Chest* 107:311–316.
 36. Morice RC, Peters EJ, Ryan MB, Putnam JB, Ali MK, Roth JA. (1992) Exercise testing in the evaluation of patients at high risk for complications from lung resection. *Chest* 101:356–361.
 37. Wilcock A, Maddocks M, Lewis M, et al. (2008) Use of a cybex NORM dynamometer to assess muscle function in patients with thoracic cancer. *BMC Palliat Care* 7:3.
 38. Nezu K, Kushibe K, Tojo T, Takahama M, Kitamura S. (1998) Recovery and limitation of exercise capacity after lung resection for lung cancer. *Chest* 113:1511–1516.
 39. Bobbio A, Chetta A, Ampollini L, et al. (2008) Preoperative pulmonary rehabilitation in patients undergoing lung resection for non-small cell lung cancer. *Eur J Cardiothorac Surg* 33:95–98.
 40. Nagamatsu Y, Maeshiro K, Kimura NY, et al. (2007) Long-term recovery of exercise capacity and pulmonary function after lobectomy. *J Thorac Cardiovasc Surg* 134:1273–1278.
 41. Nezu K, Iioka S, Kushibe K, et al. (1994) [The relationship between postoperative changes of exercise capacity and pulmonary blood flow in the residual lung after lobectomy of the lung]. *Nippon Kyobu Geka Gakkai Zasshi* 42:340–345.
 42. Bobbio A, Chetta A, Carbognani P, et al. (2005) Changes in pulmonary function test and cardio-pulmonary exercise capacity in COPD patients after lobar pulmonary resection. *Eur J Cardiothorac Surg* 28:754–758.
 43. Brunelli A, Soggi L, Refai M, Salati M, Xiume F, Sabbatini A. (2007) Quality of life before and after major lung resection for lung cancer: a prospective follow-up analysis. *Ann Thorac Surg* 84:410–416.
 44. Handy JR Jr., Asaph JW, Skokan L, et al. (2002) What happens to patients undergoing lung cancer surgery? Outcomes and quality of life before and after surgery. *Chest* 122:21–30.
 45. Schulte T, Schniewind B, Dohrmann P, Kuchler T, Kurdow R. (2008) The extent of lung parenchyma resection significantly impacts long-term quality of life in patients with non small cell lung cancer. *Chest* 135(2):322–9.
 46. Man WD, Polkey MI, Donaldson N, Gray BJ, Moxham J. (2004) Community pulmonary rehabilitation after hospitalisation for acute exacerbations of chronic obstructive pulmonary disease: randomised controlled study. *BMJ* 329:1209.
 47. Jones LW, Peddle CJ, Eves ND, et al. (2007) Effects of presurgical exercise training on cardiorespiratory fitness among patients undergoing thoracic surgery for malignant lung lesions. *Cancer* 110:590–598.
 48. Spruit MA, Janssen PP, Willemssen SC, Hochstenbag MM, Wouters EF. (2006) Exercise capacity before and after an 8-week multidisciplinary inpatient rehabilitation program in lung cancer patients: a pilot study. *Lung Cancer* 52:257–260.
 49. Mejia R, Ward J, Lentine T, Mahler DA. (1999) Target dyspnea ratings predict expected oxygen consumption as well as target heart rate values. *Am J Respir Crit Care Med* 159:1485–1489.
 50. Horowitz MB, Littenberg B, Mahler DA. (1996) Dyspnea ratings for prescribing exercise intensity in patients with COPD. *Chest* 109:1169–1175.
 51. Ries AL. (2005) Minimally clinically important difference for the UCSD shortness of breath questionnaire, borg scale, and visual analog scale. *COPD* 2(1):105–110.
 52. Jones LW, Eves ND, Peterson BL, et al. (2008) Safety and feasibility of aerobic training on cardiopulmonary function and quality of life in postsurgical nonsmall cell lung cancer patients: a pilot study. *Cancer* 113:3430–3439.

53. Cesario A, Ferri L, Galetta D, et al. (2007) Post-operative respiratory rehabilitation after lung resection for non-small cell lung cancer. *Lung Cancer* 57:175–180.
 54. Gosker HR, Lencer NH, Franssen FM, van der Vusse GJ, Wouters EF, Schols AM. (2003) Striking similarities in systemic factors contributing to decreased exercise capacity in patients with severe chronic heart failure or COPD. *Chest* 123:1416–1424.
 55. Gosselink R, Troosters T, Decramer M. (1996) Peripheral muscle weakness contributes to exercise limitation in COPD. *Am J Respir Crit Care Med* 153:976–980.
 56. Nicoletti I, Cicoira M, Zanolla L, et al. (2003) Skeletal muscle abnormalities in chronic heart failure patients: relation to exercise capacity and therapeutic implications. *Congest Heart Fail* 9:148–154.
 57. Spruit MA, Troosters T, Trappenburg JC, Decramer M, Gosselink R. (2004) Exercise training during rehabilitation of patients with COPD: a current perspective. *Patient Educ Couns* 52:243–248.
 58. Spruit MA, Wouters EFM, Gosselink R. (2005) Rehabilitation programmes in sarcoidosis: a multidisciplinary approach. In: Drent, M, Costabel, U (eds.) *European Respiratory Monograph: Sarcoidosis*. European Respiratory Society Ltd., Wakefield, UK, pp. 316–326.
 59. Probst VS, Troosters T, Pitta F, Decramer M, Gosselink R. (2006) Cardiopulmonary stress during exercise training in patients with COPD. *Eur Respir J* 27:1110–1118.
 60. Sillen MJ, Janssen PP, Akkermans MA, Wouters EF, Spruit MA. (2008) The metabolic response during resistance training and neuromuscular electrical stimulation (NMES) in patients with COPD, a pilot study. *Respir Med* 102:786–789.
 61. Vanderthommen M, Duchateau J. (2007) Electrical stimulation as a modality to improve performance of the neuromuscular system. *Exerc Sport Sci Rev* 35:180–185.
 62. Sillen MJ, et al. (2009) Effects of neuromuscular electrical stimulation of muscles of ambulation in patients with CHF or COPD: a systematic review of the English-language literature. *Chest* (in press). 136(1):44–61.
 63. Crevenna R, Marosi C, Schmidinger M, Fialka-Moser V. (2006) Neuromuscular electrical stimulation for a patient with metastatic lung cancer – a case report. *Support Care Cancer* 14:970–973.
 64. Slotman BJ, Mauer ME, Bottomley A, et al. (2009) Prophylactic cranial irradiation in extensive disease small-cell lung cancer: short-term health-related quality of life and patient reported symptoms: Results of an international phase III randomized controlled trial by the EORTC radiation oncology and lung cancer groups. *J Clin Oncol* 27:78–84.
 65. Maddocks M, Mockett S, Wilcock A. (2007) Neuromuscular electrical stimulation (NMES): a reactive palliative therapy, a proactive supportive therapy or both? *Support Care Cancer* 15:111.
 66. Dimeo FC, Thomas F, Raabe-Menssen C, Propper F, Mathias M. (2004) Effect of aerobic exercise and relaxation training on fatigue and physical performance of cancer patients after surgery. A randomised controlled trial. *Support Care Cancer* 12:774–779.
 67. Schneider CM, Hsieh CC, Sprod LK, Carter SD, Hayward R. (2007) Exercise training manages cardiopulmonary function and fatigue during and following cancer treatment in male cancer survivors. *Integr Cancer Ther* 6:235–241.
 68. Dy SM, Lorenz KA, Naeim A, Sanati H, Walling A, Asch SM. (2008, Aug 10) Evidencebased recommendations for cancer fatigue, anorexia, depression, and dyspnea. *J Clin Oncol*. 26(23):3886–95.
-
1. Parkin DM, Bray F, Ferlay J, Pisani P. (2005) Global cancer statistics, 2002. *CA Cancer J Clin* 55:74–108.
 2. Coleman MP, Quaresma M, Berrino F, et al. (2008) Cancer survival in five continents: a worldwide population-based study (CONCORD). *Lancet Oncol* 9:730–756.
 3. Jones LW, Demark-Wahnefried W. (2006) Diet, exercise, and complementary therapies after primary treatment for cancer. *Lancet Oncol* 7:1017–1026.
 4. Blair SN, Kohl HW, III, Paffenbarger RS, Jr., Clark DG, Cooper KH, Gibbons LW. (1989) Physical fitness and all-cause mortality. A prospective study of healthy men and women. *JAMA* 262:2395–2401.
 5. Kampert JB, Blair SN, Barlow CE, Kohl HW, III. (1996) Physical activity, physical fitness, and all-cause and cancer mortality: a prospective study of men and women. *Ann Epidemiol* 6:452–457.
 6. Evenson KR, Stevens J, Cai J, Thomas R, Thomas O. (2003) The effect of cardiorespiratory fitness and obesity on cancer mortality in women and men. *Med Sci Sports Exerc* 35:270–277.

7. Laukkanen JA, Lakka TA, Rauramaa R, et al. (2001) Cardiovascular fitness as a predictor of mortality in men. *Arch Intern Med* 161:825–831.
8. Lee CD, Blair SN. (2002) Cardiorespiratory fitness and smoking-related and total cancer mortality in men. *Med Sci Sports Exerc* 34:735–739.
9. Sawada SS, Muto T, Tanaka H, et al. (2003) Cardiorespiratory fitness and cancer mortality in Japanese men: a prospective study. *Med Sci Sports Exerc* 35:1546–1550.
10. Peel JB, Sui X, Matthews CE, et al. (2009) Cardiorespiratory fitness and digestive cancer mortality: findings from the aerobics center longitudinal study. *Cancer Epidemiol Biomarkers Prev* 18:1111–1117.
11. Peel JB, Sui X, Adams SA, Hebert JR, Hardin JW, Blair SN. (2009) A prospective study of cardiorespiratory fitness and breast cancer mortality. *Med Sci Sports Exerc* 41:742–748.
12. Arraiz GA, Wigle DT, Mao Y. (1992) Risk assessment of physical activity and physical fitness in the Canada health survey mortality follow-up study. *J Clin Epidemiol* 45:419–428.
13. Haskell WL, Lee IM, Pate RR, et al. (2007) Physical activity and public health: updated recommendation for adults from the American college of sports medicine and the American heart association. *Med Sci Sports Exerc* 39:1423–1434.
14. Department of Health 2004 Chief Medical Officer's Report. Evidence on the impact of physical activity and its relationship to health. 2004. London, HSMO.
15. Leitzmann MF, Park Y, Blair A, et al. (2007) Physical activity recommendations and decreased risk of mortality. *Arch Intern Med* 167:2453–2460.
16. Orsini N, Mantzoros CS, Wolk A. (2008) Association of physical activity with cancer incidence, mortality, and survival: a population-based study of men. *Br J Cancer* 98:1864–1869.
17. Inoue M, Iso H, Yamamoto S, et al. (2008) Daily total physical activity level and premature death in men and women: Results from a large-scale population-based cohort study in Japan (JPHC study). *Ann Epidemiol* 18:522–530.
18. Wannamethee G, Shaper AG, Macfarlane PW. (1993) Heart rate, physical activity, and mortality from cancer and other noncardiovascular diseases. *Am J Epidemiol* 137:735–748.
19. Hu G, Tuomilehto J, Silventoinen K, Barengo NC, Peltonen M, Jousilahti P. (2005) The effects of physical activity and body mass index on cardiovascular, cancer and all-cause mortality among 47 212 middle-aged Finnish men and women. *Int J Obes (Lond)* 29:894–902.
20. Katzmarzyk PT, Church TS, Craig CL, Bouchard C. (2009) Sitting time and mortality from all causes, cardiovascular disease, and cancer. *Med Sci Sports Exerc* 41:998–1005.
21. Rohan TE, Fu W, Hiller JE. (1995) Physical activity and survival from breast cancer. *Eur J Cancer Prev* 4:419–424.
22. Borugian MJ, Sheps SB, Kim-Sing C, et al. (2004) Insulin, macronutrient intake, and physical activity: are potential indicators of insulin resistance associated with mortality from breast cancer? *Cancer Epidemiol Biomarkers Prev* 13:1163–1172.
23. Enger SM, Bernstein L. (2004) Exercise activity, body size and premenopausal breast cancer survival. *Br J Cancer* 90:2138–2141.
24. Abrahamson PE, Gammon MD, Lund MJ, et al. (2006) Recreational physical activity and survival among young women with breast cancer. *Cancer* 107:1777–1785.
25. Friedenreich CM, Gregory J, Kopciuk KA, Mackey JR, Courneya KS. (2009) Prospective cohort study of lifetime physical activity and breast cancer survival. *Int J Cancer* 124: 1954–1962.
26. Pate RR, Pratt M, Blair SN, et al. (1995) Physical activity and public health. A recommendation from the centers for disease control and prevention and the American college of sports medicine. *JAMA* 273:402–407.
27. Haydon AM, Macinnis RJ, English DR, Giles GG. (2006) Effect of physical activity and body size on survival after diagnosis with colorectal cancer. *Gut* 55:62–67.
28. Irwin ML, Crumley D, McTiernan A, et al. (2003) Physical activity levels before and after a diagnosis of breast carcinoma: the health, eating, activity, and lifestyle (HEAL) study. *Cancer* 97:1746–1757.
29. Courneya KS, Friedenreich CM. (1997) Relationship between exercise pattern across the cancer experience and current quality of life in colorectal cancer survivors. *J Altern Complement Med* 3:215–226.
30. Holmes MD, Chen WY, Feskanich D, Kroenke CH, Colditz GA. (2005) Physical activity and survival after breast cancer diagnosis. *JAMA* 293:2479–2486.
31. Irwin ML, McTiernan A, Bernstein L, et al. (2004) Physical activity levels among breast cancer survivors. *Med Sci*

Sports Exerc 36:1484–1491.

32. Pierce JP, Stefanick ML, Flatt SW, et al. (2007) Greater survival after breast cancer in physically active women with high vegetable-fruit intake regardless of obesity. *J Clin Oncol* 25:2345–2351.
33. Holick CN, Newcomb PA, Trentham-Dietz A, et al. (2008) Physical activity and survival after diagnosis of invasive breast cancer. *Cancer Epidemiol Biomarkers Prev* 17:379–386.
34. Irwin ML, Smith AW, McTiernan A, et al. (2008) Influence of pre- and postdiagnosis physical activity on mortality in breast cancer survivors: the health, eating, activity, and lifestyle study. *J Clin Oncol* 26:3958–3964.
35. Meyerhardt JA, Heseltine D, Niedzwiecki D, et al. (2006) Impact of physical activity on cancer recurrence and survival in patients with stage III colon cancer: findings from CALGB 89803. *J Clin Oncol* 24:3535–3541.
36. Meyerhardt JA, Giovannucci EL, Holmes MD, et al. (2006) Physical activity and survival after colorectal cancer diagnosis. *J Clin Oncol* 24:3527–3534.
37. Scardino PT, Weaver R, Hudson MA. (1992) Early detection of prostate cancer. *Hum Pathol* 23:211–222.
38. Giovannucci E, Leitzmann M, Spiegelman D, et al. (1998) A prospective study of physical activity and prostate cancer in male health professionals. *Cancer Res* 58: 5117–5122.
39. Giovannucci EL, Liu Y, Leitzmann MF, Stampfer MJ, Willett WC. (2005) A prospective study of physical activity and incident and fatal prostate cancer. *Arch Intern Med* 165: 1005–1010.
40. Patel AV, Rodriguez C, Jacobs EJ, Solomon L, Thun MJ, Calle EE. (2005) Recreational physical activity and risk of prostate cancer in a large cohort of US men. *Cancer Epidemiol Biomarkers Prev* 14:275–279.
41. Nilsen TI, Romundstad PR, Vatten LJ. (2006) Recreational physical activity and risk of prostate cancer: a prospective population-based study in Norway (the HUNT study). *Int J Cancer* 119:2943–2947.
42. Crespo CJ, Garcia-Palmeri MR, Smit E, et al. (2008) Physical activity and prostate cancer mortality in Puerto Rican men. *J Phys Act Health* 5:918–929.
43. Moore SC, Peters TM, Ahn J, et al. (2008) Physical activity in relation to total, advanced, and fatal prostate cancer. *Cancer Epidemiol Biomarkers Prev* 17:2458–2466.
44. Ornish D, Weidner G, Fair WR, et al. (2005) Intensive lifestyle changes may affect the progression of prostate cancer. *J Urol* 174:1065–1069.
45. Tayek JA, Heber D, Byerley LO, Steiner B, Rajfer J, Swerdloff RS. (1990) Nutritional and metabolic effects of gonadotropin-releasing hormone agonist treatment for prostate cancer. *Metabolism* 39:1314–1319.
46. Smith MR, Finkelstein JS, McGovern FJ, et al. (2002) Changes in body composition during androgen deprivation therapy for prostate cancer. *J Clin Endocrinol Metab* 87:599–603.
47. Braga-Basaria M, Dobs AS, Muller DC, et al. (2006) Metabolic syndrome in men with prostate cancer undergoing long-term androgen-deprivation therapy. *J Clin Oncol* 24: 3979–3983.
48. Braga-Basaria M, Muller DC, Carducci MA, Dobs AS, Basaria S. (2006) Lipoprotein profile in men with prostate cancer undergoing androgen deprivation therapy. *Int J Impot Res* 18: 494–498.
49. Basaria S, Muller DC, Carducci MA, Egan J, Dobs AS. (2006) Hyperglycemia and insulin resistance in men with prostate carcinoma who receive androgen-deprivation therapy. *Cancer* 106:581–588.
50. Saigal CS, Gore JL, Krupski TL, Hanley J, Schonlau M, Litwin MS. (2007) Androgen deprivation therapy increases cardiovascular morbidity in men with prostate cancer. *Cancer* 110:1493–1500.
51. Tsai HK, D’Amico AV, Sadetsky N, Chen MH, Carroll PR. (2007) Androgen deprivation therapy for localized prostate cancer and the risk of cardiovascular mortality. *J Natl Cancer Inst* 99:1516–1524.
52. Hamer M, Stamatakis E, Saxton JM. (2009) The impact of physical activity on all-cause mortality in men and women after a cancer diagnosis. *Cancer Causes Control* 20:225–231.
53. Fairey AS, Courneya KS, Field CJ, Bell GJ, Jones LW, Mackey JR. (2005) Randomized controlled trial of exercise and blood immune function in postmenopausal breast cancer survivors. *J Appl Physiol* 98:1534–1540.
54. Hutnick NA, Williams NI, Kraemer WJ, et al. (2005) Exercise and lymphocyte activation following chemotherapy for breast cancer. *Med Sci Sports Exerc* 37:1827–1835.
55. Meijer EP, Goris AH, van Dongen JL, Bast A, Westerterp KR. (2002) Exercise-induced oxidative stress in older adults as a function of habitual activity level. *J Am Geriatr Soc* 50:349–353.
56. Campbell KL, McTiernan A. (2007) Exercise and biomarkers for cancer prevention studies. *J Nutr* 137:161S–169S.
57. Leung PS, Aronson WJ, Ngo TH, Golding LA, Barnard RJ. (2004) Exercise alters the IGF axis in vivo and increases p53 protein in prostate tumor cells in vitro. *J Appl Physiol* 96:450–454.

1. Ness KK, Wall MM, Oakes M, Robison LL, Gurney JG. (2006) Physical performance limitations and participation restrictions among cancer survivors: a population-based study. *Ann Epidemiol* 16:197–205.
2. Winningham ML, MacVicar MG, Burke CA. (1986) Exercise for cancer patients: guidelines and precautions. *Physician Sportsmed* 14:134–135.
3. Schmitz KH, Holtzman J, Courneya KS, Masse LC, Duval S, Kane R. (2005) Controlled physical activity trials in cancer survivors: a systematic review and meta-analysis. *Cancer Epidemiol Biomarkers Prev* 14:1588–1595.
4. McNeely ML, Campbell KL, Rowe BH, Klassen TP, Mackey JR, Courneya KS. (2006) Exercise interventions in breast cancer patients: a systematic review and meta-analysis of randomized controlled trials. *Can Med Assoc J* 175:34–41.
5. Holmes M, Chen WY, Feskanich D, Kroenke CH, Colditz GA. (2005) Physical activity and survival after breast cancer diagnosis. *JAMA* 293:2479–2486.
6. Holick CN, Newcomb PA, Trentham-Dietz A, Titus-Ernstoff L, Bersch AJ, Stampfer MJ, Baron JA, Egan KM, Willett WC. (2008) Physical activity and survival after diagnosis of invasive breast cancer. *Cancer Epidemiol Biomarkers Prev* 17:379–386.
7. Meyerhardt JA, Giovannucci EL, Holmes MD, Chan AT, Chan JA, Colditz GA, Fuchs CS. (2006) Physical activity and survival after colorectal cancer diagnosis. *J Clin Oncol* 24: 3527–3534.
8. Meyerhardt JA, Heseltine D, Niedzwiecki D, Hollis D, Saltz LB, Mayer RJ, Thomas J, Nelson H, Whittom R, Hantel A, Schilsky RL, Fuchs CS. (2006) Impact of physical activity on cancer recurrence and survival in patients with stage III colon cancer: findings from CALGB 89803. *J Clin Oncol* 24:3535–3541.
9. World Cancer Research Fund/American Institute for Cancer Research. (2007) *Food, Nutrition, Physical Activity and the Prevention of Cancer: A Global Perspective*. Washington, DC, American Institute for Cancer Research.
10. Cancer Council Australia. (2006) *Benefits of Healthy Diet and Physical Activity for Cancer Survivors (Position Statement)*. Sydney, Cancer Council Australia.
11. Doyle C, Kushi LH, Byers T, Courneya KS, Demark-Wahnefried W, Grant B, McTiernan A, Rock CL, Thompson C, Gansler T, Andrews KS. (2006) *Nutrition and physical activity during and after cancer treatment: an American Cancer Society guide for informed choices*. *CA Cancer J Clin* 56:323–353.
12. Demark-Wahnefried W, Aziz NM, Rowland JH, Pinto BM. (2005) Riding the crest of the teachable moment: promoting long-term health after the diagnosis of cancer. *J Clin Oncol* 23:5814–5830.
13. Demark-Wahnefried W, Peterson B, McBride C, Lipkus I, Clipp E. (2000) Current health behaviors and readiness to pursue life-style changes among men and women diagnosed with early stage prostate and breast carcinomas. *Cancer* 88:674–684.
14. Blanchard CM, Cokkinides V, Courneya KS, Nehl EJ, Stein K, Baker F. (2003) A comparison of physical activity of posttreatment breast cancer survivors and noncancer controls. *Behav Med* 28:140–149.
15. Irwin ML, McTiernan A, Bernstein L, Gilliland FD, Baumgartner R, Baumgartner K, Ballard-Barbash R. (2004) Physical activity levels among breast cancer survivors. *Med Sci Sports Exerc* 36:1484–1491.
16. Stevinson C, Faught W, Steed H, Tonkin K, Ladha AB, Vallance JK, Capstick V, Schepansky A, Courneya KS. (2006) Associations between physical activity and quality of life in ovarian cancer survivors. *Gynecol Oncol* 106:244–250.
17. Courneya KS, Karvinen KH, Campbell KL, Pearcey RG, Dundas G, Capstick V, Tonkin KS. (2005) Associations among exercise, body weight, and quality of life in a population-based sample of endometrial cancer survivors. *Gynecol Oncol* 97:422–430.
18. Karvinen KH, Courneya KS, North S, Venner P. (2007) Associations between exercise and quality of life in bladder cancer survivors: a population-based study. *Cancer Epidemiol Biomarkers Prev* 16:984–990.
19. Jones LW, Courneya KS, Vallance JK, Ladha AB, Mant MJ, Belch AR, Stewart DA, Reiman T. (2004) Association between exercise and quality of life in multiple myeloma cancer survivors. *Support Care Cancer* 12:780–788.
20. Vallance JK, Courneya KS, Jones LW, Reiman T. (2005) Differences in quality of life between non-Hodgkin's lymphoma survivors meeting and not meeting public health exercise guidelines. *Psychooncology* 14:979–991.
21. Rogers LQ, Courneya KS, Robbins KT, Malone J, Seiz A, Koch L, Rao K, Nagarkar M. (2006) Physical activity and quality of life in head and neck cancer survivors. *Support Care Cancer* 14:1012–1019.
22. Jones LW, Guill B, Keir ST, Carter BSK, Friedman HS, Bigner DD, Reardon DA. (2006) Patterns of exercise across the cancer trajectory in brain tumor patients. *Cancer* 106: 2224–3222.
23. Gaskin TA, LoBuglio A, Kelly P, Doss M, Pizitz N. (1989) STRETCH: a rehabilitative program for patients with

- breast cancer. *South Med J* 82:467–469.
24. Seifert E, Ewert S, Werle J. (1992) Physical training in the aftercare of patients with cancer of the head and neck [German]. *Rehabilitation* 31:33–37.
 25. Schulz K, Szlovak C, Schulz H, Gold S, Brechtel L, Braumann M, Koch U. (1998) Implementation and evaluation of an exercise rehabilitation training for breast cancer patients [German]. *Psychother Psychosom Med Psychol* 48:398–407.
 26. Durak EP, Lilly PC. (1998) The application of an exercise and wellness program for cancer patients: a preliminary outcomes report. *J Strength Cond Res* 6:3–6.
 27. Adamsen L, Rasmussen JM, Pedersen LS. (2001) ‘Brothers in arms’: how men with cancer experience a sense of comradeship through group intervention which combines physical activity with information relay. *J Clin Nurs* 10:528–537.
 28. Kolden G, Strauman TJ, Ward A, Kuta J, Woods TE, Schneider KL, Heerey E, Sanborn L, Burt C, Millbrandt L, Kalin NH, Stewart JA, Mullen B. (2002) A pilot study of group exercise training (GET) for women with primary breast cancer: feasibility and health benefits. *Psychooncology* 11:447–456.
 29. Adamsen L, Midtgaard J, Rorth M, Borregaard N, Andersen C, Quist M, Møller T, Zacho M, Madsen JK, Knutsen L. (2003) Feasibility, physical capacity, and health benefits of a multidimensional exercise program for cancer patients undergoing chemotherapy. *Support Care Cancer* 11:707–716.
 30. Young-McCaughan S, Mays MZ, Arzola SM, Yoder LH, Dramiga SA, Leclerc KM, Caton JR, Sheffler RL, Nowlin MU. (2003) Change in exercise tolerance, activity and sleep patterns, and quality of life in patients with cancer participating in a structured exercise program. *Oncol Nurs Forum* 30:441–454.
 31. Turner J, Hayes S, Reul-Hirche H. (2004) Improving the physical status and quality of life of women treated for breast cancer: a pilot study of a structured exercise intervention. *J Surg Oncol* 86:141–146.
 32. Christopher KA, Morrow LL. (2004) Evaluating a community-based exercise program for women cancer survivors. *Appl Nurs Res* 17:100–108.
 33. Sandel SL, Judge JO, Landry N, Faria L, Ouellette R, Majczak M. (2005) Dance and movement program improves quality-of-life measures in breast cancer survivors. *Cancer Nurs* 28:301–309.
 34. Campbell A, Mutrie N, White F, McGuire F, Kearney N. (2005) A pilot study of a supervised group exercise programme as a rehabilitation treatment for women with breast cancer receiving adjuvant treatment. *Eur J Oncol Nurs* 9:56–63.
 35. Adamsen L, Quist M, Midtgaard J, Andersen C, Møller T, Knutsen L, Tveterås A, Rorth M. (2006) The effect of a multidimensional exercise intervention on physical capacity, wellbeing and quality of life in cancer patients undergoing chemotherapy. *Support Care Cancer* 14:116–127.
 36. Losito J, Murphy S, Thomas M. (2006) The effects of group exercise on fatigue and quality of life during cancer treatment. *Oncol Nurs Forum* 33:821–825.
 37. Stevinson C, Fox KR. (2006) Feasibility of an exercise rehabilitation programme for cancer patients. *Eur J Cancer Care* 15:386–396.
 38. Midtgaard J, Rorth M, Stelter R, Adamsen L. (2006) The group matters: an explorative study of group cohesion and quality of life in cancer patients participating in physical exercise intervention during treatment. *Eur J Cancer Care* 15:25–33.
 39. Kramer MM. (1996) The effects of exercise on psychological well-being in women recovering from breast cancer. *Diss Abstr Int* 57:2157.
 40. Schwartz AL. (2000) Exercise and weight gain in breast cancer patients receiving chemotherapy. *Cancer Pract* 8:231–237.
 41. Schwartz A, Mori M, Gao R, Nail LM, King ME. (2001) Exercise reduces daily fatigue in women with breast cancer receiving chemotherapy. *Med Sci Sports Exerc* 33:718–723.
 42. Schwartz AL, Thompson JA, Masood N. (2002) Interferon-induced fatigue in patients with melanoma: a pilot study of exercise and methylphenidate. *Oncol Nurs Forum* 29:85–90.
 43. Wilson DB, Porter JS, Parker G, Kilpatrick J. (2005) Anthropometric changes using a walking intervention in African American breast cancer survivors: a pilot study. *Prev Chronic Dis* 2:1–7.
 44. Shore S, Shepard RJ. (1999) Immune responses to exercise in children treated for cancer. *J Sports Med Phys Fitness* 39:240–243.
 45. Oldervoll L, Kaasa S, Knobel H, Loge JH. (2003) Exercise reduces fatigue in chronic fatigued Hodgkins disease

- survivors - results from a pilot study. *Eur J Cancer* 39:57–63.
46. Segal R, Evans W, Johnson D, Smith J, Colletta S, Gayton J, Woodard S, Wells G, Reid R. (2001) Structured exercise improves physical functioning in women with stages I and II breast cancer: Results of a randomized controlled trial. *J Clin Oncol* 19:657–665.
 47. Midtgaard J, Tveterås A, Rørth M, Stelter R, Adamsen L. (2006) The impact of supervised exercise intervention on short-term postprogram leisure time physical activity level in cancer patients undergoing chemotherapy: 1- and 3-month follow-up on the body & cancer project. *Palliat Support Care* 4:25–35.
 48. von Gruenigen VE, Courneya KS, Gibbons HE, Kavanagh MB, Waggoner SE, Lerner E. (2008) Feasibility and effectiveness of a lifestyle intervention program in obese endometrial cancer patients: a randomized trial. *Gynecol Oncol* 109:19–26.
 49. Carmack Taylor CL, Demoor C, Smith MA, Dunn AL, Basen-Engquist K, Nielsen I, Pettaway C, Sellin R, Massey P, Gritz ER. (2006) Active for life after cancer: a randomized trial examining a lifestyle physical activity program for prostate cancer patients. *Psychooncology* 15:847–862.
 50. Basen-Engquist K, Taylor CL, Rosenblum C, Smith MA, Shinn EH, Greisinger A, Gregg X, Massey P, Valero V, Rivera E. (2006) Randomized pilot test of a lifestyle physical activity intervention for breast cancer survivors. *Patient Educ Couns* 64:225–234.
 51. Damush TM, Perkins A, Miller K. (2006) The implementation of an oncologist referred, exercise self-management program for older breast cancer survivors. *Psychooncology* 15:884–890.
 52. Vallance JKH, Courneya KS, Plotnikoff RC, Yasui Y, Mackey JR. (2007) Randomized controlled trial of the effects of print materials and step pedometers on physical activity and quality of life in breast cancer survivors. *J Clin Oncol* 25:2352–2359.
 53. Vallance JK, Courneya KS, Plotnikoff RC, Dinu I, Mackey JR. (2008) Maintenance of physical activity in breast cancer survivors after a randomized trial. *Med Sci Sports Exerc* 40:173–180.
 54. Leddy SK. (1997) Incentives and barriers to exercise in women with a history of breast cancer. *Oncol Nurs Forum* 24:885–890.
 55. Brawley LR, Culos-Reed SN, Angove J, Hoffman-Goetz L. (2002) Understanding the barriers to physical activity for cancer patients: review and recommendations. *J Psychosocial Oncol* 20:1–21.
 56. Zunft HJ, Friebe D, Seppelt B, Widhalm K, Remaut de Winter AM, Vaz de Almeida MD, Kearney JM, Gibney M. (1999) Perceived benefits and barriers to physical activity in a nationally representative sample in the European union. *Public Health Nutr* 2:153–160.
 57. Canadian Fitness and Lifestyle Research Institute. (1996) Barriers to physical activity. *Progress in Prevention Bulletin* 4:1–6.
 58. Courneya KS, Friedenreich CM, Quinney HA, Fields AL, Jones LW, Vallance JK, Fairey AS. (2005) A longitudinal study of exercise barriers in colorectal cancer survivors participating in a randomized controlled trial. *Ann Behav Med* 29:147–153.
 59. Courneya KS, Vallance JKH, Jones LW, Reiman T. (2005) Correlates of exercise intentions in non-Hodgkin's lymphoma survivors: an application of the theory of planned behavior. *J Sport Exerc Psych* 27:335–349.
 60. Servaes P, Verhagen C, Bleijenberg G. (2002) Fatigue in cancer patients during and after treatment: prevalence, correlates and interventions. *Eur J Cancer* 38:27–43.
 61. Stevinson C, Tonkin K, Capstick V, Schepansky A, Ladha AB, Vallance JK, Faught W, Steed H, Courneya KS. (in press) A population-based study of the determinants of physical activity in ovarian cancer survivors. *J Phys Activity Health*.
 62. Yoshioka H. (1994) Rehabilitation for the terminal cancer patient. *Am J Phys Med Rehabil* 73:199–206.
 63. Porock D, Kristjanson LJ, Tinnelly K, Duke T, Blight J. (2000) An exercise intervention for advanced cancer patients experiencing fatigue: a pilot study. *J Palliat Care* 16:30–36.
 64. Oldervoll LM, Loge JH, Paltiel H, Asp MB, Vidvei U, Wiken AN, Hjermstad MJ, Kaasa S. (2006) The effect of a physical exercise program in palliative care: a phase II study. *J Pain Symptom Manage* 31:421–430.
 65. Dimeo FC, Bertz H, Finke J, Fetscher S, Mertelsmann R, Keul J. (1996) An aerobic exercise program for patients with haematological malignancies after bone marrow transplantation. *Bone Marrow Transplant* 18:1157–1160.
 66. Stevinson C, Lawlor DA, Fox KR. (2004) Exercise interventions for cancer patients: systematic review of controlled trials. *Cancer Causes Control* 15:1035–1056.
 67. McNeely ML, Peddle C, Parliament M, Courneya KS. (2006) Cancer rehabilitation: recommendations

for integrating exercise programming in the clinical practice setting. *Curr Cancer Ther Rev* 2:351–360.

68. Harris SR, Niesen-Vertommen SL. (2000) Challenging the myth of exercise-induced lymphedema following breast cancer: a series of case reports. *J Surg Oncol* 74:95–98.
69. McKenzie DC, Kalda AL. (2003) Effect of upper extremity exercise on secondary lymphedema in breast cancer patients: a pilot study. *J Clin Oncol* 21:463–466.
70. Lane K, Jespersen D, McKenzie DC. (2005) The effect of a whole body exercise programme and dragon boat training on arm volume and arm circumference in women treated for breast cancer. *Eur J Cancer Care* 14:353–358.
71. Ahmed RL, Thomas W, Yee D, Schmitz KH. (2006) Randomized controlled trial of weight training and lymphedema in breast cancer survivors. *J Clin Oncol* 24:2765–2772.
72. Lane K, Worsley D, McKenzie D. (2005) Exercise and the lymphatic system: implications for breast-cancer survivors. *Sports Med* 35:461–471.
73. Mahon S. (1998) Osteoporosis: a concern for cancer survivors. *Oncol Nurs Forum* 25: 843–851.
74. Brown BW, Brauner C, Minnotte MC. (1993) Noncancer deaths in white adult cancer patients. *J Natl Cancer Inst* 85:979–987.

1. Church TS, Cheng YJ, Earnest CP, et al. (Jan 2004) Exercise capacity and body composition as predictors of mortality among men with diabetes. *Diabetes Care* 27(1):83–88.
2. Gulati M, Black HR, Shaw LJ, et al. (Aug 4 2005) The prognostic value of a nomogram for exercise capacity in women. *N Engl J Med* 353(5):468–475.
3. Gulati M, Pandey DK, Arnsdorf MF, et al. (Sep 30 2003) Exercise capacity and the risk of death in women: the St James women take heart project. *Circulation* 108(13):1554–1559.
4. Myers J, Prakash M, Froelicher V, et al. (Mar 14 2002) Exercise capacity and mortality among men referred for exercise testing. *N Engl J Med* 346(11):793–801.
5. ATS/ACCP Statement on cardiopulmonary exercise testing. (Jan 15 2003) *Am J Respir Crit Care Med* 167(2):211–277.
6. Beckles MA, Spiro SG, Colice GL. (Jan 2003) The physiologic evaluation of patients with lung cancer being considered for resectional surgery. *Chest* 123(1 Suppl):105S–114S.
7. Brunelli A, Al Refai M, Monteverde M, et al. (Apr 2002) Stair climbing test predicts cardiopulmonary complications after lung resection. *Chest* 121(4):1106–1110.
8. Brunelli A, Monteverde M, Salati M, et al. (Nov 2001) Stair-climbing test to evaluate maximum aerobic capacity early after lung resection. *Ann Thorac Surg* 72(5):1705–1710.
9. Loewen GM, Watson D, Kohman L, et al. (Jul 2007) Preoperative exercise VO₂ measurement for lung resection candidates: Results of cancer and leukemia group B protocol 9238. *J Thorac Oncol* 2(7):619–625.
10. Nomori H, Watanabe K, Ohtsuka T, et al. (Mar 2004) Six-minute walking and pulmonary function test outcomes during the early period after lung cancer surgery with special reference to patients with chronic obstructive pulmonary disease. *Jpn J Thorac Cardiovasc Surg* 52(3):113–119.
11. Pelletier C, Lapointe L, LeBlanc P. (Jul 1990) Effects of lung resection on pulmonary function and exercise capacity. *Thorax* 45(7):497–502.
12. Win T, Groves AM, Ritchie AJ, et al. (Jun 2007) The effect of lung resection on pulmonary function and exercise capacity in lung cancer patients. *Respir Care* 52(6):720–726.
13. Win T, Jackson A, Groves AM, et al. (Jan 2006) Comparison of shuttle walk with measured peak oxygen consumption in patients with operable lung cancer. *Thorax* 61(1):57–60.
14. Carlson LE, Smith D, Russell J, et al. (May 2006) Individualized exercise program for the treatment of severe fatigue in patients after allogeneic hematopoietic stem-cell transplant: a pilot study. *Bone Marrow Transplant* 37(10):945–954.
15. Courneya KS, Friedenreich CM, Quinney HA, et al. (Dec 2003) A randomized trial of exercise and quality of life in colorectal cancer survivors. *Eur J Cancer Care (Engl)* 12(4): 347–357.
16. Courneya KS, Friedenreich CM, Sela RA, et al. (Jun 2003) The group psychotherapy and home-based physical exercise (group-hope) trial in cancer survivors: physical fitness and quality of life outcomes. *Psychooncology* 12(4):357–374.
17. Courneya KS, Segal RJ, Mackey JR, et al. (Oct 1 2007) Effects of aerobic and resistance exercise in breast cancer patients receiving adjuvant chemotherapy: a multicenter randomized controlled trial. *J Clin Oncol* 25(28):4396–4404.

18. Daley AJ, Crank H, Saxton JM, et al. (May 1 2007) Randomized trial of exercise therapy in women treated for breast cancer. *J Clin Oncol* 25(13):1713–1721.
19. Dimeo F, Bertz H, Finke J, et al. (Dec 1996) An aerobic exercise program for patients with haematological malignancies after bone marrow transplantation. *Bone Marrow Transplant* 18(6):1157–1160.
20. Dimeo FC, Tilmann MH, Bertz H, et al. (May 1 1997) Aerobic exercise in the rehabilitation of cancer patients after high dose chemotherapy and autologous peripheral stem cell transplantation. *Cancer* 79(9):1717–1722.
21. Jones LW, Eves ND, Mackey JR, et al. (Feb 2007) Safety and feasibility of cardiopulmonary exercise testing in patients with advanced cancer. *Lung Cancer* 55(2):225–232.
22. Jones LW, Haykowsky M, Peddle CJ, et al. (May 2007) Cardiovascular risk profile of patients with HER2/neu-positive breast cancer treated with anthracycline-taxane-containing adjuvant chemotherapy and/or trastuzumab. *Cancer Epidemiol Biomarkers Prev* 16(5): 1026–1031.
23. Jones LW, Haykowsky M, Pituskin EN, et al. (Oct 2007) Cardiovascular reserve and risk profile of postmenopausal women after chemoendocrine therapy for hormone receptor – positive operable breast cancer. *Oncologist* 12(10):1156–1164.
24. Jones LW, Peddle CJ, Eves ND, et al. (Aug 1 2007) Effects of presurgical exercise training on cardiorespiratory fitness among patients undergoing thoracic surgery for malignant lung lesions. *Cancer* 110(3):590–598.
25. Jones LW, Eves ND, Haykowsky M, et al. (Aug 2008) Cardiorespiratory exercise testing in clinical oncology research: systematic review and practice recommendations. *Lancet Oncol* 9(8):757–765.
26. Fleg JL, Pina IL, Balady GJ, et al. (Sep 26 2000) Assessment of functional capacity in clinical and research applications: an advisory from the committee on exercise, rehabilitation, and prevention, Council on Clinical Cardiology, American Heart Association. *Circulation* 102(13):1591–1597.
27. Brooks D, Solway S, Gibbons WJ. (May 1 2003) ATS statement on six-minute walk test. *Am J Respir Crit Care Med* 167(9):1287.
28. Jones LW, Cohen RR, Mabe SK. (Feb 11 2009) Assessment of physical functioning in recurrent glioma: preliminary comparison of performance status to functional capacity testing. *J Neurooncol* 94:79–85.
29. Lederer DJ, Arcasoy SM, Wilt JS, et al. (Sep 15 2006) Six-minute-walk distance predicts waiting list survival in idiopathic pulmonary fibrosis. *Am J Respir Crit Care Med* 174(6): 659–664.
30. Celli BR, Cote CG, Marin JM, et al. (Mar 4 2004) The body-mass index, airflow obstruction, dyspnea, and exercise capacity index in chronic obstructive pulmonary disease. *N Engl J Med* 350(10):1005–1012.
31. Shah MR, Hasselblad V, Gheorghide M, et al. (Nov 1 2001) Prognostic usefulness of the six-minute walk in patients with advanced congestive heart failure secondary to ischemic or nonischemic cardiomyopathy. *Am J Cardiol* 88(9):987–993.
32. Paciocco G, Martinez FJ, Bossone E, et al. (Apr 2001) Oxygen desaturation on the six-minute walk test and mortality in untreated primary pulmonary hypertension. *Eur Respir J* 17(4): 647–652.
33. Cahalin LP, Mathier MA, Semigran MJ, et al. (Aug 1996) The six-minute walk test predicts peak oxygen uptake and survival in patients with advanced heart failure. *Chest* 110(2): 325–332.
34. Jones LW, Haykowsky MJ, Swartz JJ, et al. (Oct 9 2007) Early breast cancer therapy and cardiovascular injury. *J Am Coll Cardiol* 50(15):1435–1441.
35. Nakagawa K, Nakahara K, Miyoshi S, et al. (May 1992) Oxygen transport during incremental exercise load as a predictor of operative risk in lung cancer patients. *Chest* 101(5):1369–1375.
36. Prosnitz RG, Hubbs JL, Evans ES, et al. (Oct 15 2007) Prospective assessment of radiotherapy-associated cardiac toxicity in breast cancer patients: analysis of data 3 to 6 years after treatment. *Cancer* 110(8):1840–1850.
37. Perez EA, Suman VJ, Davidson NE, et al. (Sep 15 2004) Effect of doxorubicin plus cyclophosphamide on left ventricular ejection fraction in patients with breast cancer in the north central cancer treatment group N9831 intergroup adjuvant trial. *J Clin Oncol* 22(18):3700–3704.
38. Bohlius J, Weingart O, Trelle S, et al. (Mar 2006) Cancer-related anemia and recombinant human erythropoietin – an updated Overview. *Nat Clin Pract Oncol* 3(3):152–164.
39. Ekholm E, Rantanen V, Bergman M. (May–Jun 2000) Docetaxel and autonomic cardiovascular control in anthracycline treated breast cancer patients. *Anticancer Res* 20(3B):2045–2048.
40. Viniegra M, Marchetti M, Losso M, et al. (1990) Cardiovascular autonomic function in anthracycline-treated breast cancer patients. *Cancer Chemother Pharmacol* 26(3):227–231.
41. Mao J, Kocak Z, Zhou S, et al. (Apr 1 2007) The impact of induction chemotherapy and the associated tumor response on subsequent radiation-related changes in lung function and tumor response. *Int J Radiat Oncol Biol Phys* 67(5):1360–1369.
42. Miller KL, Zhou SM, Barrier RC, Jr. (Jul 1 2003) Long-term changes in pulmonary function tests after definitive radiotherapy for lung cancer. *Int J Radiat Oncol Biol Phys* 56(3):611–615.
43. Beckman JA, Thakore A, Kalinowski BH, et al. (Mar 1 2001) Radiation therapy impairs endothelium-dependent

- vasodilation in humans. *J Am Coll Cardiol* 37(3):761–765.
44. Batchelor TT, Taylor LP, Thaler HT, et al. (May 1997) Steroid myopathy in cancer patients. *Neurology* 48(5):1234–1238.
 45. Dropcho EJ, Soong SJ. (Aug 1991) Steroid-induced weakness in patients with primary brain tumors. *Neurology* 41(8):1235–1239.
 46. Larsen KR, Svendsen UG, Milman N, et al. (Oct 1997) Cardiopulmonary function at rest and during exercise after resection for bronchial carcinoma. *Ann Thorac Surg* 64(4):960–964.
 47. Jones LW, Eves ND, Haykowsky M. (2009). The physiologic and molecular mechanisms of exercise intolerance and the role of exercise therapy to reverse dysfunction in cancer. *Lancet Oncol* in press 10:598–605.
 48. Jones LW, Eves ND, Peterson BL, et al. (Dec 15 2008) Safety and feasibility of aerobic training on cardiopulmonary function and quality of life in postsurgical nonsmall cell lung cancer patients: a pilot study. *Cancer* 113(12):3430–3439.
 49. Jensen D, Amjadi K, Harris-McAllister V. (Feb 4 2008) Mechanisms of dyspnoea relief and improved exercise endurance after furosemide inhalation in COPD. *Thorax* 63:606–13.
 50. Spruit MA, Troosters T, Gosselink R, et al. (2007) Acute inflammatory and anabolic systemic responses to peak and constant-work-rate exercise bout in hospitalized patients with COPD. *Int J Chron Obstruct Pulmon Dis* 2(4):575–583.
 51. van't Hul A, Gosselink R, Kwakkel G. (Mar–Apr 2003) Constant-load cycle endurance performance: test-retest reliability and validity in patients with COPD. *J Cardiopulm Rehabil* 23(2):143–150.
 52. Jannik VK, Gledhill N, Shephard RJ. (Dec 2007) Revised clearance for participation in physical activity: greater screening responsibility for qualified university-educated fitness professionals. *Appl Physiol Nutr Metab* 32(6):1191–1197.
 53. Hermansen L, Saltin B. (Jan 1969) Oxygen uptake during maximal treadmill and bicycle exercise. *J Appl Physiol* 26(1):31–37.
 54. Esteban E, Villanueva N, Muniz I, et al. (Feb 2008) Pulmonary toxicity in patients treated with gemcitabine plus vinorelbine or docetaxel for advanced non-small cell lung cancer: outcome data on a randomized phase II study. *Invest New Drugs* 26(1):67–74.
 55. Fitzgerald MD, Tanaka H, Tran ZV, et al. (Jul 1997) Age-related declines in maximal aerobic capacity in regularly exercising vs. Sedentary women: a meta-analysis. *J Appl Physiol* 83(1):160–165.
 56. Wilson TM, Tanaka H. (Mar 2000) Meta-analysis of the age-associated decline in maximal aerobic capacity in men: relation to training status. *Am J Physiol Heart Circ Physiol* 278(3):H829–H834.
 57. Casaburi R, Kukafka D, Cooper CB, et al. (Mar 2005) Improvement in exercise tolerance with the combination of tiotropium and pulmonary rehabilitation in patients with COPD. *Chest* 127(3):809–817.
 58. Reardon J, Casaburi R, Morgan M, et al. (Dec 2005) Pulmonary rehabilitation for COPD. *Respir Med* 99(Suppl B):S19–S27.
 59. Borg G, Dahlstrom H. (1962) A case study of perceived exertion during a work test. *Acta Soc Med Ups* 67:91–93