

REFERENCES

Chapter 1

1. Aagaard, P, Andersen, JL, Dyhre-Poulsen, P, Leffers, AM, Wagner, A, Magnusson, SP, Halkjaer-Kristensen, J, and Simonsen, EB. A mechanism for increased contractile strength of human pennate muscle in response to strength training: changes in muscle architecture. *J. Physiol.* 534: 613-623, 2001.
2. Aagaard, P, Simonsen, EB, Andersen, JL, Magnusson, P, and Dyhre-Poulsen, P. Neural adaptation to resistance training: changes in evoked V-wave and H-reflex responses. *J. Appl. Physiol.* (1985) 92: 2309-2318, 2002.
3. Aagaard, P, Simonsen, EB, Andersen, JL, Magnusson, P, and Dyhre-Poulsen, P. Increased rate of force development and neural drive of human skeletal muscle following resistance training. *J. Appl. Physiol.* (1985) 93: 1318-1326, 2002.
4. Abernethy, PJ, Jurimae, J, Logan, PA, Taylor, AW, and Thayer, RE. Acute and chronic response of skeletal muscle to resistance exercise. *Sports Med.* 17: 22-38, 1994.
5. Adams, G. The Molecular Response of Skeletal Muscle to Resistance Training. *Deutsche Zeitschrift für Sportmedizin* 61: 61-67, 2010.
6. Adams, G, and Bamman, MM. Characterization and regulation of mechanical loading-induced compensatory muscle hypertrophy. *Comprehensive Physiology* 2829, 2012.
7. Adams, GR, and McCue, SA. Localized infusion of IGF-I results in skeletal muscle hypertrophy in rats. *J. Appl. Physiol.* 84: 1716-1722, 1998.
8. Adams, GR. Invited Review: Autocrine/paracrine IGF-I and skeletal muscle adaptation. *J. Appl. Physiol.* 93: 1159-1167, 2002.
9. Ahtiainen, JP, Pakarinen, A, Alen, M, Kraemer, WJ, and Häkkinen, K. Muscle hypertrophy, hormonal adaptations and strength development during strength training in strength-trained and untrained men. *Eur. J. Appl. Physiol.* 89: 555-563, 2003.
10. Ahtiainen, JP, Hulmi, JJ, Kraemer, WJ, Lehti, M, Nyman, K, Selanne, H, Alen, M, Pakarinen, A, Komulainen, J, Kovanen, V, Mero, AA, and Häkkinen, K. Heavy resistance exercise training and skeletal muscle androgen receptor expression in younger and older men. *Steroids* 76: 183-192, 2011.
11. Alegre, LM, Jimenez, F, Gonzalo-Orden, JM, Martin-Acero, R, and Aguado, X. Effects of dynamic resistance training on fascicle length and isometric strength. *J. Sports Sci.* 24: 501-508, 2006.
12. Alway, SE, Gonyea, WJ, and Davis, ME. Muscle fiber formation and fiber hypertrophy during the onset of stretch-overload. *Am. J. Physiol.* 259: C92-102, 1990.
13. Amiridis, IG, Martin, A, Morlon, B, Martin, L, Cometti, G, Pousson, M, and van Hoecke, J. Co-activation and tension-regulating phenomena during isokinetic knee extension in sedentary and highly skilled humans. *Eur. J. Appl. Physiol. Occup. Physiol.* 73: 149-156, 1996.
14. Andersen, JL, Klitgaard, H, and Saltin, B. Myosin heavy chain isoforms in single fibres from m. vastus lateralis of sprinters: influence of training. *Acta Physiol. Scand.* 151: 135-142, 1994.
15. Andersen, MB, Pingel, J, Kjaer, M, and Langberg, H. Interleukin-6: a growth factor stimulating collagen synthesis in human tendon. *J. Appl. Physiol.* (1985) 110: 1549-1554, 2011.
16. Antonio, J, and Gonyea, WJ. Role of muscle fiber hypertrophy and hyperplasia in intermittently stretched avian muscle. *J. Appl. Physiol.* 74: 1893-1898, 1993.
17. Antonio, J, and Gonyea, WJ. Progressive stretch overload of skeletal muscle results in hypertrophy before hyperplasia. *J. Appl. Physiol.* (1985) 75: 1263-1271, 1993.
18. Aperghis, M, Velloso, CP, Hameed, M, Brothwood, T, Bradley, L, Bouloux, PM, Harridge, SD, and Goldspink, G. Serum IGF-I levels and IGF-I gene splicing in muscle of healthy young males receiving rhGH. *Growth Horm. IGF Res.* 19: 61-67, 2009.
19. Atherton, PJ, and Smith, K. Muscle protein synthesis in response to nutrition and exercise. *J. Physiol.* 590: 1049-1057, 2012.
20. Bamman, MM, Petrella, JK, Kim, JS, Mayhew, DL, and Cross, JM. Cluster analysis tests the importance of myogenic gene expression during myofiber hypertrophy in humans. *J. Appl. Physiol.* 102: 2232-2239, 2007.
21. Barton, ER. Viral expression of insulin-like growth factor-I isoforms promotes different responses in skeletal muscle. *J. Appl. Physiol.* 100: 1778-1784, 2006.
22. Barton-Davis, ER, Shoturma, DI, and Sweeney, HL. Contribution of satellite cells to IGF-I induced hypertrophy of skeletal muscle. *Acta Physiol. Scand.* 167: 301-305, 1999.
23. Bazgir, B, Fathi, R, Rezazadeh Valojerdi, M, Mozdziaik, P, and Asgari, A. Satellite Cells Contribution to Exercise Mediated Muscle Hypertrophy and Repair. *Cell. J.* 18: 473-484, 2017.
24. Bellamy, LM, Joanisse, S, Grubb, A, Mitchell, CJ, McKay, BR, Phillips, SM, Baker, S, and Parise, G. The Acute Satellite Cell Response and Skeletal Muscle Hypertrophy following Resistance Training. *PLoS One* 9: e109739, 2014.
25. Bhasin, S, Storer, TW, Berman, N, Callegari, C, Clevenger, B, Phillips, J, Bunnell, TJ, Tricker, R, Shirazi, A, and Casaburi, R. The effects of supraphysiologic doses of testosterone on muscle size and strength in normal men. *N. Engl. J. Med.* 335: 1-7, 1996.
26. Bhasin, S, Woodhouse, L, and Storer, TW. Proof of the effect of testosterone on skeletal muscle. *J. Endocrinol.* 170: 27-38, 2001.
27. Bhasin, S, Woodhouse, L, Casaburi, R, Singh, AB, Mac, RP, Lee, M, Yarasheski, KE, Sinha-Hikim, I, Dzekov, C, Dzekov, J, Magliano, L, and Storer, TW. Older men are as responsive as young men to the anabolic effects of graded doses of testosterone on the skeletal muscle. *J. Clin. Endocrinol. Metab.* 90: 678-688, 2005.
28. Biolo, G, Williams, BD, Fleming, RY, and Wolfe, RR. Insulin action on muscle protein kinetics and amino acid transport during recovery after resistance exercise. *Diabetes* 48: 949-957, 1999.
29. Blazevich, AJ, Gill, ND, Bronks, R, and Newton, RU. Training-specific muscle architecture adaptation after 5-wk training in athletes. *Med. Sci. Sports Exerc.* 35: 2013-2022, 2003.
30. Blazevich, AJ, Cannavan, D, Coleman, DR, and Horne, S. Influence of concentric and eccentric resistance training on architectural adaptation in human quadriceps muscles. *J. Appl. Physiol.* 103: 1565-1575, 2007.
31. Bodnar, D, Geyer, N, Ruzsnavszky, O, Olah, T, Hegyi, B, Sztrettye, M, Fodor, J, Dienes, B, Balogh, A, Papp, Z, Szabo, L, Muller,

- G, Csernoch, L, and Szentesi, P. Hypermuscular mice with mutation in the myostatin gene display altered calcium signalling. *J. Physiol.* 592: 1353-1365, 2014.
32. Borst, SE. Interventions for sarcopenia and muscle weakness in older people. *Age Ageing* 33: 548-555, 2004.
33. Brahm, H, Piehl-Aulin, K, Saltin, B, and Ljunghall, S. Net fluxes over working thigh of hormones, growth factors and biomarkers of bone metabolism during short lasting dynamic exercise. *Calcif. Tissue Int.* 60: 175-180, 1997.
34. Broholm, C, and Pedersen, BK. Leukaemia inhibitory factor—an exercise-induced myokine. *Exerc. Immunol. Rev.* 16: 77-85, 2010.
35. Brook, MS, Wilkinson, DJ, Smith, K, and Atherton, PJ. It's not just about protein turnover: the role of ribosomal biogenesis and satellite cells in the regulation of skeletal muscle hypertrophy. *Eur. J. Sport. Sci.* 19: 952-963, 2019.
36. Bruusgaard, JC, Johansen, IB, Egner, IM, Rana, ZA, and Gundersen, K. Myonuclei acquired by overload exercise precede hypertrophy and are not lost on detraining. *Proc. Natl. Acad. Sci. U. S. A.* 107: 15111-15116, 2010.
37. Burd, NA, West, DW, Staples, AW, Atherton, PJ, Baker, JM, Moore, DR, Holwerda, AM, Parise, G, Rennie, MJ, Baker, SK, and Phillips, SM. Low-load high volume resistance exercise stimulates muscle protein synthesis more than high-load low volume resistance exercise in young men. *PLoS One* 5: e12033, 2010.
38. Burd, NA, Mitchell, CJ, Churchward-Venne, TA, and Phillips, SM. Bigger weights may not beget bigger muscles: evidence from acute muscle protein synthetic responses after resistance exercise. *Appl. Physiol. Nutr. Metab.* 37: 551-554, 2012.
39. Buresh, R, Berg, K, and French, J. The effect of resistive exercise rest interval on hormonal response, strength, and hypertrophy with training. *J. Strength Cond Res* 23: 62-71, 2009.
40. Campos, GER, Luecke, TJ, Wendeln, HK, Toma, K, Hagerman, FC, Murray, TE, Ragg, KE, Ratamess, NA, Kraemer, WJ, and Staron, RS. Muscular adaptations in response to three different resistance-training regimens: specificity of repetition maximum training zones. *Eur. J. Appl. Physiol.* 88: 50-60, 2002.
41. Carolan, B, and Cafarelli, E. Adaptations in coactivation after isometric resistance training. *J. Appl. Physiol.* (1985) 73: 911-917, 1992.
42. Chan, ST, Johnson, AW, Moore, MH, Kapadia, CR, and Dudley, HA. Early weight gain and glycogen-obligated water during nutritional rehabilitation. *Hum. Nutr. Clin. Nutr.* 36: 223-232, 1982.
43. Charette, SL, McEvoy, L, Pyka, G, Snow-Harter, C, Guido, D, Wiswell, RA, and Marcus, R. Muscle hypertrophy response to resistance training in older women. *J. Appl. Physiol.* (1985) 70: 1912-1916, 1991.
44. Christian, JE, and Lawrence, JC. Control of protein synthesis by insulin. *Eurekah Bioscience* 1: 711-721, 2005.
45. Coffey, VG, Shield, A, Canny, BJ, Carey, KA, Cameron-Smith, D, and Hawley, JA. Interaction of contractile activity and training history on mRNA abundance in skeletal muscle from trained athletes. *Am. J. Physiol. Endocrinol. Metab.* 290: E849-55, 2006.
46. Conboy, IM, Conboy, MJ, Wagers, AJ, Girma, ER, Weissman, IL, and Rando, TA. Rejuvenation of aged progenitor cells by exposure to a young systemic environment. *Nature* 433: 760-764, 2005.
47. Cornelison, DD, and Wold, BJ. Single-cell analysis of regulatory gene expression in quiescent and activated mouse skeletal muscle satellite cells. *Dev. Biol.* 191: 270-283, 1997.
48. Crewther, B, Keogh, J, Cronin, J, and Cook, C. Possible stimuli for strength and power adaptation: acute hormonal responses. *Sports Med.* 36: 215-238, 2006.
49. Crewther, BT, Cook, C, Cardinale, M, Weatherby, RP, and Lowe, T. Two emerging concepts for elite athletes: the short-term effects of testosterone and cortisol on the neuromuscular system and the dose-response training role of these endogenous hormones. *Sports Med.* 41: 103-123, 2011.
50. Dahmane, R, Djordjevic, S, Simunic, B, and Valencic, V. Spatial fiber type distribution in normal human muscle. Histochemical and tensiomyographical evaluation. *J. Biomech.* 38: 2451-2459, 2005.
51. Del Vecchio, A, Casolo, A, Negro, F, Scorcelletti, M, Bazzucchi, I, Enoka, R, Felici, F, and Farina, D. The increase in muscle force after 4 weeks of strength training is mediated by adaptations in motor unit recruitment and rate coding. *J. Physiol.* 597: 1873-1887, 2019.
52. Denne, SC, Liechty, EA, Liu, YM, Brechtel, G, and Baron, AD. Proteolysis in skeletal muscle and whole body in response to euglycemic hyperinsulinemia in normal adults. *Am. J. Physiol.* 261: E809-14, 1991.
53. Deshmukh, AS, Murgia, M, Nagaraj, N, Treebak, JT, Cox, J, and Mann, M. Deep proteomics of mouse skeletal muscle enables quantitation of protein isoforms, metabolic pathways, and transcription factors. *Mol. Cell. Proteomics* 14: 841-853, 2015.
54. Doessing, S, Heinemeier, KM, Holm, L, Mackey, AL, Schjerling, P, Rennie, M, Smith, K, Reitelsheder, S, Kappelgaard, AM, Rasmussen, MH, Flyvbjerg, A, and Kjaer, M. Growth hormone stimulates the collagen synthesis in human tendon and skeletal muscle without affecting myofibrillar protein synthesis. *J. Physiol.* 588: 341-351, 2010.
55. Dowling, JJ, Konert, E, Ljucovic, P, and Andrews, DM. Are humans able to voluntarily elicit maximum muscle force? *Neurosci. Lett.* 179: 25-28, 1994.
56. Duchateau, J, Semmler, JG, and Enoka, RM. Training adaptations in the behavior of human motor units. *J. Appl. Physiol.* 101: 1766-1775, 2006.
57. Dumont, NA, Wang, YX, and Rudnicki, MA. Intrinsic and extrinsic mechanisms regulating satellite cell function. *Development* 142: 1572-1581, 2015.
58. Dungan, CM, Murach, KA, Frick, KK, Jones, SR, Crow, SE, Englund, DA, Vechetti, II, Jr, Figueiredo, VC, Levitan, BM, Satin, J, McCarthy, JJ, and Peterson, CA. Elevated myonuclear density during skeletal muscle hypertrophy in response to training is reversed during detraining. *Am. J. Physiol. Cell. Physiol.* 316: C649-C654, 2019.
59. Egner, IM, Bruusgaard, JC, Eftestol, E, and Gundersen, K. A cellular memory mechanism aids overload hypertrophy in muscle long after an episodic exposure to anabolic steroids. *J. Physiol.* 591: 6221-6230, 2013.
60. Egner, IM, Bruusgaard, JC, and Gundersen, K. Satellite cell depletion prevents fiber hypertrophy in skeletal muscle. *Development* 143: 2898-2906, 2016.
61. Ehrnborg, C, and Rosen, T. Physiological and pharmacological basis for the ergogenic effects of growth hormone in elite sports. *Asian J. Androl.* 10: 373-383, 2008.
62. Febbraio, MA, and Pedersen, BK. Contraction-induced myokine production and release: is skeletal muscle an endocrine organ? *Exerc. Sport Sci. Rev.* 33: 114-119, 2005.
63. Fernandez-Gonzalo, R, Lundberg, TR, and Tesch, PA. Acute molecular responses in untrained and trained muscle subjected to aerobic and resistance exercise training versus resistance training alone. *Acta Physiol. (Oxf)* 209: 283-294, 2013.

64. Figueiredo, VC, and McCarthy, JJ. Regulation of Ribosome Biogenesis in Skeletal Muscle Hypertrophy. *Physiology (Bethesda)* 34: 30-42, 2019.
65. Fluckey, JD, Vary, TC, Jefferson, LS, and Farrell, PA. Augmented insulin action on rates of protein synthesis after resistance exercise in rats. *Am. J. Physiol.* 270: E313-9, 1996.
66. Fornaro, M, Hinken, AC, Needle, S, Hu, E, Trendelenburg, AU, Mayer, A, Rosenstiel, A, Chang, C, Meier, V, Billin, AN, Becherer, JD, Brace, AD, Evans, WJ, Glass, DJ, and Russell, AJ. Mechano-growth factor peptide, the COOH terminus of unprocessed insulin-like growth factor 1, has no apparent effect on myoblasts or primary muscle stem cells. *Am. J. Physiol. Endocrinol. Metab.* 306: E150-6, 2014.
67. Fry, CS, Kirby, TJ, Kosmac, K, McCarthy, JJ, and Peterson, CA. Myogenic Progenitor Cells Control Extracellular Matrix Production by Fibroblasts during Skeletal Muscle Hypertrophy. *Cell. Stem Cell.* 20: 56-69, 2017.
68. Fujita, S, Abe, T, Drummond, MJ, Cadenas, JG, Dreyer, HC, Sato, Y, Volpi, E, and Rasmussen, BB. Blood flow restriction during low-intensity resistance exercise increases S6K1 phosphorylation and muscle protein synthesis. *J. Appl. Physiol.* 103: 903-910, 2007.
69. Gelfand, RA, and Barrett, EJ. Effect of physiologic hyperinsulinemia on skeletal muscle protein synthesis and breakdown in man. *J. Clin. Invest.* 80: 1-6, 1987.
70. Glass, DJ. PI3 kinase regulation of skeletal muscle hypertrophy and atrophy. *Curr. Top. Microbiol. Immunol.* 346: 267-278, 2010.
71. Goldspink, G. Mechanical signals, IGF-I gene splicing, and muscle adaptation. *Physiology (Bethesda)* 20: 232-238, 2005.
72. Goldspink, G. Impairment of IGF-I gene splicing and MGF expression associated with muscle wasting. *Int. J. Biochem. Cell Biol.* 38: 481-489, 2006.
73. Goodman, CA, Mayhew, DL, and Hornberger, TA. Recent progress toward understanding the molecular mechanisms that regulate skeletal muscle mass. *Cell. Signal.* 23: 1896-1906, 2011.
74. Gordon, SE, Kraemer, WJ, Vos, NH, Lynch, JM, and Knuttgen, HG. Effect of acid-base balance on the growth hormone response to acute high-intensity cycle exercise. *J. Appl. Physiol.* 76: 821-829, 1994.
75. Goto, K, Ishii, N, Kizuka, T, and Takamatsu, K. The impact of metabolic stress on hormonal responses and muscular adaptations. *Med. Sci. Sports Exerc.* 37: 955-963, 2005.
76. Gotshalk, LA, Loebel, CC, Nindl, BC, Putukian, M, Sebastianelli, WJ, Newton, RU, Häkkinen, K, and Kraemer, WJ. Hormonal responses of multiset versus single-set heavy-resistance exercise protocols. *Can. J. Appl. Physiol.* 22: 244-255, 1997.
77. Gundersen, K. Excitation-transcription coupling in skeletal muscle: the molecular pathways of exercise. *Biol. Rev. Camb. Philos. Soc.* 86: 564-600, 2011.
78. Haddad, F, and Adams, GR. Inhibition of MAP/ERK kinase prevents IGF-I-induced hypertrophy in rat muscles. *J. Appl. Physiol.* 96: 203-210, 2004.
79. Häkkinen, K, and Pakarinen, A. Acute hormonal responses to two different fatiguing heavy-resistance protocols in male athletes. *J. Appl. Physiol.* 74: 882-887, 1993.
80. Häkkinen, K, Alen, M, Kallinen, M, Newton, RU, and Kraemer, WJ. Neuromuscular adaptation during prolonged strength training, detraining and re-strength-training in middle-aged and elderly people. *Eur. J. Appl. Physiol.* 83: 51-62, 2000.
81. Häkkinen, K, Pakarinen, A, Kraemer, WJ, Newton, RU, and Alen, M. Basal concentrations and acute responses of serum hormones and strength development during heavy resistance training in middle-aged and elderly men and women. *J. Gerontol. A Biol. Sci. Med. Sci.* 55: B95-105, 2000.
82. Häkkinen, K, Pakarinen, A, Kraemer, WJ, Häkkinen, A, Valkeinen, H, and Alen, M. Selective muscle hypertrophy, changes in EMG and force, and serum hormones during strength training in older women. *J. Appl. Physiol.* 91: 569-580, 2001.
83. Hameed, M, Lange, KH, Andersen, JL, Schjerling, P, Kjaer, M, Harridge, SD, and Goldspink, G. The effect of recombinant human growth hormone and resistance training on IGF-I mRNA expression in the muscles of elderly men. *J. Physiol.* 555: 231-240, 2004.
84. Hand, BD, Kostek, MC, Ferrell, RE, Delmonico, MJ, Douglass, LW, Roth, SM, Hagberg, JM, and Hurley, BF. Influence of promoter region variants of insulin-like growth factor pathway genes on the strength-training response of muscle phenotypes in older adults. *J. Appl. Physiol.* 103: 1678-1687, 2007.
85. Hansen, S, Kvorning, T, Kjaer, M, and Sjogaard, G. The effect of short-term strength training on human skeletal muscle: the importance of physiologically elevated hormone levels. *Scand. J. Med. Sci. Sports* 11: 347-354, 2001.
86. Hanssen, KE, Kvamme, NH, Nilsen, TS, Ronnestad, B, Ambjornsen, IK, Norheim, F, Kadi, F, Hallen, J, Drevon, CA, and Raastad, T. The effect of strength training volume on satellite cells, myogenic regulatory factors, and growth factors. *Scand. J. Med. Sci. Sports* 23: 728-739, 2013.
87. Harber, MP, Konopka, AR, Douglass, MD, Minchev, K, Kaminisky, LA, Trappe, TA, and Trappe, S. Aerobic exercise training improves whole muscle and single myofiber size and function in older women. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 297: R1452-9, 2009.
88. Harridge, SD. Plasticity of human skeletal muscle: gene expression to in vivo function. *Exp. Physiol.* 92: 783-797, 2007.
89. Haun, CT, Vann, CG, Osburn, SC, Mumford, PW, Roberson, PA, Romero, MA, Fox, CD, Johnson, CA, Parry, HA, Kavazis, AN, Moon, JR, Badisa, VLD, Mwashote, BM, Ibeanusi, V, Young, KC, and Roberts, MD. Muscle fiber hypertrophy in response to 6 weeks of high-volume resistance training in trained young men is largely attributed to sarcoplasmic hypertrophy. *PLoS One* 14: e0215267, 2019.
90. Henneman, E, Somjen, G, and Carpenter, DO. Functional Significance of Cell Size in Spinal Motoneurons. *J. Neurophysiol.* 28: 560-580, 1965.
91. Heslin, MJ, Newman, E, Wolf, RF, Pisters, PW, and Brennan, MF. Effect of hyperinsulinemia on whole body and skeletal muscle leucine carbon kinetics in humans. *Am. J. Physiol.* 262: E911-8, 1992.
92. Hill, M, Wernig, A, and Goldspink, G. Muscle satellite (stem) cell activation during local tissue injury and repair. *J. Anat.* 203: 89-99, 2003.
93. Hooper, DR, Kraemer, WJ, Focht, BC, Volek, JS, DuPont, WH, Caldwell, LK, and Maresch, CM. Endocrinological Roles for Testosterone in Resistance Exercise Responses and Adaptations. *Sports Med.* 47: 1709-1720, 2017.
94. Hornberger, TA, and Esser, KA. Mechanotransduction and the regulation of protein synthesis in skeletal muscle. *Proc. Nutr. Soc.* 63: 331-335, 2004.
95. Houtman, CJ, Stegeman, DE, Van Dijk, JP, and Zwarts, MJ. Changes in muscle fiber conduction velocity indicate recruit-

- ment of distinct motor unit populations. *J. Appl. Physiol.* 95: 1045-1054, 2003.
96. Huey, KA. Potential Roles of Vascular Endothelial Growth Factor During Skeletal Muscle Hypertrophy. *Exerc. Sport Sci. Rev.* 46: 195-202, 2018.
 97. Huxley, AF. The origin of force in skeletal muscle. *Ciba Found. Symp.* (31): 271-290, 1975.
 98. Iida, K, Itoh, E, Kim, DS, del Rincon, JP, Coschigano, KT, Kopchick, JJ, and Thorner, MO. Muscle mechano growth factor is preferentially induced by growth hormone in growth hormone-deficient lit/lit mice. *J. Physiol.* 560: 341-349, 2004.
 99. Imanaka, M, Iida, K, Murawaki, A, Nishizawa, H, Fukuoka, H, Takeno, R, Takahashi, Y, Okimura, Y, Kaji, H, and Chihara, K. Growth hormone stimulates mechano growth factor expression and activates myoblast transformation in C2C12 cells. *Kobe J. Med. Sci.* 54: E46-54, 2008.
 100. Kadi, F, Schjerling, P, Andersen, LL, Charifi, N, Madsen, JL, Christensen, LR, and Andersen, JL. The effects of heavy resistance training and detraining on satellite cells in human skeletal muscles. *J. Physiol.* 558: 1005-1012, 2004.
 101. Kamen, G, and Knight, CA. Training-related adaptations in motor unit discharge rate in young and older adults. *J. Gerontol. A Biol. Sci. Med. Sci.* 59: 1334-1338, 2004.
 102. Kami, K, and Senba, E. Localization of leukemia inhibitory factor and interleukin-6 messenger ribonucleic acids in regenerating rat skeletal muscle. *Muscle Nerve* 21: 819-822, 1998.
 103. Kelley, G. Mechanical overload and skeletal muscle fiber hyperplasia: a meta-analysis. *J. Appl. Physiol.* 81: 1584-1588, 1996.
 104. Kettelhut, IC, Wing, SS, and Goldberg, AL. Endocrine regulation of protein breakdown in skeletal muscle. *Diabetes Metab. Rev.* 4: 751-772, 1988.
 105. Kidgell, DJ, Sale, MV, and Semmler, JG. Motor unit synchronization measured by cross-correlation is not influenced by short-term strength training of a hand muscle. *Exp. Brain Res.* 175: 745-753, 2006.
 106. Kim, H, Barton, E, Muja, N, Yakar, S, Pennisi, P, and Leroith, D. Intact insulin and insulin-like growth factor-I receptor signaling is required for growth hormone effects on skeletal muscle growth and function in vivo. *Endocrinology* 146: 1772-1779, 2005.
 107. Kim, JS, Cross, JM, and Bamman, MM. Impact of resistance loading on myostatin expression and cell cycle regulation in young and older men and women. *Am. J. Physiol. Endocrinol. Metab.* 288: E1110-9, 2005.
 108. Kim, JS, Petrella, JK, Cross, JM, and Bamman, MM. Load-mediated downregulation of myostatin mRNA is not sufficient to promote myofiber hypertrophy in humans: a cluster analysis. *J. Appl. Physiol.* (1985) 103: 1488-1495, 2007.
 109. Klover, P, and Hennighausen, L. Postnatal body growth is dependent on the transcription factors signal transducers and activators of transcription 5a/b in muscle: a role for autocrine/paracrine insulin-like growth factor I. *Endocrinology* 148: 1489-1497, 2007.
 110. Knight, CA, and Kamen, G. Adaptations in muscular activation of the knee extensor muscles with strength training in young and older adults. *J. Electromyogr. Kinesiol.* 11: 405-412, 2001.
 111. Kosek, DJ, Kim, JS, Petrella, JK, Cross, JM, and Bamman, MM. Efficacy of 3 days/wk resistance training on myofiber hypertrophy and myogenic mechanisms in young vs. older adults. *J. Appl. Physiol.* 101: 531-544, 2006.
 112. Kraemer, WJ, Marchitelli, L, Gordon, SE, Harman, E, Dziados, JE, Mello, R, Frykman, P, McCurry, D, and Fleck, SJ. Hormonal and growth factor responses to heavy resistance exercise protocols. *J. Appl. Physiol.* 69: 1442-1450, 1990.
 113. Kraemer, WJ, Gordon, SE, Fleck, SJ, Marchitelli, LJ, Mello, R, Dziados, JE, Friedl, K, Harman, E, Maresh, C, and Fry, AC. Endogenous anabolic hormonal and growth factor responses to heavy resistance exercise in males and females. *Int. J. Sports Med.* 12: 228-235, 1991.
 114. Kraemer, WJ, Aguilera, BA, Terada, M, Newton, RU, Lynch, JM, Rosendaal, G, McBride, JM, Gordon, SE, and Häkkinen, K. Responses of IGF-I to endogenous increases in growth hormone after heavy-resistance exercise. *J. Appl. Physiol.* 79: 1310-1315, 1995.
 115. Kraemer, WJ, and Ratamess, NA. Hormonal responses and adaptations to resistance exercise and training. *Sports Med.* 35: 339-361, 2005.
 116. Kvorning, T, Andersen, M, Brixen, K, and Madsen, K. Suppression of endogenous testosterone production attenuates the response to strength training: a randomized, placebo-controlled, and blinded intervention study. *Am. J. Physiol. Endocrinol. Metab.* 291: E1325-32, 2006.
 117. Kvorning, T, Andersen, M, Brixen, K, Schjerling, P, Suetta, C, and Madsen, K. Suppression of testosterone does not blunt mRNA expression of myoD, myogenin, IGF, myostatin or androgen receptor post strength training in humans. *J. Physiol.* 578: 579-593, 2007.
 118. Lange, KH, Andersen, JL, Beyer, N, Isaksson, F, Larsson, B, Rasmussen, MH, Juul, A, Bulow, J, and Kjaer, M. GH administration changes myosin heavy chain isoforms in skeletal muscle but does not augment muscle strength or hypertrophy, either alone or combined with resistance exercise training in healthy elderly men. *J. Clin. Endocrinol. Metab.* 87: 513-523, 2002.
 119. Loebel, C, and Kraemer, W. A brief review: Testosterone and resistance exercise in men. *J. Strength and Conditioning Research* 12: 57-63, 1998.
 120. Loenneke, JP, Wilson, GJ, and Wilson, JM. A mechanistic approach to blood flow occlusion. *Int. J. Sports Med.* 31: 1-4, 2010.
 121. Loenneke, JP, Fahs, CA, Wilson, JM, and Bembien, MG. Blood flow restriction: the metabolite/volume threshold theory. *Med. Hypotheses* 77: 748-752, 2011.
 122. Lupu, F, Terwilliger, JD, Lee, K, Segre, GV, and Efstratiadis, A. Roles of growth hormone and insulin-like growth factor 1 in mouse postnatal growth. *Dev. Biol.* 229: 141-162, 2001.
 123. Lynn, R, and Morgan, DL. Decline running produces more sarcomeres in rat vastus intermedius muscle fibers than does incline running. *J. Appl. Physiol.* 77: 1439-1444, 1994.
 124. MacDougall, JD, Ward, GR, Sale, DG, and Sutton, JR. Biochemical adaptation of human skeletal muscle to heavy resistance training and immobilization. *J. Appl. Physiol.* 43: 700-703, 1977.
 125. MacDougall, JD, Sale, DG, Elder, GC, and Sutton, JR. Muscle ultrastructural characteristics of elite powerlifters and bodybuilders. *Eur. J. Appl. Physiol. Occup. Physiol.* 48: 117-126, 1982.
 126. MacDougall, JD, Sale, DG, Alway, SE, and Sutton, JR. Muscle fiber number in biceps brachii in bodybuilders and control subjects. *J. Appl. Physiol.* 57: 1399-1403, 1984.
 127. Madarame, H, Neya, M, Ochi, E, Nakazato, K, Sato, Y, and Ishii, N. Cross-transfer effects of resistance training with blood flow restriction. *Med. Sci. Sports Exerc.* 40: 258-263, 2008.

128. Matheny, RW, Merritt, E, Zannikos, SV, Farrar, RP, and Adamo, ML. Serum IGF-I-deficiency does not prevent compensatory skeletal muscle hypertrophy in resistance exercise. *Exp. Biol. Med.* (Maywood) 234: 164-170, 2009.
129. Matsakas, A, Foster, K, Otto, A, Macharia, R, Elashry, MI, Feist, S, Graham, I, Foster, H, Yaworsky, P, Walsh, F, Dickson, G, and Patel, K. Molecular, cellular and physiological investigation of myostatin propeptide-mediated muscle growth in adult mice. *Neuromuscul. Disord.* 19: 489-499, 2009.
130. Mayhew, DL, Hornberger, TA, Lincoln, HC, and Bamman, MM. Eukaryotic initiation factor 2B epsilon induces cap-dependent translation and skeletal muscle hypertrophy. *J. Physiol.* 589: 3023-3037, 2011.
131. McCall, GE, Byrnes, WC, Fleck, SJ, Dickinson, A, and Kraemer, WJ. Acute and chronic hormonal responses to resistance training designed to promote muscle hypertrophy. *Can. J. Appl. Physiol.* 24: 96-107, 1999.
132. McCarthy, JJ, and Esser, KA. Counterpoint: Satellite cell addition is not obligatory for skeletal muscle hypertrophy. *J Appl Physiol* 103: 1100-1102, 2007.
133. McCarthy, JJ, Murach, KA. Anabolic and catabolic signaling pathways that regulate skeletal muscle mass. In: *Nutrition and Enhanced Sports Performance*. Anonymous Cambridge, MA: Academic Press, 2019. pp. 275-290.
134. McCaulley, GO, McBride, JM, Cormie, P, Hudson, MB, Nuzzo, JL, Quindry, JC, and Travis Triplett, N. Acute hormonal and neuromuscular responses to hypertrophy, strength and power type resistance exercise. *Eur. J. Appl. Physiol.* 105: 695-704, 2009.
135. McCroskery, S, Thomas, M, Maxwell, L, Sharma, M, and Kambadur, R. Myostatin negatively regulates satellite cell activation and self-renewal. *J. Cell Biol.* 162: 1135-1147, 2003.
136. McPherron, AC, Lawler, AM, and Lee, SJ. Regulation of skeletal muscle mass in mice by a new TGF-beta superfamily member. *Nature* 387: 83-90, 1997.
137. Michaud, M, Balardy, L, Moulis, G, Gaudin, C, Peyrot, C, Vellas, B, Cesari, M, and Nourhashemi, F. Proinflammatory cytokines, aging, and age-related diseases. *J. Am. Med. Dir. Assoc.* 14: 877-882, 2013.
138. Milewska, M, Domoradzki, T, Majewska, A, Blaszczyk, M, Gajewska, M, Hulanicka, M, Ciecierska, A, and Grzelkowska-Kowalczyk, K. Interleukin-8 enhances myocilin expression, Akt-FoxO3 signaling and myogenic differentiation in rat skeletal muscle cells. *J. Cell. Physiol.* 234: 19675-19690, 2019.
139. Minderis, P, Kilikevicius, A, Baltusnikas, J, Alhindi, Y, Venckunas, T, Bunger, L, Lionikas, A, and Ratkevicius, A. Myostatin dysfunction is associated with reduction in overload induced hypertrophy of soleus muscle in mice. *Scand. J. Med. Sci. Sports* 26: 894-901, 2016.
140. Mitchell, CJ, Churchward-Venne, TA, Bellamy, L, Parise, G, Baker, SK, and Phillips, SM. Muscular and systemic correlates of resistance training-induced muscle hypertrophy. *PLoS One* 8: e78636, 2013.
141. Moritani, T, and deVries, HA. Neural factors versus hypertrophy in the time course of muscle strength gain. *Am. J. Phys. Med.* 58: 115-130, 1979.
142. Morton, RW, Sato, K, Gallaughier, MPB, Oikawa, SY, McNicholas, PD, Fujita, S, and Phillips, SM. Muscle Androgen Receptor Content but Not Systemic Hormones Is Associated With Resistance Training-Induced Skeletal Muscle Hypertrophy in Healthy, Young Men. *Front. Physiol.* 9: 1373, 2018.
143. Mosher, DS, Quignon, P, Bustamante, CD, Sutter, NB, Mellersh, CS, Parker, HG, and Ostrander, EA. A mutation in the myostatin gene increases muscle mass and enhances racing performance in heterozygote dogs. *PLoS Genet.* 3: e79, 2007.
144. Moss, FP, and Leblond, CP. Satellite cells as the source of nuclei in muscles of growing rats. *Anat. Rec.* 170: 421-435, 1971.
145. Mouser, JG, Loprinzi, PD, and Loenneke, JP. The association between physiologic testosterone levels, lean mass, and fat mass in a nationally representative sample of men in the United States. *Steroids* 115: 62-66, 2016.
146. Murach, KA, Fry, CS, Kirby, TJ, Jackson, JR, Lee, JD, White, SH, Dupont-Versteegden, EE, McCarthy, JJ, and Peterson, CA. Starving or Supporting Role? Satellite Cells and Skeletal Muscle Fiber Size Regulation. *Physiology (Bethesda)* 33: 26-38, 2018.
147. Murach, KA, Dungan, CM, Peterson, CA, and McCarthy, JJ. Muscle Fiber Splitting Is a Physiological Response to Extreme Loading in Animals. *Exerc. Sport Sci. Rev.* 47: 108-115, 2019.
148. Nader, GA, von Walden, F, Liu, C, Lindvall, J, Gutmann, L, Pistilli, EE, and Gordon, PM. Resistance exercise training modulates acute gene expression during human skeletal muscle hypertrophy. *J. Appl. Physiol.* (1985) 116: 693-702, 2014.
149. Nakada, S, Ogasawara, R, Kawada, S, Maekawa, T, and Ishii, N. Correlation between Ribosome Biogenesis and the Magnitude of Hypertrophy in Overloaded Skeletal Muscle. *PLoS One* 11: e0147284, 2016.
150. Narici, MV, Roi, GS, Landoni, L, Minetti, AE, and Cerretelli, P. Changes in force, cross-sectional area and neural activation during strength training and detraining of the human quadriceps. *Eur. J. Appl. Physiol. Occup. Physiol.* 59: 310-319, 1989.
151. Nederveen, JP, Joannis, S, Snijders, T, and Parise, G. The influence and delivery of cytokines and their mediating effect on muscle satellite cells. *Current Stem Cell Reports* 3: 192-201, 2017.
152. Nielsen, AR, Mounier, R, Plomgaard, P, Mortensen, OH, Penkowa, M, Speerschnieder, T, Pilegaard, H, and Pedersen, BK. Expression of interleukin-15 in human skeletal muscle effect of exercise and muscle fibre type composition. *J. Physiol.* 584: 305-312, 2007.
153. Nielsen, AR, and Pedersen, BK. The biological roles of exercise-induced cytokines: IL-6, IL-8, and IL-15. *Appl. Physiol. Nutr. Metab.* 32: 833-839, 2007.
154. Nindl, BC, Hymer, WC, Deaver, DR, and Kraemer, WJ. Growth hormone pulsatility profile characteristics following acute heavy resistance exercise. *J. Appl. Physiol.* 91: 163-172, 2001.
155. Norheim, KL, Cullum, CK, Andersen, JL, Kjaer, M, and Karlsen, A. Inflammation Relates to Resistance Training-induced Hypertrophy in Elderly Patients. *Med. Sci. Sports Exerc.* 49: 1079-1085, 2017.
156. Ochi, E, Ishii, N, and Nakazato, K. Time course change of IGF1/Akt/mTOR/p70s6k pathway activation in rat gastrocnemius muscle during repeated bouts of eccentric exercise. *JSSM* 9: 170-175, 2010.
157. O'Connor, RS, and Pavlath, GK. Point:Counterpoint: Satellite cell addition is/is not obligatory for skeletal muscle hypertrophy. *J. Appl. Physiol.* 103: 1099-1100, 2007.
158. Ogborn, D, and Schoenfeld, BJ. The role of fiber types in muscle hypertrophy: Implications for loading strategies. *Strength Cond J* 36: 20-25, 2014.
159. Olsson, KE, and Saltin, B. Variation in total body water with muscle glycogen changes in man. *Acta Physiol. Scand.* 80: 11-18, 1970.

160. O'Neil, TK, Duffy, LR, Frey, JW, and Hornberger, TA. The role of phosphoinositide 3-kinase and phosphatidic acid in the regulation of mammalian target of rapamycin following eccentric contractions. *J. Physiol.* 587: 3691-3701, 2009.
161. Paul, AC, and Rosenthal, N. Different modes of hypertrophy in skeletal muscle fibers. *J. Cell Biol.* 156: 751-760, 2002.
162. Peake, JM, Della Gatta, P, Suzuki, K, and Nieman, DC. Cytokine expression and secretion by skeletal muscle cells: regulatory mechanisms and exercise effects. *Exerc. Immunol. Rev.* 21: 8-25, 2015.
163. Pedersen, BK. Muscles and their myokines. *J. Exp. Biol.* 214: 337-346, 2011.
164. Pedersen, BK, and Febbraio, MA. Muscles, exercise and obesity: skeletal muscle as a secretory organ. *Nat. Rev. Endocrinol.* 8: 457-465, 2012.
165. Perez-Lopez, A, McKendry, J, Martin-Rincon, M, Morales-Alamo, D, Perez-Kohler, B, Valades, D, Bujan, J, Calbet, JAL, and Breen, L. Skeletal muscle IL-15/IL-15R α and myofibrillar protein synthesis after resistance exercise. *Scand. J. Med. Sci. Sports* 28: 116-125, 2018.
166. Petrella, JK, Kim, JS, Cross, JM, Kosek, DJ, and Bamman, MM. Efficacy of myonuclear addition may explain differential myofiber growth among resistance-trained young and older men and women. *Am. J. Physiol. Endocrinol. Metab.* 291: E937-46, 2006.
167. Petrella, JK, Kim, J, Mayhew, DL, Cross, JM, and Bamman, MM. Potent myofiber hypertrophy during resistance training in humans is associated with satellite cell-mediated myonuclear addition: a cluster analysis. *J. Appl. Physiol.* 104: 1736-1742, 2008.
168. Phillips, SM, Tipton, KD, Aarsland, A, Wolf, SE, and Wolfe, RR. Mixed muscle protein synthesis and breakdown after resistance exercise in humans. *Am. J. Physiol.* 273: E99-107, 1997.
169. Phillips, SM. Physiologic and molecular bases of muscle hypertrophy and atrophy: impact of resistance exercise on human skeletal muscle (protein and exercise dose effects). *Appl. Physiol. Nutr. Metab.* 34: 403-410, 2009.
170. Pierce, JR, Clark, BC, Ploutz-Snyder, LL, and Kanaley, JA. Growth hormone and muscle function responses to skeletal muscle ischemia. *J. Appl. Physiol.* 101: 1588-1595, 2006.
171. Pillon, NJ, Bilan, PJ, Fink, LN, and Klip, A. Cross-talk between skeletal muscle and immune cells: muscle-derived mediators and metabolic implications. *Am. J. Physiol. Endocrinol. Metab.* 304: E453-65, 2013.
172. Pistilli, EE, and Quinn, LS. From anabolic to oxidative: reconsidering the roles of IL-15 and IL-15R α in skeletal muscle. *Exerc. Sport Sci. Rev.* 41: 100-106, 2013.
173. Ploutz, LL, Tesch, PA, Biro, RL, and Dudley, GA. Effect of resistance training on muscle use during exercise. *J. Appl. Physiol.* (1985) 76: 1675-1681, 1994.
174. Pucci, AR, Griffin, L, and Cafarelli, E. Maximal motor unit firing rates during isometric resistance training in men. *Exp. Physiol.* 91: 171-178, 2006.
175. Qiao, C, Li, J, Jiang, J, Zhu, X, Wang, B, Li, J, and Xiao, X. Myostatin propeptide gene delivery by adeno-associated virus serotype 8 vectors enhances muscle growth and ameliorates dystrophic phenotypes in mdx mice. *Hum. Gene Ther.* 19: 241-254, 2008.
176. Quinn, LS, Anderson, BG, Conner, JD, Pistilli, EE, and Wolden-Hanson, T. Overexpression of interleukin-15 in mice promotes resistance to diet-induced obesity, increased insulin sensitivity, and markers of oxidative skeletal muscle metabolism. *Int J Interferon Cytokine Mediator Res* 3: 29-42, 2011.
177. Quinn, LS. Interleukin-15: a muscle-derived cytokine regulating fat-to-lean body composition. *J. Anim. Sci.* 86: E75-83, 2008.
178. Randrianarison-Huetz, V, Papaefthymiou, A, Herledan, G, Novello, C, Faradova, U, Collard, L, Pincini, A, Schol, E, Decaux, JF, Maire, P, Vassilopoulos, S, and Sotiropoulos, A. Srf controls satellite cell fusion through the maintenance of actin architecture. *J. Cell Biol.* 217: 685-700, 2018.
179. Raue, U, Trappe, TA, Estrem, ST, Qian, HR, Helvering, LM, Smith, RC, and Trappe, S. Transcriptome signature of resistance exercise adaptations: mixed muscle and fiber type specific profiles in young and old adults. *J. Appl. Physiol.* (1985) 112: 1625-1636, 2012.
180. Raught, B, and Gingras, AC. eIF4E activity is regulated at multiple levels. *Int. J. Biochem. Cell Biol.* 31: 43-57, 1999.
181. Rayagiri, SS, Ranaldi, D, Raven, A, Mohamad Azhar, NIF, Lefebvre, O, Zammit, PS, and Borycki, AG. Basal lamina remodeling at the skeletal muscle stem cell niche mediates stem cell self-renewal. *Nat. Commun.* 9: 1075-018-03425-3, 2018.
182. Reeves, GV, Kraemer, RR, Hollander, DB, Clavier, J, Thomas, C, Francois, M, and Castracane, VD. Comparison of hormone responses following light resistance exercise with partial vascular occlusion and moderately difficult resistance exercise without occlusion. *J. Appl. Physiol.* 101: 1616-1622, 2006.
183. Reidy, PT, Borack, MS, Markofski, MM, Dickinson, JM, Fry, CS, Deer, RR, Volpi, E, and Rasmussen, BB. Post-absorptive muscle protein turnover affects resistance training hypertrophy. *Eur. J. Appl. Physiol.* 117: 853-866, 2017.
184. Reihmane, D, and Dela, F. Interleukin-6: possible biological roles during exercise. *Eur. J. Sport. Sci.* 14: 242-250, 2014.
185. Ribeiro, AS, Avelar, A, Schoenfeld, BJ, Ritti Dias, RM, Altimari, LR, and Cyrino, ES. Resistance training promotes increase in intracellular hydration in men and women. *Eur. J. Sport. Sci.* 14: 578-585, 2014.
186. Riechman, SE, Balasekaran, G, Roth, SM, and Ferrell, RE. Association of interleukin-15 protein and interleukin-15 receptor genetic variation with resistance exercise training responses. *J. Appl. Physiol.* 97: 2214-2219, 2004.
187. Rieu, I, Magne, H, Savary-Auzeloux, I, Averous, J, Bos, C, Peyron, MA, Combaret, L, and Dardevet, D. Reduction of low grade inflammation restores blunting of postprandial muscle anabolism and limits sarcopenia in old rats. *J. Physiol.* 587: 5483-5492, 2009.
188. Rigamonti, AE, Locatelli, L, Cella, SG, Bonomo, SM, Giunta, M, Molinari, F, Sartorio, A, and Muller, EE. Muscle expressions of MGF, IGF-IEa, and myostatin in intact and hypophysectomized rats: effects of rhGH and testosterone alone or combined. *Horm. Metab. Res.* 41: 23-29, 2009.
189. Rommel, C, Bodine, SC, Clarke, BA, Rossman, R, Nunez, L, Stitt, TN, Yancopoulos, GD, and Glass, DJ. Mediation of IGF-1-induced skeletal myotube hypertrophy by PI(3)K/Akt/mTOR and PI(3)K/Akt/GSK3 pathways. *Nat. Cell Biol.* 3: 1009-1013, 2001.
190. Ronnestad, BR, Nygaard, H, and Raastad, T. Physiological elevation of endogenous hormones results in superior strength training adaptation. *Eur. J. Appl. Physiol.* 111: 2249-2259, 2011.
191. Rossetti, ML, Steiner, JL, and Gordon, BS. Androgen-mediated regulation of skeletal muscle protein balance. *Mol. Cell. Endocrinol.* 447: 35-44, 2017.
192. Rubin, MR, Kraemer, WJ, Maresch, CM, Volek, JS, Ratamess, NA, Vanheest, JL, Silvestre, R, French, DN, Sharman, MJ, Judelson, DA, Gomez, AL, Vescovi, JD, and Hymer, WC. High-affinity growth hormone binding protein and acute heavy resistance exercise. *Med. Sci. Sports Exerc.* 37: 395-403, 2005.

193. Sabourin, LA, and Rudnicki, MA. The molecular regulation of myogenesis. *Clin. Genet.* 57: 16-25, 2000.
194. Saclier, M, Yacoub-Youssef, H, Mackey, AL, Arnold, L, Ardjoune, H, Magnan, M, Sailhan, F, Chelly, J, Pavlath, GK, Mounier, R, Kjaer, M, and Chazaud, B. Differentially activated macrophages orchestrate myogenic precursor cell fate during human skeletal muscle regeneration. *Stem Cells* 31: 384-396, 2013.
195. Sahlin, K, Soderlund, K, Tonkonogi, M, and Hirakoba, K. Phosphocreatine content in single fibers of human muscle after sustained submaximal exercise. *Am. J. Physiol.* 273: C172-8, 1997.
196. Sale, DG. Neural adaptation to resistance training. *Med. Sci. Sports Exerc.* 20: S135-45, 1988.
197. Sandri, M. Signaling in muscle atrophy and hypertrophy. *Physiology (Bethesda)* 23: 160-170, 2008.
198. Schoenfeld, BJ. The mechanisms of muscle hypertrophy and their application to resistance training. *J. Strength Cond Res.* 24: 2857-2872, 2010.
199. Schoenfeld, BJ. Postexercise hypertrophic adaptations: a reexamination of the hormone hypothesis and its applicability to resistance training program design. *J. Strength Cond Res.* 27: 1720-1730, 2013.
200. Schuelke, M, Wagner, KR, Stolz, LE, Hubner, C, Riebel, T, Komen, W, Braun, T, Tobin, JE, and Lee, SJ. Myostatin mutation associated with gross muscle hypertrophy in a child. *N. Engl. J. Med.* 350: 2682-2688, 2004.
201. Schuenke, MD, Herman, JR, Gliders, RM, Hagerman, FC, Hikida, RS, Rana, SR, Ragg, KE, and Staron, RS. Early-phase muscular adaptations in response to slow-speed versus traditional resistance-training regimens. *Eur. J. Appl. Physiol.* 112: 3585-3595, 2012.
202. Schwartz, LM. Skeletal Muscles Do Not Undergo Apoptosis During Either Atrophy or Programmed Cell Death-Revisiting the Myonuclear Domain Hypothesis. *Front. Physiol.* 9: 1887, 2019.
203. Sculthorpe, N, Solomon, AM, Sinanan, AC, Bouloux, PM, Grace, F, and Lewis, MP. Androgens Affect Myogenesis In Vitro and Increase Local IGF-1 Expression. *Med. Sci. Sports Exerc.* 44: 610-615, 2012.
204. Semmler, JG, and Nordstrom, MA. Motor unit discharge and force tremor in skill- and strength-trained individuals. *Exp. Brain Res.* 119: 27-38, 1998.
205. Semsarian, C, Wu, MJ, Ju, YK, Marciniak, T, Yeoh, T, Allen, DG, Harvey, RP, and Graham, RM. Skeletal muscle hypertrophy is mediated by a Ca²⁺-dependent calcineurin signalling pathway. *Nature* 400: 576-581, 1999.
206. Serrano, AL, Baeza-Raja, B, Perdiguero, E, Jardi, M, and Munoz-Canoves, P. Interleukin-6 is an essential regulator of satellite cell-mediated skeletal muscle hypertrophy. *Cell. Metab.* 7: 33-44, 2008.
207. Seynnes, OR, de Boer, M, and Narici, MV. Early skeletal muscle hypertrophy and architectural changes in response to high-intensity resistance training. *J. Appl. Physiol.* 102: 368-373, 2007.
208. Shan, T, Xu, Z, Wu, W, Liu, J, and Wang, Y. Roles of Notch1 Signaling in Regulating Satellite Cell Fates Choices and Postnatal Skeletal Myogenesis. *J. Cell. Physiol.* 232: 2964-2967, 2017.
209. Siff, M. Supertraining. Denver, CO; Supertraining Institute, 2009.
210. Sinha-Hikim, I, Cornford, M, Gaytan, H, Lee, ML, and Bhasin, S. Effects of testosterone supplementation on skeletal muscle fiber hypertrophy and satellite cells in community-dwelling older men. *J. Clin. Endocrinol. Metab.* 91: 3024-3033, 2006.
211. Smiliotis, I, Piliandis, T, Karamouzis, M, and Tokmakidis, SP. Hormonal responses after various resistance exercise protocols. *Med. Sci. Sports Exerc.* 35: 644-654, 2003.
212. Solomon, AM, and Bouloux, PM. Modifying muscle mass - the endocrine perspective. *J. Endocrinol.* 191: 349-360, 2006.
213. Sotiropoulos, A, Ohanna, M, Kedzia, C, Menon, RK, Kopchick, JJ, Kelly, PA, and Pende, M. Growth hormone promotes skeletal muscle cell fusion independent of insulin-like growth factor 1 up-regulation. *Proc. Natl. Acad. Sci. U. S. A.* 103: 7315-7320, 2006.
214. Spangenburg, EE, Le Roith, D, Ward, CW, and Bodine, SC. A functional insulin-like growth factor receptor is not necessary for load-induced skeletal muscle hypertrophy. *J. Physiol.* 586: 283-291, 2008.
215. Spangenburg, EE. Changes in muscle mass with mechanical load: possible cellular mechanisms. *Appl. Physiol. Nutr. Metab.* 34: 328-335, 2009.
216. Spiering, BA, Kraemer, WJ, Vingren, JL, Ratamess, NA, Anderson, JM, Armstrong, LE, Nindl, BC, Volek, JS, Häkkinen, K, and Maresh, CM. Elevated endogenous testosterone concentrations potentiate muscle androgen receptor responses to resistance exercise. *J. Steroid Biochem. Mol. Biol.* 114: 195-199, 2009.
217. Staron, RS, Leonardi, MJ, Karapondo, DL, Malicky, ES, Falkel, JE, Hagerman, FC, and Hikida, RS. Strength and skeletal muscle adaptations in heavy-resistance-trained women after detraining and retraining. *J. Appl. Physiol.* 70: 631-640, 1991.
218. Suga, T, Okita, K, Morita, N, Yokota, T, Hirabayashi, K, Horiuchi, M, Takada, S, Omokawa, M, Kinugawa, S, and Tsutsui, H. Dose effect on intramuscular metabolic stress during low-intensity resistance exercise with blood flow restriction. *J. Appl. Physiol.* 108: 1563-1567, 2010.
219. Takano, H, Morita, T, Iida, H, Asada, K, Kato, M, Uno, K, Hirose, K, Matsumoto, A, Takenaka, K, Hirata, Y, Eto, F, Nagai, R, Sato, Y, and Nakajima, T. Hemodynamic and hormonal responses to a short-term low-intensity resistance exercise with the reduction of muscle blood flow. *Eur. J. Appl. Physiol.* 95: 65-73, 2005.
220. Takarada, Y, Nakamura, Y, Aruga, S, Onda, T, Miyazaki, S, and Ishii, N. Rapid increase in plasma growth hormone after low-intensity resistance exercise with vascular occlusion. *J. Appl. Physiol.* 88: 61-65, 2000.
221. Tatsumi, R, and Allen, RE. Active hepatocyte growth factor is present in skeletal muscle extracellular matrix. *Muscle Nerve* 30: 654-658, 2004.
222. Tatsumi, R. Mechano-biology of skeletal muscle hypertrophy and regeneration: possible mechanism of stretch-induced activation of resident myogenic stem cells. *Anim. Sci. J.* 81: 11-20, 2010.
223. Tatsumi, R, Hattori, A, Ikeuchi, Y, Anderson, JE, and Allen, RE. Release of hepatocyte growth factor from mechanically stretched skeletal muscle satellite cells and role of pH and nitric oxide. *Mol. Biol. Cell* 13: 2909-2918, 2002.
224. Tesch, PA, and Larsson, L. Muscle hypertrophy in bodybuilders. *Eur. J. Appl. Physiol. Occup. Physiol.* 49: 301-306, 1982.
225. Tesch, PA. Skeletal muscle adaptations consequent to long-term heavy resistance exercise. *Med. Sci. Sports Exerc.* 20: S132-4, 1988.
226. Thomson, DM. The Role of AMPK in the Regulation of Skeletal Muscle Size, Hypertrophy, and Regeneration. *Int. J. Mol. Sci.* 19: 10.3390/ijms19103125, 2018.

227. Timmons, JA. Variability in training-induced skeletal muscle adaptation. *J. Appl. Physiol.* 110: 846-853, 2011.
228. Toigo, M, and Boutellier, U. New fundamental resistance exercise determinants of molecular and cellular muscle adaptations. *Eur. J. Appl. Physiol.* 97: 643-663, 2006.
229. Tomiya, A, Aizawa, T, Nagatomi, R, Sensui, H, and Kokubun, S. Myofibers express IL-6 after eccentric exercise. *Am. J. Sports Med.* 32: 503-508, 2004.
230. Trappe, S, Luden, N, Minchev, K, Raue, U, Jemiolo, B, and Trappe, TA. Skeletal muscle signature of a champion sprint runner. *J. Appl. Physiol.* (1985) 118: 1460-1466, 2015.
231. Trappe, TA, Raue, U, and Tesch, PA. Human soleus muscle protein synthesis following resistance exercise. *Acta Physiol. Scand.* 182: 189-196, 2004.
232. Travison, TG, Vesper, HW, Orwoll, E, Wu, F, Kaufman, JM, Wang, Y, Lapauw, B, Fiers, T, Matsumoto, AM, and Bhasin, S. Harmonized Reference Ranges for Circulating Testosterone Levels in Men of Four Cohort Studies in the United States and Europe. *J. Clin. Endocrinol. Metab.* 102: 1161-1173, 2017.
233. Urban, RJ, Bodenbun, YH, Gilkison, C, Foxworth, J, Coggan, AR, Wolfe, RR, and Ferrando, A. Testosterone administration to elderly men increases skeletal muscle strength and protein synthesis. *Am. J. Physiol.* 269: E820-6, 1995.
234. Van Cutsem, M, Duchateau, J, and Hainaut, K. Changes in single motor unit behaviour contribute to the increase in contraction speed after dynamic training in humans. *J. Physiol.* 513 (Pt 1): 295-305, 1998.
235. van der Pijl, R, Strom, J, Conijn, S, Lindqvist, J, Labeit, S, Granzier, H, and Ottenheijm, C. Titin-based mechanosensing modulates muscle hypertrophy. *J. Cachexia Sarcopenia Muscle* 9: 947-961, 2018.
236. van Wessel, T, de Haan, A, van der Laarse, WJ, and Jaspers, RT. The muscle fiber type-fiber size paradox: hypertrophy or oxidative metabolism? *Eur. J. Appl. Physiol.* 110: 665-694, 2010.
237. Veldhuis, JD, Keenan, DM, Mielke, K, Miles, JM, and Bowers, CY. Testosterone supplementation in healthy older men drives GH and IGF-I secretion without potentiating peptidyl secretagogue efficacy. *Eur. J. Endocrinol.* 153: 577-586, 2005.
238. Velloso, CP. Regulation of muscle mass by growth hormone and IGF-I. *Br. J. Pharmacol.* 154: 557-568, 2008.
239. Vierck, J, O'Reilly, B, Hossner, K, Antonio, J, Byrne, K, Bucci, L, and Dodson, M. Satellite cell regulation following myotrauma caused by resistance exercise. *Cell Biol. Int.* 24: 263-272, 2000.
240. Vingren, JL, Kraemer, WJ, Ratamess, NA, Anderson, JM, Volek, JS, and Maresh, CM. Testosterone physiology in resistance exercise and training: the up-stream regulatory elements. *Sports Med.* 40: 1037-1053, 2010.
241. Viru, M, Jansson, E, Viru, A, and Sundberg, CJ. Effect of restricted blood flow on exercise-induced hormone changes in healthy men. *Eur. J. Appl. Physiol. Occup. Physiol.* 77: 517-522, 1998.
242. Vollestad, NK, Vaage, O, and Hermansen, L. Muscle glycogen depletion patterns in type I and subgroups of type II fibres during prolonged severe exercise in man. *Acta Physiol. Scand.* 122: 433-441, 1984.
243. Wang, Q, and McPherron, AC. Myostatin inhibition induces muscle fibre hypertrophy prior to satellite cell activation. *J. Physiol.* 590: 2151-2165, 2012.
244. Wen, Y, Alimov, AP, and McCarthy, JJ. Ribosome Biogenesis is Necessary for Skeletal Muscle Hypertrophy. *Exerc. Sport Sci. Rev.* 44: 110-115, 2016.
245. West, DW, Kujbida, GW, Moore, DR, Atherton, P, Burd, NA, Padzik, JP, De Lisio, M, Tang, JE, Parise, G, Rennie, MJ, Baker, SK, and Phillips, SM. Resistance exercise-induced increases in putative anabolic hormones do not enhance muscle protein synthesis or intracellular signalling in young men. *J. Physiol.* 587: 5239-5247, 2009.
246. West, DW, Burd, NA, Tang, JE, Moore, DR, Staples, AW, Holwerda, AM, Baker, SK, and Phillips, SM. Elevations in ostensibly anabolic hormones with resistance exercise enhance neither training-induced muscle hypertrophy nor strength of the elbow flexors. *J. Appl. Physiol.* 108: 60-67, 2010.
247. West, DW, and Phillips, SM. Anabolic processes in human skeletal muscle: restoring the identities of growth hormone and testosterone. *Phys. Sportsmed* 38: 97-104, 2010.
248. West, DW, and Phillips, SM. Associations of exercise-induced hormone profiles and gains in strength and hypertrophy in a large cohort after weight training. *Eur. J. Appl. Physiol.* 112: 2693-2702, 2012.
249. Yamaguchi, A, Fujikawa, T, Shimada, S, Kanbayashi, I, Tateoka, M, Soya, H, Takeda, H, Morita, I, Matsubara, K, and Hirai, T. Muscle IGF-I Ea, MGF, and myostatin mRNA expressions after compensatory overload in hypophysectomized rats. *Pflugers Arch.* 453: 203-210, 2006.
250. Yang, SY, and Goldspink, G. Different roles of the IGF-I E peptide (MGF) and mature IGF-I in myoblast proliferation and differentiation. *FEBS Lett.* 522: 156-160, 2002.
251. Yao, W, Fuglevand, RJ, and Enoka, RM. Motor-unit synchronization increases EMG amplitude and decreases force steadiness of simulated contractions. *J. Neurophysiol.* 83: 441-452, 2000.
252. Yarasheski, KE, Campbell, JA, Smith, K, Rennie, MJ, Holloszy, JO, and Bier, DM. Effect of growth hormone and resistance exercise on muscle growth in young men. *Am. J. Physiol.* 262: E261-7, 1992.
253. Yarasheski, KE, Zachwieja, JJ, Campbell, JA, and Bier, DM. Effect of growth hormone and resistance exercise on muscle growth and strength in older men. *Am. J. Physiol.* 268: E268-76, 1995.
254. Zammit, PS. All muscle satellite cells are equal, but are some more equal than others? *J. Cell. Sci.* 121: 2975-2982, 2008.
255. Zanou, N, and Gailly, P. Skeletal muscle hypertrophy and regeneration: interplay between the myogenic regulatory factors (MRFs) and insulin-like growth factors (IGFs) pathways. *Cell Mol. Life Sci.* 70: 4117-4130, 2013.
256. Zhao, W, Pan, J, Zhao, Z, Wu, Y, Bauman, WA, and Cardozo, CP. Testosterone protects against dexamethasone-induced muscle atrophy, protein degradation and MAFbx upregulation. *J. Steroid Biochem. Mol. Biol.* 110: 125-129, 2008.
257. Zou, K, Meador, BM, Johnson, B, Huntsman, HD, Mahmassani, Z, Valero, MC, Huey, KA, and Boppart, MD. The alpha(7)beta(1)-integrin increases muscle hypertrophy following multiple bouts of eccentric exercise. *J. Appl. Physiol.* 111: 1134-1141, 2011.

Chapter 2

1. Abe, T, Beekley, MD, Hinata, S, Koizumi, K, and Sato, Y. Day-to-day change in muscle strength and MRI-measured skeletal muscle size during 7 days KAATSU resistance training: A case study. 2005;1:71-6. 39. *Int J Kaatsu Training Res.* 1: 71-76, 2005.
2. Abe, T, Yasuda, T, Midorikawa, T, Sato, Y, Kearns, C, Inoue, K, Koizumi, K, and Ishii, N. Skeletal muscle size and circulating IGF-1 are increased after two weeks of twice daily KAATSU resistance training. *Int J Kaatsu Training Res.* 1: 6-12, 2005.

3. Abe, T, Kearns, CE, and Sato, Y. Muscle size and strength are increased following walk training with restricted venous blood flow from the leg muscle, Kaatsu-walk training. *J. Appl. Physiol.* 100: 1460-1466, 2006.
4. Adams, G. The Molecular Response of Skeletal Muscle to Resistance Training. *Deutsche Zeitschrift für Sportmedizin* 61: 61-67, 2010.
5. Akima, H, and Saito, A. Activation of quadriceps femoris including vastus intermedius during fatiguing dynamic knee extensions. *Eur. J. Appl. Physiol.* 113: 2829-2840, 2013.
6. Allen, DG, Whitehead, NP, and Yeung, EW. Mechanisms of stretch-induced muscle damage in normal and dystrophic muscle: role of ionic changes. *J. Physiol.* 567: 723-735, 2005.
7. Aronson, D, Violan, MA, Dufresne, SD, Zangen, D, Fielding, RA, and Goodyear, LJ. Exercise stimulates the mitogen-activated protein kinase pathway in human skeletal muscle. *J. Clin. Invest.* 99: 1251-1257, 1997.
8. Aronson, D, Wojtaszewski, JF, Thorell, A, Nygren, J, Zangen, D, Richter, EA, Ljungqvist, O, Fielding, RA, and Goodyear, LJ. Extracellular-regulated protein kinase cascades are activated in response to injury in human skeletal muscle. *Am. J. Physiol.* 275: C555-61, 1998.
9. Atherton, PJ, Phillips, BE, and Wilkinson, DJ. Exercise and regulation of protein metabolism. In: *Progress in molecular biology and translational science*. Anonymous Cambridge, MA: Academic Press, 2015. pp. 75-98.
10. Bamman, MM, Shipp, JR, Jiang, J, Gower, BA, Hunter, GR, Goodman, A, McLafferty, CL, Jr, and Urban, RJ. Mechanical load increases muscle IGF-I and androgen receptor mRNA concentrations in humans. *Am. J. Physiol. Endocrinol. Metab.* 280: E383-90, 2001.
11. Bamman, MM. Take two NSAIDs and call on your satellite cells in the morning. *J. Appl. Physiol.* 103: 415-416, 2007.
12. Barash, IA, Mathew, L, Ryan, AF, Chen, J, and Lieber, RL. Rapid muscle-specific gene expression changes after a single bout of eccentric contractions in the mouse. *Am. J. Physiol. Cell. Physiol.* 286: C355-64, 2004.
13. Barton, ER, Morris, L, Musaro, A, Rosenthal, N, and Sweeney, HL. Muscle-specific expression of insulin-like growth factor I counters muscle decline in mdx mice. *J. Cell Biol.* 157: 137-148, 2002.
14. Bassel-Duby, R, and Olson, EN. Signaling pathways in skeletal muscle remodeling. *Annu. Rev. Biochem.* 75: 19-37, 2006.
15. Baum, C, Kennedy, DL, and Forbes, MB. Utilization of nonsteroidal antiinflammatory drugs. *Arthritis Rheum.* 28: 686-692, 1985.
16. Behm, DG. Neuromuscular implications and applications of resistance training. *J Strength Cond Res* 9: 264-274, 1995.
17. Belcastro, AN, Shewchuk, LD, and Raj, DA. Exercise-induced muscle injury: a calpain hypothesis. *Mol. Cell. Biochem.* 179: 135-145, 1998.
18. Biazon, TMPC, Ugrinowitsch, C, Soligon, SD, Oliveira, RM, Bergamasco, JG, Borghi-Silva, A, and Libardi, CA. The Association Between Muscle Deoxygenation and Muscle Hypertrophy to Blood Flow Restricted Training Performed at High and Low Loads. *Front. Physiol.* 10: 446, 2019.
19. Blaauw, B, Schiaffino, S, and Reggiani, C. Mechanisms modulating skeletal muscle phenotype. *Compr. Physiol.* 3: 1645-1687, 2013.
20. Bodine, SC, Stitt, TN, Gonzalez, M, Kline, WO, Stover, GL, Bauerlein, R, Zlotchenko, E, Scrimgeour, A, Lawrence, JC, Glass, DJ, and Yancopoulos, GD. Akt/mTOR pathway is a crucial regulator of skeletal muscle hypertrophy and can prevent muscle atrophy in vivo. *Nat. Cell Biol.* 3: 1014-1019, 2001.
21. Bondesen, BA, Mills, ST, Kegley, KM, and Pavlath, GK. The COX-2 pathway is essential during early stages of skeletal muscle regeneration. *Am. J. Physiol. Cell. Physiol.* 287: C475-83, 2004.
22. Bondesen, BA, Mills, ST, Kegley, KM, and Pavlath, GK. The COX-2 pathway is essential during early stages of skeletal muscle regeneration. *Am. J. Physiol., Cell Physiol.* 287: 475-483, 2004.
23. Bondesen, BA, Mills, ST, and Pavlath, GK. The COX-2 pathway regulates growth of atrophied muscle via multiple mechanisms. *Am. J. Physiol., Cell Physiol.* 290: 1651-1659, 2006.
24. Brentano, MA, and Martins Krueel, LF. A review on strength exercise-induced muscle damage: applications, adaptation mechanisms and limitations. *J. Sports Med. Phys. Fitness* 51: 1-10, 2011.
25. Brown, D, Hikim, AP, Kovacheva, EL, and Sinha-Hikim, I. Mouse model of testosterone-induced muscle fiber hypertrophy: involvement of p38 mitogen-activated protein kinase-mediated Notch signaling. *J. Endocrinol.* 201: 129-139, 2009.
26. Bruunsgaard, H, Galbo, H, Halkjaer-Kristensen, J, Johansen, TL, MacLean, DA, and Pedersen, BK. Exercise-induced increase in serum interleukin-6 in humans is related to muscle damage. *J. Physiol.* 499 (Pt 3): 833-841, 1997.
27. Bruunsgaard, H. Physical activity and modulation of systemic low-level inflammation. *J. Leukoc. Biol.* 78: 819-835, 2005.
28. Burd, NA, Dickinson, JM, Lemoine, JK, Carroll, CC, Sullivan, BE, Haus, JM, Jemiolo, B, Trappe, SW, Hughes, GM, Sanders, CE, Jr, and Trappe, TA. Effect of a cyclooxygenase-2 inhibitor on postexercise muscle protein synthesis in humans. *Am. J. Physiol. Endocrinol. Metab.* 298: E354-61, 2010.
29. Buresh, R, Berg, K, and French, J. The effect of resistive exercise rest interval on hormonal response, strength, and hypertrophy with training. *J Strength Cond Res* 23: 62-71, 2009.
30. Burian, M, and Geisslinger, G. COX-dependent mechanisms involved in the antinociceptive action of NSAIDs at central and peripheral sites. *Pharmacol. Ther.* 107: 139-154, 2005.
31. Burkholder, TJ. Mechanotransduction in skeletal muscle. *Front. Biosci.* 12: 174-191, 2007.
32. Campos, GER, Luecke, TJ, Wendeln, HK, Toma, K, Hagerman, FC, Murray, TF, Ragg, KE, Ratamess, NA, Kraemer, WJ, and Staron, RS. Muscular adaptations in response to three different resistance-training regimens: specificity of repetition maximum training zones. *Eur. J. Appl. Physiol.* 88: 50-60, 2002.
33. Carruthers, NJ, and Stemmer, PM. Methionine oxidation in the calmodulin-binding domain of calcineurin disrupts calmodulin binding and calcineurin activation. *Biochemistry (N. Y.)* 47: 3085-3095, 2008.
34. Cerda-Kohler, H, Henriquez-Olguin, C, Casas, M, Jensen, TE, Llanos, P, and Jaimovich, E. Lactate administration activates the ERK1/2, mTORC1, and AMPK pathways differentially according to skeletal muscle type in mouse. *Physiol. Rep.* 6: e13800, 2018.
35. Chan, MHS, Carey, AL, Watt, MJ, and Febbraio, MA. Cytokine gene expression in human skeletal muscle during concentric contraction: evidence that IL-8, like IL-6, is influenced by glycogen availability. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 287: 322-327, 2004.
36. Chen, TC, Yang, TJ, Huang, MJ, Wang, HS, Tseng, KW, Chen, HL, and Nosaka, K. Damage and the repeated bout effect of

- arm, leg, and trunk muscles induced by eccentric resistance exercises. *Scand. J. Med. Sci. Sports* 29: 725-735, 2019.
37. Chen, TC, Lin, K, Chen, H, Lin, M, and Nosaka, K. Comparison in eccentric exercise-induced muscle damage among four limb muscles. *Eur. J. Appl. Physiol.* 111: 211-223, 2011.
 38. Chin, ER. Role of Ca²⁺/calmodulin-dependent kinases in skeletal muscle plasticity. *J. Appl. Physiol.* 99: 414-423, 2005.
 39. Choi, J, Takahashi, H, and Itai, Y. The difference between effects of 'power-up type' and 'bulk-up type' strength training exercises: with special reference to muscle cross-sectional area. *Jpn J Phys Fitness Sports Med* 47: 119-129, 1998.
 40. Cirillo, F, Resmini, G, Ghiroldi, A, Piccoli, M, Bergante, S, Tettamanti, G, and Anastasia, L. Activation of the hypoxia-inducible factor 1 α promotes myogenesis through the noncanonical Wnt pathway, leading to hypertrophic myotubes. *FASEB J.* 31: 2146-2156, 2017.
 41. Clarke, MS, and Feedback, DL. Mechanical load induces sarcoplasmic wounding and FGF release in differentiated human skeletal muscle cultures. *FASEB J.* 10: 502-509, 1996.
 42. Clarke, MS, Bamman, MM, and Feedback, DL. Bed rest decreases mechanically induced myofiber wounding and consequent wound-mediated FGF release. *J. Appl. Physiol.* 85: 593-600, 1998.
 43. Clarkson, PM, Byrnes, WC, McCormick, KM, Turcotte, LP, and White, JS. Muscle soreness and serum creatine kinase activity following isometric, eccentric, and concentric exercise. *Int. J. Sports Med.* 7: 152-155, 1986.
 44. Clarkson, PM, Nosaka, K, and Braun, B. Muscle function after exercise-induced muscle damage and rapid adaptation. *Med. Sci. Sports Exerc.* 24: 512-520, 1992.
 45. Clarkson, PM, and Hubal, MJ. Exercise-induced muscle damage in humans. *Am. J. Phys. Med. Rehabil.* 81: 52-69, 2002.
 46. Cook, SB, Murphy, BG, and Labarbera, KE. Neuromuscular function after a bout of low-load blood flow-restricted exercise. *Med. Sci. Sports Exerc.* 45: 67-74, 2013.
 47. Croisier, JL, Camus, G, Venneman, I, Deby-Dupont, G, Juchmes-Ferir, A, Lamy, M, Crielaard, JM, Deby, C, and Duchateau, J. Effects of training on exercise-induced muscle damage and interleukin 6 production. *Muscle Nerve* 22: 208-212, 1999.
 48. Crossland, H, Kazi, AA, Lang, CH, Timmons, JA, Pierre, P, Wilkinson, DJ, Smith, K, Szweczyk, NJ, and Atherton, PJ. Focal adhesion kinase is required for IGF-I-mediated growth of skeletal muscle cells via a TSC2/mTOR/S6K1-associated pathway. *Am. J. Physiol. Endocrinol. Metab.* 305: E183-93, 2013.
 49. Damas, F, Phillips, SM, Libardi, CA, Vechin, FC, Lixandrao, ME, Jannig, PR, Costa, LA, Bacurau, AV, Snijders, T, Parise, G, Tricoli, V, Roschel, H, and Ugrinowitsch, C. Resistance training-induced changes in integrated myofibrillar protein synthesis are related to hypertrophy only after attenuation of muscle damage. *J. Physiol.* 594: 5209-5222, 2016.
 50. Dangott, B, Schultz, E, and Mozdziak, PE. Dietary creatine monohydrate supplementation increases satellite cell mitotic activity during compensatory hypertrophy. *Int. J. Sports Med.* 21: 13-16, 2000.
 51. Debold, EP. Recent Insights into the Molecular Basis of Muscular Fatigue. *Med. Sci. Sports Exerc.* , 2012.
 52. Dhawan, J, and Rando, TA. Stem cells in postnatal myogenesis: molecular mechanisms of satellite cell quiescence, activation and replenishment. *Trends Cell Biol.* 15: 666-673, 2005.
 53. Dimitrova, NA, and Dimitrov, GV. Interpretation of EMG changes with fatigue: facts, pitfalls, and fallacies. *J. Electromyogr. Kinesiol.* 13: 13-36, 2003.
 54. Dreyer, HC, Fujita, S, Cadenas, JG, Chinkes, DL, Volpi, E, and Rasmussen, BB. Resistance exercise increases AMPK activity and reduces 4E-BP1 phosphorylation and protein synthesis in human skeletal muscle. *J. Physiol.* 576: 613-624, 2006.
 55. Drummond, MJ, Fujita, S, Abe, T, Dreyer, HC, Volpi, E, and Rasmussen, BB. Human muscle gene expression following resistance exercise and blood flow restriction. *Med. Sci. Sports Exerc.* 40: 691-698, 2008.
 56. Dungan, CM. Less is more: the role of mTORC1 activation in the progression of ageing-mediated anabolic resistance. *J. Physiol.* 595: 2781-2782, 2017.
 57. Dunn, SE, Burns, JL, and Michel, RN. Calcineurin is required for skeletal muscle hypertrophy. *J. Biol. Chem.* 274: 21908-21912, 1999.
 58. Dunn, SE, Chin, ER, and Michel, RN. Matching of calcineurin activity to upstream effectors is critical for skeletal muscle fiber growth. *J. Cell Biol.* 151: 663-672, 2000.
 59. Ebbeling, CB, and Clarkson, PM. Exercise-induced muscle damage and adaptation. *Sports Med.* 7: 207-234, 1989.
 60. Eliasson, J, Elfegoun, T, Nilsson, J, Kohnke, R, Ekblom, B, and Blomstrand, E. Maximal lengthening contractions increase p70 S6 kinase phosphorylation in human skeletal muscle in the absence of nutritional supply. *Am. J. Physiol. Endocrinol. Metab.* 291: 1197-1205, 2006.
 61. Elkina, Y, von Haehling, S, Anker, SD, and Springer, J. The role of myostatin in muscle wasting: an overview. *J. Cachexia Sarcopenia Muscle* 2: 143-151, 2011.
 62. Enoka, RM. Eccentric contractions require unique activation strategies by the nervous system. *J. Appl. Physiol.* 81: 2339-2346, 1996.
 63. Evans, WJ, and Cannon, JG. The metabolic effects of exercise-induced muscle damage. *Exerc. Sport Sci. Rev.* 19: 99-9125, 1991.
 64. Farup, J, de Paoli, F, Bjerg, K, Riis, S, Ringgard, S, and Vissing, K. Blood flow restricted and traditional resistance training performed to fatigue produce equal muscle hypertrophy. *Scand. J. Med. Sci. Sports* 25: 754-763, 2015.
 65. Febbraio, MA, and Pedersen, BK. Muscle-derived interleukin-6: mechanisms for activation and possible biological roles. *FASEB J.* 16: 1335-1347, 2002.
 66. Fernandez-Gonzalo, R, Lundberg, TR, and Tesch, PA. Acute molecular responses in untrained and trained muscle subjected to aerobic and resistance exercise training versus resistance training alone. *Acta Physiol. (Oxf)* 209: 283-294, 2013.
 67. Figueiredo, VC, and McCarthy, JJ. Regulation of Ribosome Biogenesis in Skeletal Muscle Hypertrophy. *Physiology (Bethesda)* 34: 30-42, 2019.
 68. Finkenzeller, G, Newsome, W, Lang, F, and Haussinger, D. Increase of c-jun mRNA upon hypo-osmotic cell swelling of rat hepatoma cells. *FEBS Lett.* 340: 163-166, 1994.
 69. Flann, KL, LaStayo, PC, McClain, DA, Hazel, M, and Lindstedt, SL. Muscle damage and muscle remodeling: no pain, no gain? *J. Exp. Biol.* 214: 674-679, 2011.
 70. Foley, JM, Jayaraman, RC, Prior, BM, Pivarnik, JM, and Meyer, RA. MR measurements of muscle damage and adaptation after eccentric exercise. *J. Appl. Physiol.* 87: 2311-2318, 1999.
 71. Folland, JP, Irish, CS, Roberts, JC, Tarr, JE, and Jones, DA. Fatigue is not a necessary stimulus for strength gains during resistance training. *Br. J. Sports Med.* 36: 370-373, 2002.
 72. Formigli, L, Lombardo, LD, Adembri, C, Brunelleschi, S, Ferrari, E, and Novelli, GP. Neutrophils as mediators of human skeletal muscle ischemia-reperfusion syndrome. *Hum. Pathol.* 23: 627-634, 1992.

73. Foster, WH, Tidball, JG, and Wang, Y. P38gamma Activity is Required for Maintenance of Slow Skeletal Muscle Size. *Muscle Nerve* 45: 266-273, 2012.
74. Frey, JW, Farley, EE, O'Neil, TK, Burkholder, TJ, and Hornberger, TA. Evidence that mechanosensors with distinct biomechanical properties allow for specificity in mechanotransduction. *Biophys. J.* 97: 347-356, 2009.
75. Friedmann, B, Kinscherf, R, Borisch, S, Richter, G, Bartsch, P, and Billeter, R. Effects of low-resistance/high-repetition strength training in hypoxia on muscle structure and gene expression. *Pflugers Arch.* 446: 742-751, 2003.
76. Frigeri, A, Nicchia, GP, Verbavatz, JM, Valenti, G, and Svelto, M. Expression of aquaporin-4 in fast-twitch fibers of mammalian skeletal muscle. *J. Clin. Invest.* 102: 695-703, 1998.
77. Fry, AC. The role of resistance exercise intensity on muscle fibre adaptations. *Sports Med.* 34: 663-679, 2004.
78. Fry, CS, Glynn, EL, Drummond, MJ, Timmerman, KL, Fujita, S, Abe, T, Dhanani, S, Volpi, E, and Rasmussen, BB. Blood flow restriction exercise stimulates mTORC1 signaling and muscle protein synthesis in older men. *J. Appl. Physiol.* 108: 1199-1209, 2010.
79. Fujino, H, Xu, W, and Regan, JW. Prostaglandin E2 induced functional expression of early growth response factor-1 by EP4, but not EP2, prostanoid receptors via the phosphatidylinositol 3-kinase and extracellular signal-regulated kinases. *J. Biol. Chem.* 278: 12151-12156, 2003.
80. Fujita, T, Brechue, WF, Kurita, K, Sato, Y, and Abe, T. Increased muscle volume and strength following six days of low-intensity resistance training with restricted muscle blood flow. *Int J KAATSU Training Res* 4: 1-8, 2008.
81. Fujita, S, Abe, T, Drummond, MJ, Cadenas, JG, Dreyer, HC, Sato, Y, Volpi, E, and Rasmussen, BB. Blood flow restriction during low-intensity resistance exercise increases S6K1 phosphorylation and muscle protein synthesis. *J. Appl. Physiol.* 103: 903-910, 2007.
82. Garma, T, Kobayashi, C, Haddad, F, Adams, GR, Bodell, PW, and Baldwin, KM. Similar acute molecular responses to equivalent volumes of isometric, lengthening, or shortening mode resistance exercise. *J. Appl. Physiol.* 102: 135-143, 2007.
83. Gibala, MJ, MacDougall, JD, Tarnopolsky, MA, Stauber, WT, and Elorriaga, A. Changes in human skeletal muscle ultrastructure and force production after acute resistance exercise. *J. Appl. Physiol.* 78: 702-708, 1995.
84. Gibala, MJ, Interisano, SA, Tarnopolsky, MA, Roy, BD, MacDonald, JR, Yarasheski, KE, and MacDougall, JD. Myofibrillar disruption following acute concentric and eccentric resistance exercise in strength-trained men. *Can. J. Physiol. Pharmacol.* 78: 656-661, 2000.
85. Giordani, L, Parisi, A, and Le Grand, F. Satellite cell self-renewal. In: *Current topics in developmental biology*. Anonymous Cambridge, MA: Academic Press., 2018. pp. 177-203.
86. Glass, DJ. PI3 kinase regulation of skeletal muscle hypertrophy and atrophy. *Curr. Top. Microbiol. Immunol.* 346: 267-278, 2010.
87. Godfrey, RJ, Whyte, GP, Buckley, J, and Quinlivan, R. The role of lactate in the exercise-induced human growth hormone response: evidence from McArdle disease. *Br. J. Sports Med.* 43: 521-525, 2009.
88. Goldberg, AL, Etlinger, JD, Goldspink, DE, and Jablecki, C. Mechanism of work-induced hypertrophy of skeletal muscle. *Med. Sci. Sports* 7: 185-198, 1975.
89. Gomez-Cabrera, MC, Domenech, E, and Vina, J. Moderate exercise is an antioxidant: upregulation of antioxidant genes by training. *Free Radic. Biol. Med.* 44: 126-131, 2008.
90. Goodman, CA, Mayhew, DL, and Hornberger, TA. Recent progress toward understanding the molecular mechanisms that regulate skeletal muscle mass. *Cell. Signal.* 23: 1896-1906, 2011.
91. Goodman, CA, and Hornberger, TA. New roles for Smad signaling and phosphatidic acid in the regulation of skeletal muscle mass. *F1000Prime Rep.* 6: 20-20. eCollection 2014, 2014.
92. Goodman, CA. Role of mTORC1 in mechanically induced increases in translation and skeletal muscle mass. *J. Appl. Physiol.* (1985) 127: 581-590, 2019.
93. Gordon, SE, Kraemer, WJ, Vos, NH, Lynch, JM, and Knuttgen, HG. Effect of acid-base balance on the growth hormone response to acute high-intensity cycle exercise. *J. Appl. Physiol.* 76: 821-829, 1994.
94. Goto, K, Ishii, N, Kizuka, T, and Takamatsu, K. The impact of metabolic stress on hormonal responses and muscular adaptations. *Med. Sci. Sports Exerc.* 37: 955-963, 2005.
95. Gotshalk, LA, Loebel, CC, Nindl, BC, Putukian, M, Sebastianelli, WJ, Newton, RU, Häkkinen, K, and Kraemer, WJ. Hormonal responses of multiset versus single-set heavy-resistance exercise protocols. *Can. J. Appl. Physiol.* 22: 244-255, 1997.
96. Gumucio, JP, Sugg, KB, and Mendias, CL. TGF-beta superfamily signaling in muscle and tendon adaptation to resistance exercise. *Exerc. Sport Sci. Rev.* 43: 93-99, 2015.
97. Gundermann, D. Mechanisms of Blood Flow Restriction Exercise in Skeletal Muscle Adaptations. , 2016.
98. Gundermann, DM, Fry, CS, Dickinson, JM, Walker, DK, Timmerman, KL, Drummond, MJ, Volpi, E, and Rasmussen, BB. Reactive hyperemia is not responsible for stimulating muscle protein synthesis following blood flow restriction exercise. *J. Appl. Physiol.*, 2012.
99. Gute, DC, Ishida, T, Yarimizu, K, and Korthuis, RJ. Inflammatory responses to ischemia and reperfusion in skeletal muscle. *Mol. Cell. Biochem.* 179: 169-187, 1998.
100. Guyton, A. Textbook of medical physiology. In: Anonymous Philadelphia, PA: WB Saunders, 1986. pp. 366-368.
101. Haddad, F, and Adams, GR. Inhibition of MAP/ERK kinase prevents IGF-I-induced hypertrophy in rat muscles. *J. Appl. Physiol.* 96: 203-210, 2004.
102. Häkkinen, K, and Pakarinen, A. Acute hormonal responses to two different fatiguing heavy-resistance protocols in male athletes. *J. Appl. Physiol.* 74: 882-887, 1993.
103. Hall, JK, Banks, GB, Chamberlain, JS, and Olwin, BB. Prevention of muscle aging by myofiber-associated satellite cell transplantation. *Sci. Transl. Med.* 2: 57ra83, 2010.
104. Handayaniingsih, A, Iguchi, G, Fukuoka, H, Nishizawa, H, Takahashi, M, Yamamoto, M, Herningtyas, E, Okimura, Y, Kaji, H, Chihara, K, Seino, S, and Takahashi, Y. Reactive oxygen species play an essential role in IGF-I signaling and IGF-I-induced myocyte hypertrophy in C2C12 myocytes. *Endocrinology* 152: 912-921, 2011.
105. Hardy, D, Besnard, A, Latil, M, Jouvion, G, Briand, D, Thepenier, C, Pascal, Q, Guguin, A, Gayraud-Morel, B, Cavaillon, JM, Tajbakhsh, S, Rocheteau, P, and Chretien, F. Comparative Study of Injury Models for Studying Muscle Regeneration in Mice. *PLoS One* 11: e0147198, 2016.
106. Harridge, SD. Plasticity of human skeletal muscle: gene expression to in vivo function. *Exp. Physiol.* 92: 783-797, 2007.
107. Haussinger, D, Lang, E, and Gerok, W. Regulation of cell function by the cellular hydration state. *Am. J. Physiol.* 267: E343-55, 1994.

108. Haussinger, D. The role of cellular hydration in the regulation of cell function. *Biochem. J.* 313 (Pt 3): 697-710, 1996.
109. Hawley, JA. Molecular responses to strength and endurance training: are they incompatible? *Appl Physiol Nutr Metab* 34: 355-361, 2009.
110. Henneman, E, Somjen, G, and Carpenter, DO. Functional Significance of Cell Size in Spinal Motoneurons. *J. Neurophysiol.* 28: 560-580, 1965.
111. Hill, M, and Goldspink, G. Expression and splicing of the insulin-like growth factor gene in rodent muscle is associated with muscle satellite (stem) cell activation following local tissue damage. *J. Physiol. (Lond.)* 549: 409-418, 2003.
112. Hornberger, TA, McLoughlin, TJ, Leszczynski, JK, Armstrong, DD, Jameson, RR, Bowen, PE, Hwang, ES, Hou, H, Moustafa, ME, Carlson, BA, Hatfield, DL, Diamond, AM, and Esser, KA. Selenoprotein-deficient transgenic mice exhibit enhanced exercise-induced muscle growth. *J. Nutr.* 133: 3091-3097, 2003.
113. Hornberger, TA, Chu, WK, Mak, YW, Hsiung, JW, Huang, SA, and Chien, S. The role of phospholipase D and phosphatidic acid in the mechanical activation of mTOR signaling in skeletal muscle. *Proc. Natl. Acad. Sci. U. S. A.* 103: 4741-4746, 2006.
114. Hornberger, TA, Chu, WK, Mak, YW, Hsiung, JW, Huang, SA, and Chien, S. The role of phospholipase D and phosphatidic acid in the mechanical activation of mTOR signaling in skeletal muscle. *Proc. Natl. Acad. Sci. U. S. A.* 103: 4741-4746, 2006.
115. Horsley, V, and Pavlath, GK. Prostaglandin F2(alpha) stimulates growth of skeletal muscle cells via an NFATC2-dependent pathway. *J. Cell Biol.* 161: 111-118, 2003.
116. Houtman, CJ, Stegeman, DE, Van Dijk, JP, and Zwarts, MJ. Changes in muscle fiber conduction velocity indicate recruitment of distinct motor unit populations. *J. Appl. Physiol.* 95: 1045-1054, 2003.
117. Howatson, G, and Milak, A. Exercise-induced muscle damage following a bout of sport specific repeated sprints. *J Strength Cond Res* 23: 2419-2424, 2009.
118. Howell, JN, Chleboun, G, and Conatser, R. Muscle stiffness, strength loss, swelling and soreness following exercise-induced injury in humans. *J. Physiol. (Lond.)* 464: 183-196, 1993.
119. Hudson, MB, and Price, SR. Calcineurin: a poorly understood regulator of muscle mass. *Int. J. Biochem. Cell Biol.* 45: 2173-2178, 2013.
120. Huey, KA. Potential Roles of Vascular Endothelial Growth Factor During Skeletal Muscle Hypertrophy. *Exerc. Sport Sci. Rev.* 46: 195-202, 2018.
121. Hyldahl, RD, and Hubal, MJ. Lengthening our perspective: morphological, cellular, and molecular responses to eccentric exercise. *Muscle Nerve* 49: 155-170, 2014.
122. Hyldahl, RD, Chen, TC, and Nosaka, K. Mechanisms and Mediators of the Skeletal Muscle Repeated Bout Effect. *Exerc. Sport Sci. Rev.* 45: 24-33, 2017.
123. Ingemann-Hansen, T, Halkjaer-Kristensen, J, and Halskov, O. Skeletal muscle phosphagen and lactate concentrations in ischaemic dynamic exercise. *Eur. J. Appl. Physiol. Occup. Physiol.* 46: 261-270, 1981.
124. Ito, N, Ruegg, UT, and Takeda, S. ATP-Induced Increase in Intracellular Calcium Levels and Subsequent Activation of mTOR as Regulators of Skeletal Muscle Hypertrophy. *Int. J. Mol. Sci.* 19: 10.3390/ijms19092804, 2018.
125. Jacinto, E, and Hall, MN. Tor signalling in bugs, brain and brawn. *Nat. Rev. Mol. Cell Biol.* 4: 117-126, 2003.
126. Jackson, MJ. Free radicals generated by contracting muscle: by-products of metabolism or key regulators of muscle function? *Free Radic. Biol. Med.* 44: 132-141, 2008.
127. Ji, LL, Gomez-Cabrera, MC, and Vina, J. Exercise and hormesis: activation of cellular antioxidant signaling pathway. *Ann. N. Y. Acad. Sci.* 1067: 425-435, 2006.
128. Katch, VL, Katch, FI, Moffatt, R, and Gittleson, M. Muscular development and lean body weight in body builders and weight lifters. *Med. Sci. Sports Exerc.* 12: 340-344, 1980.
129. Kawada, S, and Ishii, N. Skeletal muscle hypertrophy after chronic restriction of venous blood flow in rats. *Med. Sci. Sports Exerc.* 37: 1144-1150, 2005.
130. Kefaloyianni, E, Gaitanaki, C, and Beis, I. ERK1/2 and p38-MAPK signalling pathways, through MSK1, are involved in NF-kappaB transactivation during oxidative stress in skeletal myoblasts. *Cell. Signal.* 18: 2238-2251, 2006.
131. Koh, TJ, and Pizza, FX. Do inflammatory cells influence skeletal muscle hypertrophy? *Front. Biosci. (Elite Ed)* 1: 60-71, 2009.
132. Komulainen, J, Kalliokoski, R, Koskinen, SO, Drost, MR, Kuipers, H, and Hesselink, MK. Controlled lengthening or shortening contraction-induced damage is followed by fiber hypertrophy in rat skeletal muscle. *Int. J. Sports Med.* 21: 107-112, 2000.
133. Kon, M, Ikeda, T, Homma, T, and Suzuki, Y. Effects of low-intensity resistance exercise under acute systemic hypoxia on hormonal responses. *J. Strength Cond Res.* 26: 611-617, 2012.
134. Kosek, DJ, Kim, JS, Petrella, JK, Cross, JM, and Bamman, MM. Efficacy of 3 days/wk resistance training on myofiber hypertrophy and myogenic mechanisms in young vs. older adults. *J. Appl. Physiol.* 101: 531-544, 2006.
135. Kraemer, WJ, Marchitelli, L, Gordon, SE, Harman, E, Dziados, JE, Mello, R, Frykman, P, McCurry, D, and Fleck, SJ. Hormonal and growth factor responses to heavy resistance exercise protocols. *J. Appl. Physiol.* 69: 1442-1450, 1990.
136. Kraemer, WJ, Gordon, SE, Fleck, SJ, Marchitelli, LJ, Mello, R, Dziados, JE, Friedl, K, Harman, E, Maresh, C, and Fry, AC. Endogenous anabolic hormonal and growth factor responses to heavy resistance exercise in males and females. *Int. J. Sports Med.* 12: 228-235, 1991.
137. Kraemer, WJ, Fleck, SJ, Dziados, JE, Harman, EA, Marchitelli, LJ, Gordon, SE, Mello, R, Frykman, PN, Koziris, LP, and Triplett, NT. Changes in hormonal concentrations after different heavy-resistance exercise protocols in women. *J. Appl. Physiol.* 75: 594-604, 1993.
138. Kraemer, WJ, Aguilera, BA, Terada, M, Newton, RU, Lynch, JM, Rosendaal, G, McBride, JM, Gordon, SE, and Häkkinen, K. Responses of IGF-I to endogenous increases in growth hormone after heavy-resistance exercise. *J. Appl. Physiol.* 79: 1310-1315, 1995.
139. Kraemer, WJ, and Ratamess, NA. Hormonal responses and adaptations to resistance exercise and training. *Sports Med.* 35: 339-361, 2005.
140. Kraemer, WJ, Adams, K, Cafarelli, E, Dudley, GA, Dooly, C, Feigenbaum, MS, Fleck, SJ, Franklin, B, Fry, AC, Hoffman, JR, Newton, RU, Pottenger, J, Stone, MH, Ratamess, NA, and Triplett-McBride, T. American College of Sports Medicine position stand. Progression models in resistance training for healthy adults. *Med. Sci. Sports Exerc.* 34: 364-380, 2002.
141. Kramer, HF, and Goodyear, LJ. Exercise, MAPK, and NF-kappaB signaling in skeletal muscle. *J. Appl. Physiol.* (1985) 103: 388-395, 2007.
142. Krentz, JR, Quest, B, Farthing, JP, Quest, DW, and Chilibeck, PD. The effects of ibuprofen on muscle hypertrophy, strength,

- and soreness during resistance training. *Appl. Physiol. Nutr. Metab.* 33: 470-475, 2008.
143. Krentz, JR, and Farthing, JP. Neural and morphological changes in response to a 20-day intense eccentric training protocol. *Eur. J. Appl. Physiol.* 110: 333-340, 2010.
 144. Kuipers, H. Exercise-induced muscle damage. *Int. J. Sports Med.* 15: 132-135, 1994.
 145. Lambert, CP, and Flynn, MG. Fatigue during high-intensity intermittent exercise: application to bodybuilding. *Sports Med.* 32: 511-522, 2002.
 146. Lambert, IH, Hoffmann, EK, and Pedersen, SF. Cell volume regulation: physiology and pathophysiology. *Acta Physiol. (Oxf)* 194: 255-282, 2008.
 147. Lang, F, Busch, GL, Ritter, M, Volkl, H, Waldegger, S, Gulbins, E, and Haussinger, D. Functional significance of cell volume regulatory mechanisms. *Physiol. Rev.* 78: 247-306, 1998.
 148. Lang, F. Mechanisms and significance of cell volume regulation. *J. Am. Coll. Nutr.* 26: 613S-623S, 2007.
 149. Latham, T, Mackay, L, Sproul, D, Karim, M, Culley, J, Harrison, DJ, Hayward, L, Langridge-Smith, P, Gilbert, N, and Ramsahoye, BH. Lactate, a product of glycolytic metabolism, inhibits histone deacetylase activity and promotes changes in gene expression. *Nucleic Acids Res.* 40: 4794-4803, 2012.
 150. Laurentino, GC, Ugrinowitsch, C, Roschel, H, Aoki, MS, Soares, AG, Neves, M, Jr, Aihara, AY, Fernandes Ada, R, and Tricoli, V. Strength training with blood flow restriction diminishes myostatin gene expression. *Med. Sci. Sports Exerc.* 44: 406-412, 2012.
 151. Lehman, N, Ledford, B, Di Fulvio, M, Frondorf, K, McPhail, LC, and Gomez-Cambronero, J. Phospholipase D2-derived phosphatidic acid binds to and activates ribosomal p70 S6 kinase independently of mTOR. *FASEB J.* 21: 1075-1087, 2007.
 152. Lessard, SJ, MacDonald, TL, Pathak, P, Han, MS, Coffey, VG, Edge, J, Rivas, DA, Hirshman, MF, Davis, RJ, and Goodyear, LJ. JNK regulates muscle remodeling via myostatin/SMAD inhibition. *Nat. Commun.* 9: 3030-018-05439-3, 2018.
 153. Lilja, M, Mandic, M, Apro, W, Melin, M, Olsson, K, Rosenberg, S, Gustafsson, T, and Lundberg, TR. High doses of anti-inflammatory drugs compromise muscle strength and hypertrophic adaptations to resistance training in young adults. *Acta Physiol. (Oxf)* , 2017.
 154. Loenneke, JP, Wilson, GJ, and Wilson, JM. A mechanistic approach to blood flow occlusion. *Int. J. Sports Med.* 31: 1-4, 2010.
 155. Loenneke, JP, Fahs, CA, Wilson, JM, and Bemben, MG. Blood flow restriction: the metabolite/volume threshold theory. *Med. Hypotheses* 77: 748-752, 2011.
 156. Loenneke, JP, Wilson, JM, Marin, PJ, Zourdos, MC, and Bemben, MG. Low intensity blood flow restriction training: a meta-analysis. *Eur. J. Appl. Physiol.* , 2011.
 157. Loenneke, JP, Thiebaud, RS, and Abe, T. Does blood flow restriction result in skeletal muscle damage? A critical review of available evidence. *Scand. J. Med. Sci. Sports* 24: e415-422, 2014.
 158. Low, SY, Rennie, MJ, and Taylor, PM. Signaling elements involved in amino acid transport responses to altered muscle cell volume. *FASEB J.* 11: 1111-1117, 1997.
 159. Lu, SS, Lau, CP, Tung, YF, Huang, SW, Chen, YH, Shih, HC, Tsai, SC, Lu, CC, Wang, SW, Chen, JJ, Chien, EJ, Chien, CH, and Wang, PS. Lactate and the effects of exercise on testosterone secretion: evidence for the involvement of a cAMP-mediated mechanism. *Med. Sci. Sports Exerc.* 29: 1048-1054, 1997.
 160. MacDougall, JD, Ray, S, Sale, DG, McCartney, N, Lee, P, and Garner, S. Muscle substrate utilization and lactate production. *Can. J. Appl. Physiol.* 24: 209-215, 1999.
 161. Mackey, AL, Kjaer, M, Dandanell, S, Mikkelsen, KH, Holm, L, Dossing, S, Kadi, F, Koskinen, SO, Jensen, CH, Schroder, HD, and Langberg, H. The influence of anti-inflammatory medication on exercise-induced myogenic precursor cell responses in humans. *J. Appl. Physiol.* 103: 425-431, 2007.
 162. MacNeil, LG, Melov, S, Hubbard, AE, Baker, SK, and Tarnopolsky, MA. Eccentric exercise activates novel transcriptional regulation of hypertrophic signaling pathways not affected by hormone changes. *PLoS One* 5: e10695, 2010.
 163. Mahmassani, ZS, Son, K, Pincu, Y, Munroe, M, Drnevich, J, Chen, J, and Boppart, MD. $\alpha 7 \beta 1$ Integrin regulation of gene transcription in skeletal muscle following an acute bout of eccentric exercise. *Am. J. Physiol. Cell. Physiol.* 312: C638-C650, 2017.
 164. Malm, C. Exercise-induced muscle damage and inflammation: fact or fiction? *Acta Physiol. Scand.* 171: 233-239, 2001.
 165. Manini, TM, and Clark, BC. Blood flow restricted exercise and skeletal muscle health. *Exerc. Sport Sci. Rev.* 37: 78-85, 2009.
 166. Manini, TM, Vincent, KR, Leeuwenburgh, CL, Lees, HA, Kavazis, AN, Borst, SE, and Clark, BC. Myogenic and proteolytic mRNA expression following blood flow restricted exercise. *Acta Physiol. (Oxf)* 201: 255-263, 2011.
 167. Martineau, LC, and Gardiner, PF. Insight into skeletal muscle mechanotransduction: MAPK activation is quantitatively related to tension. *J. Appl. Physiol.* 91: 693-702, 2001.
 168. Martineau, LC, and Gardiner, PF. Skeletal muscle is sensitive to the tension-time integral but not to the rate of change of tension, as assessed by mechanically induced signaling. *J. Biomech.* 35: 657-663, 2002.
 169. Martins, KJ, St-Louis, M, Murdoch, GK, MacLean, IM, McDonald, P, Dixon, WT, Putman, CT, and Michel, RN. Nitric oxide synthase inhibition prevents activity-induced calcineurin-NFATc1 signalling and fast-to-slow skeletal muscle fibre type conversions. *J. Physiol.* 590: 1427-1442, 2012.
 170. Mascher, H, Tannerstedt, J, Brink-Elfegoun, T, Ekblom, B, Gustafsson, T, and Blomstrand, E. Repeated resistance exercise training induces different changes in mRNA expression of MAFbx and MuRF-1 in human skeletal muscle. *Am. J. Physiol. Endocrinol. Metab.* 294: E43-51, 2008.
 171. Masuda, K, Choi, JY, Shimojo, H, and Katsuta, S. Maintenance of myoglobin concentration in human skeletal muscle after heavy resistance training. *Eur. J. Appl. Physiol. Occup. Physiol.* 79: 347-352, 1999.
 172. Mayhew, DL, Hornberger, TA, Lincoln, HC, and Bamman, MM. Eukaryotic initiation factor 2B epsilon induces cap-dependent translation and skeletal muscle hypertrophy. *J. Physiol.* 589: 3023-3037, 2011.
 173. McCarthy, JJ, Murach, KA. Anabolic and catabolic signaling pathways that regulate skeletal muscle mass. In: *Nutrition and Enhanced Sports Performance*. Anonymous Cambridge, MA: Academic Press, 2019. pp. 275-290.
 174. McCaulley, GO, McBride, JM, Cormie, P, Hudson, MB, Nuzzo, JL, Quindry, JC, and Travis Triplett, N. Acute hormonal and neuromuscular responses to hypertrophy, strength and power type resistance exercise. *Eur. J. Appl. Physiol.* 105: 695-704, 2009.
 175. McGee, SL, Mustard, KJ, Hardie, DG, and Baar, K. Normal hypertrophy accompanied by phosphorylation and activation of AMP-activated protein kinase $\alpha 1$ following overload in LKB1 knockout mice. *J. Physiol.* 586: 1731-1741, 2008.

176. McGinley, C, Shafat, A, and Donnelly, AE. Does antioxidant vitamin supplementation protect against muscle damage? *Sports Med.* 39: 1011-1032, 2009.
177. McHugh, MP. Recent advances in the understanding of the repeated bout effect: the protective effect against muscle damage from a single bout of eccentric exercise. *Scand. J. Med. Sci. Sports* 13: 88-97, 2003.
178. McKay, BR, O'Reilly, CE, Phillips, SM, Tarnopolsky, MA, and Parise, G. Co-expression of IGF-1 family members with myogenic regulatory factors following acute damaging muscle-lengthening contractions in humans. *J. Physiol.* 586: 5549-5560, 2008.
179. McKinsey, TA, Zhang, CL, and Olson, EN. Signaling chromatin to make muscle. *Curr. Opin. Cell Biol.* 14: 763-772, 2002.
180. Mendias, CL, Tatsumi, R, and Allen, RE. Role of cyclooxygenase-1 and -2 in satellite cell proliferation, differentiation, and fusion. *Muscle Nerve* 30: 497-500, 2004.
181. Merry, TL, and Ristow, M. Do antioxidant supplements interfere with skeletal muscle adaptation to exercise training? *J. Physiol.* 594: 5135-5147, 2016.
182. Meyer, RA. Does blood flow restriction enhance hypertrophic signaling in skeletal muscle? *J. Appl. Physiol.* 100: 1443-1444, 2006.
183. Michel, RN, Dunn, SE, and Chin, ER. Calcineurin and skeletal muscle growth. *Proc. Nutr. Soc.* 63: 341-349, 2004.
184. Mikkelsen, UR, Langberg, H, Helmark, IC, Skovgaard, D, Andersen, LL, Kjaer, M, and Mackey, AL. Local NSAID infusion inhibits satellite cell proliferation in human skeletal muscle after eccentric exercise. *J. Appl. Physiol.* 107: 1600-1611, 2009.
185. Mikkelsen, UR, Schjerling, P, Helmark, IC, Reitelseder, S, Holm, L, Skovgaard, D, Langberg, H, Kjaer, M, and Heine-meier, KM. Local NSAID infusion does not affect protein synthesis and gene expression in human muscle after eccentric exercise. *Scand. J. Med. Sci. Sports* 21: 630-644, 2011.
186. Miller, KJ, Garland, SJ, Ivanova, T, and Ohtsuki, T. Motor-unit behavior in humans during fatiguing arm movements. *J. Neurophysiol.* 75: 1629-1636, 1996.
187. Miyazaki, M, and Esser, KA. Cellular mechanisms regulating protein synthesis and skeletal muscle hypertrophy in animals. *J. Appl. Physiol.* 106: 1367-1373, 2009.
188. Miyazaki, M, McCarthy, JJ, Fedele, MJ, and Esser, KA. Early activation of mTORC1 signalling in response to mechanical overload is independent of phosphoinositide 3-kinase/Akt signalling. *J. Physiol.* 589: 1831-1846, 2011.
189. Moriya, N, and Miyazaki, M. Akt1 deficiency diminishes skeletal muscle hypertrophy by reducing satellite cell proliferation. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 314: R741-R751, 2018.
190. Morton, RW, Sonne, MW, Farias Zuniga, A, Mohammad, IYZ, Jones, A, McGlory, C, Keir, PJ, Potvin, JR, and Phillips, SM. Muscle fibre activation is unaffected by load and repetition duration when resistance exercise is performed to task failure. *J. Physiol.* 597: 4601-4613, 2019.
191. Muddle, TWD, Colquhoun, RJ, Magrini, MA, Luera, MJ, DeFreitas, JM, and Jenkins, NDM. Effects of fatiguing, submaximal high- versus low-torque isometric exercise on motor unit recruitment and firing behavior. *Physiol. Rep.* 6: e13675, 2018.
192. Murach, KA, Englund, DA, Dupont-Versteegden, EE, McCarthy, JJ, and Peterson, CA. Myonuclear Domain Flexibility Challenges Rigid Assumptions on Satellite Cell Contribution to Skeletal Muscle Fiber Hypertrophy. *Front. Physiol.* 9: 635, 2018.
193. Nader, GA, and Esser, KA. Intracellular signaling specificity in skeletal muscle in response to different modes of exercise. *J. Appl. Physiol.* (1985) 90: 1936-1942, 2001.
194. Nakashima, K, and Yakabe, Y. AMPK activation stimulates myofibrillar protein degradation and expression of atrophy-related ubiquitin ligases by increasing FOXO transcription factors in C2C12 myotubes. *Biosci. Biotechnol. Biochem.* 71: 1650-1656, 2007.
195. Naya, FJ, Mercer, B, Shelton, J, Richardson, JA, Williams, RS, and Olson, EN. Stimulation of slow skeletal muscle fiber gene expression by calcineurin in vivo. *J. Biol. Chem.* 275: 4545-4548, 2000.
196. Nguyen, HX, and Tidball, JG. Null mutation of gp91phox reduces muscle membrane lysis during muscle inflammation in mice. *J. Physiol. (Lond.)* 553: 833-841, 2003.
197. Nielsen, AR, and Pedersen, BK. The biological roles of exercise-induced cytokines: IL-6, IL-8, and IL-15. *Appl. Physiol. Nutr. Metab.* 32: 833-839, 2007.
198. Nishimura, A, Sugita, M, Kato, K, Fukuda, A, Sudo, A, and Uchida, A. Hypoxia increases muscle hypertrophy induced by resistance training. *Int. J. Sports Physiol. Perform.* 5: 497-508, 2010.
199. Nosaka, K, Lavender, A, Newton, M, and Sacco, P. Muscle damage in resistance training: Is muscle damage necessary for strength gain and muscle hypertrophy? *IJSHS* 1: 1-8, 2003.
200. Nosaka, K, and Clarkson, PM. Changes in indicators of inflammation after eccentric exercise of the elbow flexors. *Med. Sci. Sports Exerc.* 28: 953-961, 1996.
201. Novak, ML, Billich, W, Smith, SM, Sukhija, KB, McLoughlin, TJ, Hornberger, TA, and Koh, TJ. COX-2 inhibitor reduces skeletal muscle hypertrophy in mice. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 296: R1132-9, 2009.
202. Ogasawara, R, and Sugino, T. Rapamycin-insensitive mechanistic target of rapamycin regulates basal and resistance exercise-induced muscle protein synthesis. *FASEB J.* : fj201701422R, 2018.
203. Ogasawara, R, Jensen, TE, Goodman, CA, and Hornberger, TA. Resistance Exercise-Induced Hypertrophy: A Potential Role for Rapamycin-Insensitive mTOR. *Exerc. Sport Sci. Rev.* 47: 188-194, 2019.
204. Ohno, Y, Ando, K, Ito, T, Suda, Y, Matsui, Y, Oyama, A, Kaneko, H, Yokoyama, S, Egawa, T, and Goto, K. Lactate Stimulates a Potential for Hypertrophy and Regeneration of Mouse Skeletal Muscle. *Nutrients* 11: 10.3390/nu11040869, 2019.
205. Oishi, Y, Tsukamoto, H, Yokokawa, T, Hirotsu, K, Shimazu, M, Uchida, K, Tomi, H, Higashida, K, Iwanaka, N, and Hashimoto, T. Mixed lactate and caffeine compound increases satellite cell activity and anabolic signals for muscle hypertrophy. *J. Appl. Physiol.* (1985) 118: 742-749, 2015.
206. O'Neil, TK, Duffy, LR, Frey, JW, and Hornberger, TA. The role of phosphoinositide 3-kinase and phosphatidic acid in the regulation of mammalian target of rapamycin following eccentric contractions. *J. Physiol.* 587: 3691-3701, 2009.
207. Otis, JS, Burkholder, TJ, and Pavlath, GK. Stretch-induced myoblast proliferation is dependent on the COX2 pathway. *Exp. Cell Res.* 310: 417-425, 2005.
208. Palmer, RM. Prostaglandins and the control of muscle protein synthesis and degradation. *Prostaglandins Leukot. Essent. Fatty Acids* 39: 95-104, 1990.
209. Parsons, SA, Millay, DP, Wilkins, BJ, Bueno, OF, Tsika, GL, Neilson, JR, Liberatore, CM, Yutzey, KE, Crabtree, GR, Tsika, RW, and Molkentin, JD. Genetic loss of calcineurin blocks mechanical overload-induced skeletal muscle fiber type

- switching but not hypertrophy. *J. Biol. Chem.* 279: 26192-26200, 2004.
210. Pasiakos, SM. Exercise and amino acid anabolic cell signaling and the regulation of skeletal muscle mass. *Nutrients* 4: 740-758, 2012.
 211. Pasiakos, SM, and Carbone, JW. Assessment of skeletal muscle proteolysis and the regulatory response to nutrition and exercise. *IUBMB Life* 66: 478-484, 2014.
 212. Paulsen, G, Egner, IM, Drange, M, Langberg, H, Benestad, HB, Fjeld, JG, Hallen, J, and Raastad, T. A COX-2 inhibitor reduces muscle soreness, but does not influence recovery and adaptation after eccentric exercise. *Scand. J. Med. Sci. Sports* 20: e195-207, 2010.
 213. Pedersen, BK, Ostrowski, K, Rohde, T, and Bruunsgaard, H. The cytokine response to strenuous exercise. *Can. J. Physiol. Pharmacol.* 76: 505-511, 1998.
 214. Petersen, SG, Beyer, N, Hansen, M, Holm, L, Aagaard, P, Mackey, AL, and Kjaer, M. Nonsteroidal anti-inflammatory drug or glucosamine reduced pain and improved muscle strength with resistance training in a randomized controlled trial of knee osteoarthritis patients. *Arch. Phys. Med. Rehabil.* 92: 1185-1193, 2011.
 215. Petersen, SG, Miller, BF, Hansen, M, Kjaer, M, and Holm, L. Exercise and NSAIDs: effect on muscle protein synthesis in patients with knee osteoarthritis. *Med. Sci. Sports Exerc.* 43: 425-431, 2011.
 216. Petrella, JK, Kim, JS, Cross, JM, Kosek, DJ, and Bamman, MM. Efficacy of myonuclear addition may explain differential myofiber growth among resistance-trained young and older men and women. *Am. J. Physiol. Endocrinol. Metab.* 291: E937-46, 2006.
 217. Petrella, JK, Kim, J, Mayhew, DL, Cross, JM, and Bamman, MM. Potent myofiber hypertrophy during resistance training in humans is associated with satellite cell-mediated myonuclear addition: a cluster analysis. *J. Appl. Physiol.* 104: 1736-1742, 2008.
 218. Pierce, JR, Clark, BC, Ploutz-Snyder, LL, and Kanaley, JA. Growth hormone and muscle function responses to skeletal muscle ischemia. *J. Appl. Physiol.* 101: 1588-1595, 2006.
 219. Pistilli, EE, and Quinn, LS. From anabolic to oxidative: reconsidering the roles of IL-15 and IL-15R α in skeletal muscle. *Exerc. Sport Sci. Rev.* 41: 100-106, 2013.
 220. Proske, U, and Morgan, DL. Muscle damage from eccentric exercise: mechanism, mechanical signs, adaptation and clinical applications. *J. Physiol.* 537: 333-345, 2001.
 221. Quinn, LS. Interleukin-15: a muscle-derived cytokine regulating fat-to-lean body composition. *J. Anim. Sci.* 86: E75-83, 2008.
 222. Raue, U, Trappe, TA, Estrem, ST, Qian, HR, Helvering, LM, Smith, RC, and Trappe, S. Transcriptome signature of resistance exercise adaptations: mixed muscle and fiber type specific profiles in young and old adults. *J. Appl. Physiol.* (1985) 112: 1625-1636, 2012.
 223. Reeves, GV, Kraemer, RR, Hollander, DB, Clavier, J, Thomas, C, Francois, M, and Castracane, VD. Comparison of hormone responses following light resistance exercise with partial vascular occlusion and moderately difficult resistance exercise without occlusion. *J. Appl. Physiol.* 101: 1616-1622, 2006.
 224. Riechman, SE, Balasekaran, G, Roth, SM, and Ferrell, RE. Association of interleukin-15 protein and interleukin-15 receptor genetic variation with resistance exercise training responses. *J. Appl. Physiol.* 97: 2214-2219, 2004.
 225. Rieu, I, Magne, H, Savary-Auzeloux, I, Averous, J, Bos, C, Peyron, MA, Combaret, L, and Dardevet, D. Reduction of low grade inflammation restores blunting of postprandial muscle anabolism and limits sarcopenia in old rats. *J. Physiol.* 587: 5483-5492, 2009.
 226. Rindom, E, Kristensen, AM, Overgaard, K, Vissing, K, and de Paoli, FV. Activation of mTORC1 signalling in rat skeletal muscle is independent of the EC-coupling sequence but dependent on tension per se in a dose-response relationship. *Acta Physiol. (Oxf)* 227: e13336, 2019.
 227. Robbins, DW, Goodale, TL, Docherty, D, Behm, DG, and Tran, QT. The effects of load and training pattern on acute neuromuscular responses in the upper body. *J. Strength Cond Res.* 24: 2996-3007, 2010.
 228. Rodemann, HP, and Goldberg, AL. Arachidonic acid, prostaglandin E₂ and F₂ α influence rates of protein turnover in skeletal and cardiac muscle. *J. Biol. Chem.* 257: 1632-1638, 1982.
 229. Rodriguez, J, Vernus, B, Chelhel, I, Cassar-Malek, I, Gabilard, JC, Hadj Sassi, A, Seiliez, I, Picard, B, and Bonniieu, A. Myostatin and the skeletal muscle atrophy and hypertrophy signaling pathways. *Cell Mol. Life Sci.* 71: 4361-4371, 2014.
 230. Rooney, KJ, Herbert, RD, and Balnave, RJ. Fatigue contributes to the strength training stimulus. *Med. Sci. Sports Exerc.* 26: 1160-1164, 1994.
 231. Roux, PP, and Blenis, J. ERK and p38 MAPK-activated protein kinases: a family of protein kinases with diverse biological functions. *Microbiol. Mol. Biol. Rev.* 68: 320-344, 2004.
 232. Rubin, MR, Kraemer, WJ, Maresh, CM, Volek, JS, Ratamess, NA, Vanheest, JL, Silvestre, R, French, DN, Sharman, MJ, Judelson, DA, Gomez, AL, Vescovi, JD, and Hymer, WC. High-affinity growth hormone binding protein and acute heavy resistance exercise. *Med. Sci. Sports Exerc.* 37: 395-403, 2005.
 233. Russell, B, Dix, DJ, Haller, DL, and Jacobs-El, J. Repair of injured skeletal muscle: a molecular approach. *Med. Sci. Sports Exerc.* 24: 189-196, 1992.
 234. Sahlin, K, Soderlund, K, Tonkonogi, M, and Hirakoba, K. Phosphocreatine content in single fibers of human muscle after sustained submaximal exercise. *Am. J. Physiol.* 273: C172-8, 1997.
 235. Saxton, JM, Donnelly, AE, and Roper, HP. Indices of free-radical-mediated damage following maximum voluntary eccentric and concentric muscular work. *Eur. J. Appl. Physiol. Occup. Physiol.* 68: 189-193, 1994.
 236. Schliess, F, Schreiber, R, and Haussinger, D. Activation of extracellular signal-regulated kinases Erk-1 and Erk-2 by cell swelling in H4IIE hepatoma cells. *Biochem. J.* 309 (Pt 1): 13-17, 1995.
 237. Schliess, F, Richter, L, vom Dahl, S, and Haussinger, D. Cell hydration and mTOR-dependent signalling. *Acta Physiol. (Oxf)* 187: 223-229, 2006.
 238. Schmidtbleicher, D, Buehrle, M. Neuronal adaptation and increase of cross-sectional area studying different strength training methods. In: Biomechanics X-B volume 6-B. Champaign (IL): Human Kinetics, 1987: 615-20. Jonsson, GB, ed. Champaign, IL: Human Kinetics, 1987. pp. 615-620.
 239. Schoenfeld, BJ. Potential mechanisms for a role of metabolic stress in hypertrophic adaptations to resistance training. *Sports Med.* , In Press.
 240. Schoenfeld, BJ. The mechanisms of muscle hypertrophy and their application to resistance training. *J. Strength Cond Res.* 24: 2857-2872, 2010.

241. Schoenfeld, BJ. Does exercise-induced muscle damage play a role in skeletal muscle hypertrophy? *J. Strength Cond Res.* 26: 1441-1453, 2012.
242. Schoenfeld, BJ, Contreras, B, Willardson, JM, Fontana, E, and Tiryaki-Sonmez, G. Muscle activation during low- versus high-load resistance training in well-trained men. *Eur. J. Appl. Physiol.* 114: 2491-2497, 2014.
243. Schoenfeld, BJ, Ratamess, NA, Peterson, MD, Contreras, B, Tiryaki-Sonmez, G, and Alvar, BA. Effects of different volume-equated resistance training loading strategies on muscular adaptations in well-trained men. *J. Strength Cond Res.* 28: 2909-2918, 2014.
244. Schott, J, McCully, K, and Rutherford, OM. The role of metabolites in strength training. II. Short versus long isometric contractions. *Eur. J. Appl. Physiol. Occup. Physiol.* 71: 337-341, 1995.
245. Schultz, E, Jaryszak, DL, and Valliere, CR. Response of satellite cells to focal skeletal muscle injury. *Muscle Nerve* 8: 217-222, 1985.
246. Schwane, JA, Johnson, SR, Vandenakker, CB, and Armstrong, RB. Delayed-onset muscular soreness and plasma CPK and LDH activities after downhill running. *Med. Sci. Sports Exerc.* 15: 51-56, 1983.
247. Scott, BR, Slaterry, KM, and Dascombe, BJ. Intermittent hypoxic resistance training: Is metabolic stress the key moderator? *Med. Hypotheses* 84: 145-149, 2015.
248. Sengupta, S, Peterson, TR, and Sabatini, DM. Regulation of the mTOR complex 1 pathway by nutrients, growth factors, and stress. *Mol. Cell* 40: 310-322, 2010.
249. Shill, DD, Polley, KR, Willingham, TB, Call, JA, Murrow, JR, McCully, KK, and Jenkins, NT. Experimental intermittent ischemia augments exercise-induced inflammatory cytokine production. *J. Appl. Physiol.* (1985) 123: 434-441, 2017.
250. Shinohara, M, Kouzaki, M, Yoshihisa, T, and Fukunaga, T. Efficacy of tourniquet ischemia for strength training with low resistance. *Eur. J. Appl. Physiol. Occup. Physiol.* 77: 189-191, 1998.
251. Sinha-Hikim, I, Cornford, M, Gaytan, H, Lee, ML, and Bhasin, S. Effects of testosterone supplementation on skeletal muscle fiber hypertrophy and satellite cells in community-dwelling older men. *J. Clin. Endocrinol. Metab.* 91: 3024-3033, 2006.
252. Sjogaard, G, Adams, RP, and Saltin, B. Water and ion shifts in skeletal muscle of humans with intense dynamic knee extension. *Am. J. Physiol.* 248: R190-6, 1985.
253. Sjogaard, G. Water and electrolyte fluxes during exercise and their relation to muscle fatigue. *Acta Physiol. Scand. Suppl.* 556: 129-136, 1986.
254. Smilios, I, Pilianidis, T, Karamouzis, M, and Tokmakidis, SP. Hormonal responses after various resistance exercise protocols. *Med. Sci. Sports Exerc.* 35: 644-654, 2003.
255. Smith, RC, and Rutherford, OM. The role of metabolites in strength training. I. A comparison of eccentric and concentric contractions. *Eur. J. Appl. Physiol. Occup. Physiol.* 71: 332-336, 1995.
256. Soltow, QA, Betters, JL, Sellman, JE, Lira, VA, Long, JH, and Criswell, DS. Ibuprofen inhibits skeletal muscle hypertrophy in rats. *Med. Sci. Sports Exerc.* 38: 840-846, 2006.
257. Soltow, QA, Betters, JL, Sellman, JE, Lira, VA, Long, JHD, and Criswell, DS. Ibuprofen inhibits skeletal muscle hypertrophy in rats. *Med. Sci. Sports Exerc.* 38: 840-846, 2006.
258. Spangenburg, EE. Changes in muscle mass with mechanical load: possible cellular mechanisms. *Appl. Physiol. Nutr. Metab.* 34: 328-335, 2009.
259. Street, SF. Lateral transmission of tension in frog myofibers: a myofibrillar network and transverse cytoskeletal connections are possible transmitters. *J. Cell. Physiol.* 114: 346-364, 1983.
260. Suga, T, Okita, K, Morita, N, Yokota, T, Hirabayashi, K, Horiuchi, M, Takada, S, Omokawa, M, Kinugawa, S, and Tsutsui, H. Dose effect on intramuscular metabolic stress during low-intensity resistance exercise with blood flow restriction. *J. Appl. Physiol.* 108: 1563-1567, 2010.
261. Suga, T, Okita, K, Morita, N, Yokota, T, Hirabayashi, K, Horiuchi, M, Takada, S, Takahashi, T, Omokawa, M, Kinugawa, S, and Tsutsui, H. Intramuscular metabolism during low-intensity resistance exercise with blood flow restriction. *J. Appl. Physiol.* 106: 1119-1124, 2009.
262. Suzuki, YJ, and Ford, GD. Redox regulation of signal transduction in cardiac and smooth muscle. *J. Mol. Cell. Cardiol.* 31: 345-353, 1999.
263. Takada, S, Okita, K, Suga, T, Omokawa, M, Kadoguchi, T, Sato, T, Takahashi, M, Yokota, T, Hirabayashi, K, Morita, N, Horiuchi, M, Kinugawa, S, and Tsutsui, H. Low-intensity exercise can increase muscle mass and strength proportionally to enhanced metabolic stress under ischemic conditions. *J. Appl. Physiol.* (1985) 113: 199-205, 2012.
264. Takano, H, Morita, T, Iida, H, Asada, K, Kato, M, Uno, K, Hirose, K, Matsumoto, A, Takenaka, K, Hirata, Y, Eto, F, Nagai, R, Sato, Y, and Nakajima, T. Hemodynamic and hormonal responses to a short-term low-intensity resistance exercise with the reduction of muscle blood flow. *Eur. J. Appl. Physiol.* 95: 65-73, 2005.
265. Takarada, Y, Nakamura, Y, Aruga, S, Onda, T, Miyazaki, S, and Ishii, N. Rapid increase in plasma growth hormone after low-intensity resistance exercise with vascular occlusion. *J. Appl. Physiol.* 88: 61-65, 2000.
266. Takarada, Y, Takazawa, H, Sato, Y, Takebayashi, S, Tanaka, Y, and Ishii, N. Effects of resistance exercise combined with moderate vascular occlusion on muscular function in humans. *J. Appl. Physiol.* 88: 2097-2106, 2000.
267. Talmadge, RJ, Otis, JS, Rittler, MR, Garcia, ND, Spencer, SR, Lees, SJ, and Naya, FJ. Calcineurin activation influences muscle phenotype in a muscle-specific fashion. *BMC Cell Biol.* 5: 28, 2004.
268. Tamaki, T, Uchiyama, S, Tamura, T, and Nakano, S. Changes in muscle oxygenation during weight-lifting exercise. *Eur. J. Appl. Physiol. Occup. Physiol.* 68: 465-469, 1994.
269. Tatsumi, R, Hattori, A, Ikeuchi, Y, Anderson, JE, and Allen, RE. Release of hepatocyte growth factor from mechanically stretched skeletal muscle satellite cells and role of pH and nitric oxide. *Mol. Biol. Cell* 13: 2909-2918, 2002.
270. Tee, JC, Bosch, AN, and Lambert, MI. Metabolic consequences of exercise-induced muscle damage. *Sports Med.* 37: 827-836, 2007.
271. Terzis, G, Georgiadis, G, Stratakos, G, Vogiatzis, I, Kavouras, S, Manta, P, Mascher, H, and Blomstrand, E. Resistance exercise-induced increase in muscle mass correlates with p70S6 kinase phosphorylation in human subjects. *Eur. J. Appl. Physiol.* 102: 145-152, 2008.
272. Tesch, PA, Colliander, EB, and Kaiser, P. Muscle metabolism during intense, heavy-resistance exercise. *Eur. J. Appl. Physiol. Occup. Physiol.* 55: 362-366, 1986.
273. Thannickal, VJ, and Fanburg, BL. Reactive oxygen species in cell signaling. *Am. J. Physiol. Lung Cell. Mol. Physiol.* 279: L1005-28, 2000.
274. Thomas, G, and Hall, MN. TOR signalling and control of cell growth. *Curr. Opin. Cell Biol.* 9: 782-787, 1997.

275. Thomson, DM, and Gordon, SE. Impaired overload-induced muscle growth is associated with diminished translational signalling in aged rat fast-twitch skeletal muscle. *J. Physiol.* 574: 291-305, 2006.
276. Tidball, JG. Mechanical signal transduction in skeletal muscle growth and adaptation. *J. Appl. Physiol.* 98: 1900-1908, 2005.
277. Tidball, JG. Inflammatory processes in muscle injury and repair. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 288: 345-353, 2005.
278. Toft, AD, Jensen, LB, Bruunsgaard, H, Ibfelt, T, Halkjaer-Kristensen, J, Febbraio, M, and Pedersen, BK. Cytokine response to eccentric exercise in young and elderly humans. *Am. J. Physiol. Cell Physiol.* 283: 289-295, 2002.
279. Toigo, M, and Boutellier, U. New fundamental resistance exercise determinants of molecular and cellular muscle adaptations. *Eur. J. Appl. Physiol.* 97: 643-663, 2006.
280. Trappe, TA, White, F, Lambert, CP, Cesar, D, Hellerstein, M, and Evans, WJ. Effect of ibuprofen and acetaminophen on postexercise muscle protein synthesis. *Am. J. Physiol. Endocrinol. Metab.* 282: E551-6, 2002.
281. Trappe, TA, Carroll, CC, Dickinson, JM, LeMoine, JK, Haus, JM, Sullivan, BE, Lee, JD, Jemio, B, Weinheimer, EM, and Hollon, CJ. Influence of acetaminophen and ibuprofen on skeletal muscle adaptations to resistance exercise in older adults. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 300: R655-62, 2011.
282. Tsukamoto, S, Shibasaki, A, Naka, A, Saito, H, and Iida, K. Lactate Promotes Myoblast Differentiation and Myotube Hypertrophy via a Pathway Involving MyoD In Vitro and Enhances Muscle Regeneration In Vivo. *Int. J. Mol. Sci.* 19: 10.3390/ijms19113649, 2018.
283. Uchiyama, S, Tsukamoto, H, Yoshimura, S, and Tamaki, T. Relationship between oxidative stress in muscle tissue and weight-lifting-induced muscle damage. *Pflugers Arch.* 452: 109-116, 2006.
284. Usher-Smith, JA, Fraser, JA, Bailey, PS, Griffin, JL, and Huang, CL. The influence of intracellular lactate and H⁺ on cell volume in amphibian skeletal muscle. *J. Physiol.* 573: 799-818, 2006.
285. van der Pijl, R, Strom, J, Conijn, S, Lindqvist, J, Labeit, S, Granzier, H, and Ottenheijm, C. Titin-based mechanosensing modulates muscle hypertrophy. *J. Cachexia Sarcopenia Muscle* 9: 947-961, 2018.
286. van Wessel, T, de Haan, A, van der Laarse, WJ, and Jaspers, RT. The muscle fiber type-fiber size paradox: hypertrophy or oxidative metabolism? *Eur. J. Appl. Physiol.* 110: 665-694, 2010.
287. Vandenburg, HH, Hatfaludy, S, Sohar, I, and Shansky, J. Stretch-induced prostaglandins and protein turnover in cultured skeletal muscle. *Am. J. Physiol.* 259: C232-40, 1990.
288. Vane, JR, and Botting, RM. Anti-inflammatory drugs and their mechanism of action. *Inflamm. Res.* 47 Suppl 2: S78-87, 1998.
289. Vierck, J, O'Reilly, B, Hossner, K, Antonio, J, Byrne, K, Bucci, L, and Dodson, M. Satellite cell regulation following myotrauma caused by resistance exercise. *Cell Biol. Int.* 24: 263-272, 2000.
290. Vijayan, K, Thompson, JL, Norenberg, KM, Fitts, RH, and Riley, DA. Fiber-type susceptibility to eccentric contraction-induced damage of hindlimb-unloaded rat AL muscles. *J. Appl. Physiol.* 90: 770-776, 2001.
291. Viru, M, Jansson, E, Viru, A, and Sundberg, CJ. Effect of restricted blood flow on exercise-induced hormone changes in healthy men. *Eur. J. Appl. Physiol. Occup. Physiol.* 77: 517-522, 1998.
292. Vissing, K, McGee, S, Farup, J, Kjolhede, T, Vendelbo, M, and Jessen, N. Differentiated mTOR but not AMPK signaling after strength vs endurance exercise in training-accustomed individuals. *Scand. J. Med. Sci. Sports* 23: 355-366, 2013.
293. Vollestad, NK, Vaage, O, and Hermansen, L. Muscle glycogen depletion patterns in type I and subgroups of type II fibres during prolonged severe exercise in man. *Acta Physiol. Scand.* 122: 433-441, 1984.
294. Wackerhage, H, Schoenfeld, BJ, Hamilton, DL, Lehti, M, and Hulmi, JJ. Stimuli and sensors that initiate skeletal muscle hypertrophy following resistance exercise. *J. Appl. Physiol.* (1985) 126: 30-43, 2019.
295. Walton, RG, Kosmac, K, Mula, J, Fry, CS, Peck, BD, Groshong, JS, Finlin, BS, Zhu, B, Kern, PA, and Peterson, CA. Human skeletal muscle macrophages increase following cycle training and are associated with adaptations that may facilitate growth. *Sci. Rep.* 9: 969-018-37187-1, 2019.
296. Wang, XD, Kawano, F, Matsuoka, Y, Fukunaga, K, Terada, M, Sudoh, M, Ishihara, A, and Ohira, Y. Mechanical load-dependent regulation of satellite cell and fiber size in rat soleus muscle. *Am. J. Physiol., Cell Physiol.* 290: 981-989, 2006.
297. Warner, DC, Schnepf, G, Barrett, MS, Dian, D, and Swigonski, NL. Prevalence, attitudes, and behaviors related to the use of nonsteroidal anti-inflammatory drugs (NSAIDs) in student athletes. *J. Adolesc. Health* 30: 150-153, 2002.
298. Wernbom, M, Jarrebring, R, Andreasson, MA, and Augustsson, J. Acute effects of blood flow restriction on muscle activity and endurance during fatiguing dynamic knee extensions at low load. *J. Strength Cond Res.* 23: 2389-2395, 2009.
299. Wernbom, M, Paulsen, G, Nilsen, TS, Hisdal, J, and Raastad, T. Contractile function and sarcolemmal permeability after acute low-load resistance exercise with blood flow restriction. *Eur. J. Appl. Physiol.* 112: 2051-2063, 2012.
300. Wernig, A, Irintchev, A, and Weisshaupt, P. Muscle injury, cross-sectional area and fibre type distribution in mouse soleus after intermittent wheel-running. *J. Physiol.* 428: 639-652, 1990.
301. White, JP. Control of skeletal muscle cell growth and size through adhesion GPCRs. In: *Adhesion G Protein-coupled Receptors*. Anonymous New York, NY: Springer, 2016. pp. 299-308.
302. Widegren, U, Ryder, JW, and Zierath, JR. Mitogen-activated protein kinase signal transduction in skeletal muscle: effects of exercise and muscle contraction. *Acta Physiol. Scand.* 172: 227-238, 2001.
303. Willkomm, L, Schubert, S, Jung, R, Elsen, M, Borde, J, Gehlert, S, Suhr, F, and Bloch, W. Lactate regulates myogenesis in C2C12 myoblasts in vitro. *Stem Cell. Res.* 12: 742-753, 2014.
304. Wilson, JM, Lowery, RP, Joy, JM, Loenneke, JP, and Naimo, MA. Practical Blood Flow Restriction Training Increases Acute Determinants of Hypertrophy Without Increasing Indices of Muscle Damage. *J. Strength Cond Res.*, 2013.
305. Winter, JN, Jefferson, LS, and Kimball, SR. ERK and Akt signaling pathways function through parallel mechanisms to promote mTORC1 signaling. *Am. J. Physiol. Cell. Physiol.* 300: C1172-80, 2011.
306. Yasuda, T, Abe, T, Sato, Y, Midorikawa, T, Kearns, CF, Inoue, K, Ryushi, T, and Ishii, N. Muscle fiber cross-sectional area is increased after two weeks of twice daily KAATSU-resistance training. *Int J KAATSU Train Res* 1: 65-70, 2005.
307. Zanchi, NE, and Lancha, AH, Jr. Mechanical stimuli of skeletal muscle: implications on mTOR/p70s6k and protein synthesis. *Eur. J. Appl. Physiol.* 102: 253-263, 2008.

308. Zou, K, Meador, BM, Johnson, B, Huntsman, HD, Mahmassani, Z, Valero, MC, Huey, KA, and Boppart, MD. The $\alpha(7)\beta(1)$ -integrin increases muscle hypertrophy following multiple bouts of eccentric exercise. *J. Appl. Physiol.* 111: 1134-1141, 2011.
- ### Chapter 3
1. Aagaard, P, Andersen, JL, Dyhre-Poulsen, P, Leffers, AM, Wagner, A, Magnusson, SP, Halkjaer-Kristensen, J, and Simonsen, EB. A mechanism for increased contractile strength of human pennate muscle in response to strength training: changes in muscle architecture. *J. Physiol.* 534: 613-623, 2001.
 2. Abe, T, Loenneke, JP, and Thiebaud, RS. Morphological and functional relationships with ultrasound measured muscle thickness of the lower extremity: a brief review. *Ultrasound* 23: 166-173, 2015.
 3. Abe, T, Dankel, SJ, and Loenneke, JP. Body Fat Loss Automatically Reduces Lean Mass by Changing the Fat-Free Component of Adipose Tissue. *Obesity (Silver Spring)* 27: 357-358, 2019.
 4. Ahtiainen, JP, Hoffren, M, Hulmi, JJ, Pietikainen, M, Mero, AA, Avela, J, and Häkkinen, K. Panoramic ultrasonography is a valid method to measure changes in skeletal muscle cross-sectional area. *Eur. J. Appl. Physiol.* 108: 273-279, 2010.
 5. Antonio, J, Kenyon, M, Ellerbroek, A, Carson, C, Burgess, V, Tyler-Palmer, D, Mike, J, Roberts, J, Angeli, G, and and Peacock, C. Comparison of Dual-Energy X-Ray Absorptiometry (DXA) versus a Multi-frequency Bioelectrical Impedance (InBody 770) Device for Body Composition Assessment after a 4-Week Hypoenergetic Diet. *Journal of Functional Morphology and Kinesiology* 4: 23, 2019.
 6. Barber, L, Barrett, R, and Lichtwark, G. Validation of a free-hand 3D ultrasound system for morphological measures of the medial gastrocnemius muscle. *J. Biomech.* 42: 1313-1319, 2009.
 7. Beneke, R, Neuerburg, J, and Bohndorf, K. Muscle cross-section measurement by magnetic resonance imaging. *Eur. J. Appl. Physiol. Occup. Physiol.* 63: 424-429, 1991.
 8. Blazeovich, AJ, Gill, ND, and Zhou, S. Intra- and intermuscular variation in human quadriceps femoris architecture assessed in vivo. *J. Anat.* 209: 289-310, 2006.
 9. Brodie, D, Moscrip, V, and Hutcheon, R. Body composition measurement: a review of hydrodensitometry, anthropometry, and impedance methods. *Nutrition* 14: 296-310, 1998.
 10. Buchholz, AC, Bartok, C, and Schoeller, DA. The validity of bioelectrical impedance models in clinical populations. *Nutr. Clin. Pract.* 19: 433-446, 2004.
 11. Buckinx, F, Landi, F, Cesari, M, Fielding, RA, Visser, M, Engelke, K, Maggi, S, Dennison, E, Al-Daghri, NM, Allepaerts, S, Bauer, J, Bautmans, I, Brandi, ML, Bruyere, O, Cederholm, T, Cerreta, F, Cherubini, A, Cooper, C, Cruz-Jentoft, A, McCloskey, E, Dawson-Hughes, B, Kaufman, JM, Laslop, A, Petermans, J, Reginster, JY, Rizzoli, R, Robinson, S, Rolland, Y, Rueda, R, Vellas, B, and Kanis, JA. Pitfalls in the measurement of muscle mass: a need for a reference standard. *J. Cachexia Sarcopenia Muscle* 9: 269-278, 2018.
 12. Bunt, JC, Lohman, TG, and Boileau, RA. Impact of total body water fluctuations on estimation of body fat from body density. *Med. Sci. Sports Exerc.* 21: 96-100, 1989.
 13. Clark, RV, Walker, AC, O'Connor-Semmes, RL, Leonard, MS, Miller, RR, Stimpson, SA, Turner, SM, Ravussin, E, Cefalu, WT, Hellerstein, MK, and Evans, WJ. Total body skeletal muscle mass: estimation by creatine (methyl-d3) dilution in humans. *J. Appl. Physiol.* (1985) 116: 1605-1613, 2014.
 14. Clark, RV, Walker, AC, Miller, RR, O'Connor-Semmes, RL, Ravussin, E, and Cefalu, WT. Creatine (methyl-d3) dilution in urine for estimation of total body skeletal muscle mass: accuracy and variability vs. MRI and DXA. *J. Appl. Physiol.* (1985) 124: 1-9, 2018.
 15. Cumberledge, EA, Myers, C, Venditti, JJ, Dixon, CB, and Andreacci, JL. The Effect of the Menstrual Cycle on Body Composition Determined by Contact-Electrode Bioelectrical Impedance Analyzers. *Int. J. Exerc. Sci.* 11: 625-632, 2018.
 16. Cyrino, ES, Okano, AH, Glaner, ME, Romanzini, M, Gobbo, LA, Makoski, A, Bruna, N, Melo, JC, and Tassi, GN. Impact of the use of different skinfold calipers for the analysis of the body composition. *Revista Brasileira de Medicina do Esporte* 9: 150-153, 2003.
 17. Dahmane, R, Djordjevic, S, Simunic, B, and Valencic, V. Spatial fiber type distribution in normal human muscle. Histological and tensiomyographical evaluation. *J. Biomech.* 38: 2451-2459, 2005.
 18. DeFreitas, JM, Beck, TW, Stock, MS, Dillon, MA, Sherk, VD, Stout, JR, and Cramer, JT. A comparison of techniques for estimating training-induced changes in muscle cross-sectional area. *J. Strength Cond Res.* 24: 2383-2389, 2010.
 19. Delisle-Houde, P, Reid, RER, Insogna, JA, Prokop, NW, Buchan, TA, Fontaine, SL, and Andersen, RE. Comparing DXA and Air Displacement Plethysmography to Assess Body Composition of Male Collegiate Hockey Players. *J. Strength Cond Res.* 33: 474-478, 2019.
 20. Delmonico, MJ, Kostek, MC, Johns, J, Hurley, BF, and Conway, JM. Can dual energy X-ray absorptiometry provide a valid assessment of changes in thigh muscle mass with strength training in older adults? *Eur. J. Clin. Nutr.* 62: 1372-1378, 2008.
 21. Eng, CM, Abrams, GD, Smallwood, LR, Lieber, RL, and Ward, SR. Muscle geometry affects accuracy of forearm volume determination by magnetic resonance imaging (MRI). *J. Biomech.* 40: 3261-3266, 2007.
 22. Engstrom, CM, Loeb, GE, Reid, JG, Forrest, WJ, and Avruch, L. Morphometry of the human thigh muscles. A comparison between anatomical sections and computer tomographic and magnetic resonance images. *J. Anat.* 176: 139-156, 1991.
 23. Esco, MR, Snarr, RL, Leatherwood, MD, Chamberlain, NA, Redding, ML, Flatt, AA, Moon, JR, and Williford, HN. Comparison of total and segmental body composition using DXA and multifrequency bioimpedance in collegiate female athletes. *J. Strength Cond Res.* 29: 918-925, 2015.
 24. Esmarck, B, Andersen, JL, Olsen, S, Richter, EA, Mizuno, M, and Kjaer, M. Timing of postexercise protein intake is important for muscle hypertrophy with resistance training in elderly humans. *J. Physiol.* 535: 301-311, 2001.
 25. Evans, EM, Saunders, MJ, Spano, MA, Arngimsson, SA, Lewis, RD, and Cureton, KJ. Body-composition changes with diet and exercise in obese women: a comparison of estimates from clinical methods and a 4-component model. *Am. J. Clin. Nutr.* 70: 5-12, 1999.
 26. Faria, SL, Faria, OP, Cardeal, MD, and Ito, MK. Validation study of multi-frequency bioelectrical impedance with dual-energy X-ray absorptiometry among obese patients. *Obes. Surg.* 24: 1476-1480, 2014.
 27. Farup, J, Kjolhede, T, Sorensen, H, Dalgas, U, Moller, AB, Vestergaard, PE, Ringgaard, S, Bojsen-Moller, J, and Vissing, K. Muscle morphological and strength adaptations to endurance vs. resistance training. *J. Strength Cond Res.* 26: 398-407, 2012.
 28. Fields, DA, Higgins, PB, and Hunter, GR. Assessment of body composition by air-displacement plethysmography:

- influence of body temperature and moisture. *Dyn. Med.* 3: 3-5918-3-3, 2004.
29. Fields, DA, Higgins, PB, and Radley, D. Air-displacement plethysmography: here to stay. *Curr. Opin. Clin. Nutr. Metab. Care* 8: 624-629, 2005.
 30. Fosbol, MO, and Zerahn, B. Contemporary methods of body composition measurement. *Clin. Physiol. Funct. Imaging* 35: 81-97, 2015.
 31. Franchi, MV, Atherton, PJ, Reeves, ND, Fluck, M, Williams, J, Mitchell, WK, Selby, A, Beltran Valls, RM, and Narici, MV. Architectural, functional and molecular responses to concentric and eccentric loading in human skeletal muscle. *Acta Physiol. (Oxf)* 210: 642-654, 2014.
 32. Franchi, MV, Longo, S, Mallinson, J, Quinlan, JJ, Taylor, T, Greenhaff, PL, and Narici, MV. Muscle thickness correlates to muscle cross-sectional area in the assessment of strength training-induced hypertrophy. *Scand. J. Med. Sci. Sports* 28: 846-853, 2018.
 33. Franchi, MV, Raiteri, BJ, Longo, S, Sinha, S, Narici, MV, and Csapo, R. Muscle Architecture Assessment: Strengths, Shortcomings and New Frontiers of in Vivo Imaging Techniques. *Ultrasound Med. Biol.* 44: 2492-2504, 2018.
 34. Frisard, MI, Greenway, FL, and Delany, JP. Comparison of methods to assess body composition changes during a period of weight loss. *Obes. Res.* 13: 845-854, 2005.
 35. Frontera, WR, Meredith, CN, O'Reilly, KP, Knuttgen, HG, and Evans, WJ. Strength conditioning in older men: skeletal muscle hypertrophy and improved function. *J. Appl. Physiol.* (1985) 64: 1038-1044, 1988.
 36. Frontera, WR, Meredith, CN, O'Reilly, KP, Knuttgen, HG, and Evans, WJ. Strength conditioning in older men: skeletal muscle hypertrophy and improved function. *J. Appl. Physiol.* (1985) 64: 1038-1044, 1988.
 37. Gaba, A, Kapus, O, Cuberek, R, and Botek, M. Comparison of multi- and single-frequency bioelectrical impedance analysis with dual-energy X-ray absorptiometry for assessment of body composition in post-menopausal women: effects of body mass index and accelerometer-determined physical activity. *J. Hum. Nutr. Diet.* 28: 390-400, 2015.
 38. Geisler, C, Pourhassan, M, Braun, W, Schweitzer, L, and Muller, MJ. The prediction of total skeletal muscle mass in a Caucasian population - comparison of Magnetic resonance imaging (MRI) and Dual-energy X-ray absorptiometry (DXA). *Clin. Physiol. Funct. Imaging* 37: 168-172, 2017.
 39. Gleichauf, CN, and Roe, DA. The menstrual cycle's effect on the reliability of bioimpedance measurements for assessing body composition. *Am. J. Clin. Nutr.* 50: 903-907, 1989.
 40. Grgic, J, and Schoenfeld, BJ. Are the Hypertrophic Adaptations to High and Low-Load Resistance Training Muscle Fiber Type Specific? *Front. Physiol.* 9: 402, 2018.
 41. Haun, CT, Vann, CG, Osburn, SC, Mumford, PW, Roberson, PA, Romero, MA, Fox, CD, Johnson, CA, Parry, HA, Kavazis, AN, Moon, JR, Badisa, VLD, Mwashote, BM, Ibeanusi, V, Young, KC, and Roberts, MD. Muscle fiber hypertrophy in response to 6 weeks of high-volume resistance training in trained young men is largely attributed to sarcoplasmic hypertrophy. *PLoS One* 14: e0215267, 2019.
 42. Haun, CT, Vann, CG, Roberts, BM, Vigotsky, AD, Schoenfeld, BJ, and Roberts, MD. A Critical Evaluation of the Biological Construct Skeletal Muscle Hypertrophy: Size Matters but So Does the Measurement. *Front. Physiol.* 10: 247, 2019.
 43. Henriksson-Larsén, K. Distribution, total number and size of different types of fibres in male and female skeletal muscles: an enzyme histochemical study of whole muscle cross-sections. *Umeå universitet*, 1984.
 44. Heymsfield, SB, Arteaga, C, McManus, C, Smith, J, and Mof-fitt, S. Measurement of muscle mass in humans: validity of the 24-hour urinary creatinine method. *Am. J. Clin. Nutr.* 37: 478-494, 1983.
 45. Heyward, VH. Evaluation of body composition. Current issues. *Sports Med.* 22: 146-156, 1996.
 46. Jensen, B, Moritoyo, T, Kaufer-Horwitz, M, Peine, S, Norman, K, Maisch, MJ, Matsumoto, A, Masui, Y, Velazquez-Gonzalez, A, Dominguez-Garcia, J, Fonz-Enriquez, E, Salgado-Moctezuma, SG, and Bosy-Westphal, A. Ethnic differences in fat and muscle mass and their implication for interpretation of bioelectrical impedance vector analysis. *Appl. Physiol. Nutr. Metab.* 44: 619-626, 2019.
 47. Kendall, KL, Fukuda, DH, Hyde, PN, Smith-Ryan, AE, Moon, JR, and Stout, JR. Estimating fat-free mass in elite-level male rowers: a four-compartment model validation of laboratory and field methods. *J. Sports Sci.* 35: 624-633, 2017.
 48. Kilduff, LP, Lewis, S, Kingsley, MI, Owen, NJ, and Dietz, RE. Reliability and detecting change following short-term creatine supplementation: comparison of two-component body composition methods. *J. Strength Cond Res.* 21: 378-384, 2007.
 49. Kim, J, Heshka, S, Gallagher, D, Kotler, DP, Mayer, L, Albu, J, Shen, W, Freda, PU, and Heymsfield, SB. Intermuscular adipose tissue-free skeletal muscle mass: estimation by dual-energy X-ray absorptiometry in adults. *J. Appl. Physiol.* (1985) 97: 655-660, 2004.
 50. Kyle, UG, Bosaeus, I, De Lorenzo, AD, Deurenberg, P, Elia, M, Gomez, JM, Heitmann, BL, Kent-Smith, L, Melchior, JC, Pirlich, M, Scharfetter, H, Schols, AM, Pichard, C, and Composition of the ESPEN Working Group. Bioelectrical impedance analysis--part I: review of principles and methods. *Clin. Nutr.* 23: 1226-1243, 2004.
 51. Levine, JA, Abboud, L, Barry, M, Reed, JE, Sheedy, PF, and Jensen, MD. Measuring leg muscle and fat mass in humans: comparison of CT and dual-energy X-ray absorptiometry. *J. Appl. Physiol.* 88: 452-456, 2000.
 52. Loenneke, JP, Dankel, SJ, Bell, ZW, Spitz, RW, Abe, T, and Yasuda, T. Ultrasound and MRI measured changes in muscle mass gives different estimates but similar conclusions: a Bayesian approach. *Eur. J. Clin. Nutr.* 73: 1203-1205, 2019.
 53. Lohman, T, Wang, Z, and Going, SB. Human Body Composition. Champaign, IL; Human Kinetics, 2005.
 54. Lukaski, HC. Methods for the assessment of human body composition: traditional and new. *Am. J. Clin. Nutr.* 46: 537-556, 1987.
 55. Mahon, AK, Flynn, MG, Iglay, HB, Stewart, LK, Johnson, CA, McFarlin, BK, and Campbell, WW. Measurement of body composition changes with weight loss in postmenopausal women: comparison of methods. *J. Nutr. Health Aging* 11: 203-213, 2007.
 56. Matta, T, Simao, R, de Salles, BF, Spinetti, J, and Oliveira, LF. Strength training's chronic effects on muscle architecture parameters of different arm sites. *J. Strength Cond Res.* 25: 1711-1717, 2011.
 57. Matthie, JR. Bioimpedance measurements of human body composition: critical analysis and outlook. *Expert Rev. Med. Devices* 5: 239-261, 2008.
 58. Mazonakis, M, and Damilakis, J. Computed tomography: What and how does it measure? *Eur. J. Radiol.* 85: 1499-1504, 2016.

59. McCall, GE, Byrnes, WC, Dickinson, A, Pattany, PM, and Fleck, SJ. Muscle fiber hypertrophy, hyperplasia, and capillary density in college men after resistance training. *J. Appl. Physiol.* (1985) 81: 2004-2012, 1996.
60. McKee, JE, and Cameron, N. Bioelectrical impedance changes during the menstrual cycle. *Am. J. Hum. Biol.* 9: 155-161, 1997.
61. McLester, CN, Nickerson, BS, Kliszczewicz, BM, and McLester, JR. Reliability and Agreement of Various InBody Body Composition Analyzers as Compared to Dual-Energy X-Ray Absorptiometry in Healthy Men and Women. *J. Clin. Densitom.* , 2018.
62. Mitsiopoulos, N, Baumgartner, RN, Heymsfield, SB, Lyons, W, Gallagher, D, and Ross, R. Cadaver validation of skeletal muscle measurement by magnetic resonance imaging and computerized tomography. *J. Appl. Physiol.* (1985) 85: 115-122, 1998.
63. Miyatani, M, Kanehisa, H, Ito, M, Kawakami, Y, and Fukunaga, T. The accuracy of volume estimates using ultrasound muscle thickness measurements in different muscle groups. *Eur. J. Appl. Physiol.* 91: 264-272, 2004.
64. Moon, JR. Body composition in athletes and sports nutrition: an examination of the bioimpedance analysis technique. *Eur. J. Clin. Nutr.* 67 Suppl 1: S54-9, 2013.
65. Nakajima, T, Koide, S, Yasuda, T, Hasegawa, T, Yamasoba, T, Obi, S, Toyoda, S, Nakamura, E, Inoue, T, Poole, DC, and Kano, Y. Muscle hypertrophy following blood flow-restricted, low-force isometric electrical stimulation in rat tibialis anterior: role for muscle hypoxia. *J. Appl. Physiol.* (1985) 125: 134-145, 2018.
66. Narici, MV, Roi, GS, Landoni, L, Minetti, AE, and Cerretelli, P. Changes in force, cross-sectional area and neural activation during strength training and detraining of the human quadriceps. *Eur. J. Appl. Physiol. Occup. Physiol.* 59: 310-319, 1989.
67. Narici, MV, Hoppeler, H, Kayser, B, Landoni, L, Claassen, H, Gavardi, C, Conti, M, and Cerretelli, P. Human quadriceps cross-sectional area, torque and neural activation during 6 months strength training. *Acta Physiol. Scand.* 157: 175-186, 1996.
68. Nilwik, R, Snijders, T, Leenders, M, Groen, BB, van Kranenburg, J, Verdijk, LB, and van Loon, LJ. The decline in skeletal muscle mass with aging is mainly attributed to a reduction in type II muscle fiber size. *Exp. Gerontol.* 48: 492-498, 2013.
69. Noorkoiv, M, Nosaka, K, and Blazevich, AJ. Assessment of quadriceps muscle cross-sectional area by ultrasound extended-field-of-view imaging. *Eur. J. Appl. Physiol.* 109: 631-639, 2010.
70. Oppliger, RA, Nielsen, DH, Shetler, AC, Crowley, ET, and Albright, JP. Body composition of collegiate football players: bioelectrical impedance and skinfolds compared to hydrostatic weighing. *J. Orthop. Sports Phys. Ther.* 15: 187-192, 1992.
71. Peeters, MW, and Claessens, AL. Effect of deviating clothing schemes on the accuracy of body composition measurements by air-displacement plethysmography. *International Journal of Body Composition Research* 7, 2009.
72. Peeters, MW. Subject positioning in the BOD POD(R) only marginally affects measurement of body volume and estimation of percent body fat in young adult men. *PLoS One* 7: e32722, 2012.
73. Prado, CM, and Heymsfield, SB. Lean tissue imaging: a new era for nutritional assessment and intervention. *JPEN J. Parenter. Enteral Nutr.* 38: 940-953, 2014.
74. Ritz, P, Salle, A, Audran, M, and Rohmer, V. Comparison of different methods to assess body composition of weight loss in obese and diabetic patients. *Diabetes Res. Clin. Pract.* 77: 405-411, 2007.
75. Roman, WJ, Fleckenstein, J, Stray-Gundersen, J, Alway, SE, Peshock, R, and Gonyea, WJ. Adaptations in the elbow flexors of elderly males after heavy-resistance training. *J. Appl. Physiol.* (1985) 74: 750-754, 1993.
76. Schmidt, PK, and Carter, JE. Static and dynamic differences among five types of skinfold calipers. *Hum. Biol.* 62: 369-388, 1990.
77. Schoenfeld, BJ, Nickerson, BS, Wilborn, CD, Urbina, SL, Hayward, SB, Krieger, J, Aragon, AA, and Tinsley, GM. Comparison of Multifrequency Bioelectrical Impedance vs. Dual-Energy X-ray Absorptiometry for Assessing Body Composition Changes After Participation in a 10-Week Resistance Training Program. *J. Strength Cond Res.* , 2018.
78. Scott, JM, Martin, DS, Ploutz-Snyder, R, Matz, T, Caine, T, Downs, M, Hackney, K, Buxton, R, Ryder, JW, and Ploutz-Snyder, L. Panoramic ultrasound: a novel and valid tool for monitoring change in muscle mass. *J. Cachexia Sarcopenia Muscle* 8: 475-481, 2017.
79. Scott, SH, Engstrom, CM, and Loeb, GE. Morphometry of human thigh muscles. Determination of fascicle architecture by magnetic resonance imaging. *J. Anat.* 182 (Pt 2): 249-257, 1993.
80. Shaw, G, Kerr, A. Non-imaging method: Air displacement plethysmography (bod pod). (pp. 87-99). In: *Best Practice Protocols for Physique Assessment in Sport*. Anonymous Singapore: Springer, 2018. pp. 87-99.
81. Sillanpaa, E, Häkkinen, A, and Häkkinen, K. Body composition changes by DXA, BIA and skinfolds during exercise training in women. *Eur. J. Appl. Physiol.* 113: 2331-2341, 2013.
82. Silva, AM, Fields, DA, Quiterio, AL, and Sardinha, LB. Are skinfold-based models accurate and suitable for assessing changes in body composition in highly trained athletes? *J. Strength Cond Res.* 23: 1688-1696, 2009.
83. Stimpson, SA, Turner, SM, Clifton, LG, Poole, JC, Mohammed, HA, Shearer, TW, Waitt, GM, Hagerty, LL, Remlinger, KS, Hellerstein, MK, and Evans, WJ. Total-body creatine pool size and skeletal muscle mass determination by creatine-(methyl-D3) dilution in rats. *J. Appl. Physiol.* (1985) 112: 1940-1948, 2012.
84. Stimpson, SA, Leonard, MS, Clifton, LG, Poole, JC, Turner, SM, Shearer, TW, Remlinger, KS, Clark, RV, Hellerstein, MK, and Evans, WJ. Longitudinal changes in total body creatine pool size and skeletal muscle mass using the D3-creatine dilution method. *J. Cachexia Sarcopenia Muscle* , 2013.
85. Tavoian, D, Ampomah, K, Amano, S, Law, TD, and Clark, BC. Changes in DXA-derived lean mass and MRI-derived cross-sectional area of the thigh are modestly associated. *Sci. Rep.* 9: 10028-019-46428-w, 2019.
86. Thomson, R, Brinkworth, GD, Buckley, JD, Noakes, M, and Clifton, PM. Good agreement between bioelectrical impedance and dual-energy X-ray absorptiometry for estimating changes in body composition during weight loss in overweight young women. *Clin. Nutr.* 26: 771-777, 2007.
87. Tingart, MJ, Apreleva, M, Lehtinen, JT, Capell, B, Palmer, WE, and Warner, JJ. Magnetic resonance imaging in quantitative analysis of rotator cuff muscle volume. *Clin. Orthop. Relat. Res.* (415):104-110. doi: 104-110, 2003.
88. Tinsley, GM, Morales, E, Forsse, JS, and Grandjean, PW. Impact of Acute Dietary Manipulations on DXA and BIA

- Body Composition Estimates. *Med. Sci. Sports Exerc.* 49: 823-832, 2017.
89. van Marken Lichtenbelt, WD, Hartgens, F, Vollaard, NB, Ebbing, S, and Kuipers, H. Body composition changes in bodybuilders: a method comparison. *Med. Sci. Sports Exerc.* 36: 490-497, 2004.
 90. Verdijk, LB, Gleeson, BG, Jonkers, RA, Meijer, K, Savelberg, HH, Dendale, P, and van Loon, LJ. Skeletal muscle hypertrophy following resistance training is accompanied by a fiber type-specific increase in satellite cell content in elderly men. *J. Gerontol. A Biol. Sci. Med. Sci.* 64: 332-339, 2009.
 91. Vissing, K, Brink, M, Lonbro, S, Sorensen, H, Overgaard, K, Danborg, K, Mortensen, J, Elstrom, O, Rosenhoj, N, Ringgaard, S, Andersen, JL, and Aagaard, P. Muscle adaptations to plyometric vs. resistance training in untrained young men. *J Strength Cond Res* 22: 1799-1810, 2008.
 92. Volgyi, E, Tylavsky, FA, Lyytikainen, A, Suominen, H, Alen, M, and Cheng, S. Assessing body composition with DXA and bioimpedance: effects of obesity, physical activity, and age. *Obesity (Silver Spring)* 16: 700-705, 2008.
 93. Wang, Z, Pi-Sunyer, FX, Kotler, DP, Wielopolski, L, Withers, RT, Pierson, RN, Jr, and Heymsfield, SB. Multicomponent methods: evaluation of new and traditional soft tissue mineral models by in vivo neutron activation analysis. *Am. J. Clin. Nutr.* 76: 968-974, 2002.
 94. Ward, LC. Segmental bioelectrical impedance analysis: an update. *Curr. Opin. Clin. Nutr. Metab. Care* 15: 424-429, 2012.
 95. Wells, JC, and Fewtrell, MS. Measuring body composition. *Arch. Dis. Child.* 91: 612-617, 2006.
 96. Weyers, AM, Mazzetti, SA, Love, DM, Gomez, AL, Kraemer, WJ, and Volek, JS. Comparison of methods for assessing body composition changes during weight loss. *Med. Sci. Sports Exerc.* 34: 497-502, 2002.
 97. Williams, JE, Wells, JC, Wilson, CM, Haroun, D, Lucas, A, and Fewtrell, MS. Evaluation of Lunar Prodigy dual-energy X-ray absorptiometry for assessing body composition in healthy persons and patients by comparison with the criterion 4-component model. *Am. J. Clin. Nutr.* 83: 1047-1054, 2006.
 98. Yasuda, T, Abe, T, Sato, Y, Midorikawa, T, Kearns, CE, Inoue, K, Ryushi, T, and Ishii, N. Muscle fiber cross-sectional area is increased after two weeks of twice daily KAATSU-resistance training. *Int J KAATSU Train Res* 1: 65-70, 2005.
 - resistance training for healthy adults. *Med. Sci. Sports Exerc.* 41: 687-708, 2009.
 6. Amirthalingam, T, Mavros, Y, Wilson, GC, Clarke, JL, Mitchell, L, and Hackett, DA. Effects of a Modified German Volume Training Program on Muscular Hypertrophy and Strength. *J. Strength Cond Res.* , 2016.
 7. Ampomah, K, Amano, S, Wages, NP, Volz, L, Clift, R, Ludin, AFM, Nakazawa, M, Law, TD, Manini, TM, Thomas, JS, Russ, DW, and Clark, BC. Blood Flow-restricted Exercise Does Not Induce a Cross-Transfer of Effect: A Randomized Controlled Trial. *Med. Sci. Sports Exerc.* 51: 1817-1827, 2019.
 8. Arazi, H, and Asadi, A. Effects of 8 Weeks Equal-Volume Resistance Training with Different Workout Frequency on Maximal Strength, Endurance and Body Composition. *International Journal of Sports Science and Engineering* 5: 112-118, 2011.
 9. Assis-Pereira, PE, Motoyama, YL, Esteves, GJ, Quinelato, WC, Botter, L, Tanaka, KH, and Azevedo, P. Resistance training with slow speed of movement is better for hypertrophy and muscle strength gains than fast speed of movement. *Int J Appl Exerc Physiol* 5: 37-43, 2016.
 10. Avelar, A, Ribeiro, AS, Nunes, JP, Schoenfeld, BJ, Papst, RR, Trindade, MCC, Bottaro, M, and Cyrino, ES. Effects of order of resistance training exercises on muscle hypertrophy in young adult men. *Appl. Physiol. Nutr. Metab.* 44: 420-424, 2019.
 11. Balsalobre, C, Santos-Concejero, J, Baz, E, and Schoenfeld, BJ. The effects of exercise variation in muscle thickness, maximal strength and motivation in resistance trained men. *Plos One* , 2019.
 12. Bamman, MM, Shipp, JR, Jiang, J, Gower, BA, Hunter, GR, Goodman, A, McLafferty, CL, Jr, and Urban, RJ. Mechanical load increases muscle IGF-I and androgen receptor mRNA concentrations in humans. *Am. J. Physiol. Endocrinol. Metab.* 280: E383-90, 2001.
 13. Barbalho, M, Coswig, VS, Steele, J, Fisher, JP, Giessing, J, and Gentil, P. Evidence of a Ceiling Effect for Training Volume in Muscle Hypertrophy and Strength in Trained Men - Less is More? *Int. J. Sports Physiol. Perform.* : 1-23, 2019.
 14. Barbalho, M, Coswig, VS, Steele, J, Fisher, JP, Paoli, A, and Gentil, P. Evidence for an Upper Threshold for Resistance Training Volume in Trained Women. *Med. Sci. Sports Exerc.* 51: 515-522, 2019.
 15. Barcelos, C, Damas, F, Nobrega, SR, Ugrinowitsch, C, Lixandrao, ME, Marcelino Eder Dos Santos, L, and Libardi, CA. High-frequency resistance training does not promote greater muscular adaptations compared to low frequencies in young untrained men. *Eur. J. Sport. Sci.* : 1-6, 2018.
 16. Baz-Valle, E, Fontes-Villalba, M, and Santos-Concejero, J. Total Number of Sets as a Training Volume Quantification Method for Muscle Hypertrophy: A Systematic Review. *J. Strength Cond Res.* , 2018.
 17. Behm, DG. Neuromuscular implications and applications of resistance training. *J Strength Cond Res* 9: 264-274, 1995.
 18. Belcastro, AN, Shewchuk, LD, and Raj, DA. Exercise-induced muscle injury: a calpain hypothesis. *Mol. Cell. Biochem.* 179: 135-145, 1998.
 19. Ben-Sira, D, Ayalon, A, and Tavi, M. The effect of different types of strength training on concentric strength in women. *J Strength Cond Res.* 9: 143-148, 1995.
 20. Benton, MJ, Kasper, MJ, Raab, SA, Waggner, GT, and Swan, PD. Short-term effects of resistance training frequency on body composition and strength in middle-aged women. *J. Strength Cond Res.* 25: 3142-3149, 2011.

Chapter 4

1. Abdessemed, D, Duche, P, Hautier, C, Poumarat, G, and Bedu, M. Effect of recovery duration on muscular power and blood lactate during the bench press exercise. *Int. J. Sports Med.* 20: 368-373, 1999.
2. Ahtiainen, JP, Walker, S, Silvennoinen, M, Kyrolainen, H, Nindl, BC, Häkkinen, K, Nyman, K, Selanne, H, and Hulmi, JJ. Exercise type and volume alter signaling pathways regulating skeletal muscle glucose uptake and protein synthesis. *Eur. J. Appl. Physiol.* 115: 1835-1845, 2015.
3. Ahtiainen, JP, Pakarinen, A, Alen, M, Kraemer, WJ, and Häkkinen, K. Short vs. long rest period between the sets in hypertrophic resistance training: influence on muscle strength, size, and hormonal adaptations in trained men. *J Strength Cond Res* 19: 572-582, 2005.
4. Allen, DG, Whitehead, NP, and Yeung, EW. Mechanisms of stretch-induced muscle damage in normal and dystrophic muscle: role of ionic changes. *J. Physiol.* 567: 723-735, 2005.
5. American College of Sports Medicine. American College of Sports Medicine position stand. Progression models in

21. Bickel, CS, Slade, J, Mahoney, E, Haddad, F, Dudley, GA, and Adams, GR. Time course of molecular responses of human skeletal muscle to acute bouts of resistance exercise. *J. Appl. Physiol.* 98: 482-488, 2005.
22. Bjornsen, T, Wernbom, M, Kirketeig, A, Paulsen, G, Samnøy, L, Baekken, L, Cameron-Smith, D, Berntsen, S, and Raastad, T. Type 1 Muscle Fiber Hypertrophy after Blood Flow-restricted Training in Powerlifters. *Med. Sci. Sports Exerc.* 51: 288-298, 2019.
23. Bjornsen, T, Wernbom, M, Lovstad, A, Paulsen, G, D'Souza, RF, Cameron-Smith, D, Flesche, A, Hisdal, J, Berntsen, S, and Raastad, T. Delayed myonuclear addition, myofiber hypertrophy, and increases in strength with high-frequency low-load blood flow restricted training to volitional failure. *J. Appl. Physiol.* (1985) 126: 578-592, 2019.
24. Blazevich, AJ, Cannavan, D, Coleman, DR, and Horne, S. Influence of concentric and eccentric resistance training on architectural adaptation in human quadriceps muscles. *J. Appl. Physiol.* 103: 1565-1575, 2007.
25. Bloomquist, K, Langberg, H, Karlsen, S, Madsgaard, S, Boesen, M, and Raastad, T. Effect of range of motion in heavy load squatting on muscle and tendon adaptations. *Eur. J. Appl. Physiol.* 113: 2133-2142, 2013.
26. Bottaro, M, Veloso, J, Wagner, D, and Gentil, P. Resistance training for strength and muscle thickness: Effect of number of sets and muscle group trained. *Sci Sports* 26: 259-264, 2011.
27. Brigatto, FA, Braz, TV, Zanini, T, CDC, Germano, MD, Aoki, MS, Schoenfeld, BJ, Marchetti, PH, and Lopes, CR. Effect of Resistance Training Frequency on Neuromuscular Performance and Muscle Morphology after Eight Weeks in Trained Men. *J. Strength Cond Res.*, 2018.
28. Brown, JM, Solomon, C, and Paton, M. Further evidence of functional differentiation within biceps brachii. *Electromyogr. Clin. Neurophysiol.* 33: 301-309, 1993.
29. Burd, NA, Holwerda, AM, Selby, KC, West, DW, Staples, AW, Cain, NE, Cashaback, JG, Potvin, JR, Baker, SK, and Phillips, SM. Resistance exercise volume affects myofibrillar protein synthesis and anabolic signalling molecule phosphorylation in young men. *J. Physiol.* 588: 3119-3130, 2010.
30. Burd, NA, West, DW, Staples, AW, Atherton, PJ, Baker, JM, Moore, DR, Holwerda, AM, Parise, G, Rennie, MJ, Baker, SK, and Phillips, SM. Low-load high volume resistance exercise stimulates muscle protein synthesis more than high-load low volume resistance exercise in young men. *PLoS One* 5: e12033, 2010.
31. Burd, NA, West, DW, Moore, DR, Atherton, PJ, Staples, AW, Prior, T, Tang, JE, Rennie, MJ, Baker, SK, and Phillips, SM. Enhanced amino acid sensitivity of myofibrillar protein synthesis persists for up to 24 h after resistance exercise in young men. *J. Nutr.* 141: 568-573, 2011.
32. Burd, NA, Andrews, RJ, West, DW, Little, JP, Cochran, AJ, Hector, AJ, Cashaback, JG, Gibala, MJ, Potvin, JR, Baker, SK, and Phillips, SM. Muscle time under tension during resistance exercise stimulates differential muscle protein sub-fractional synthetic responses in men. *J. Physiol.* 590: 351-362, 2012.
33. Burd, NA, Mitchell, CJ, Churchward-Venne, TA, and Phillips, SM. Bigger weights may not beget bigger muscles: evidence from acute muscle protein synthetic responses after resistance exercise. *Appl. Physiol. Nutr. Metab.* 37: 551-554, 2012.
34. Buresh, R, Berg, K, and French, J. The effect of resistive exercise rest interval on hormonal response, strength, and hypertrophy with training. *J Strength Cond Res* 23: 62-71, 2009.
35. Cadore, EL, Izquierdo, M, Pinto, SS, Alberton, CL, Pinto, RS, Baroni, BM, Vaz, MA, Lanferdini, FJ, Radaelli, R, Gonzalez-Izal, M, Bottaro, M, and Krusel, LF. Neuromuscular adaptations to concurrent training in the elderly: effects of intrasession exercise sequence. *Age (Dordr)* 35: 891-903, 2013.
36. Calatayud, J, Vinstrop, J, Jakobsen, MD, Sundstrup, E, Brandt, M, Jay, K, Colado, JC, and Andersen, LL. Importance of mind-muscle connection during progressive resistance training. *Eur. J. Appl. Physiol.* 116: 527-533, 2016.
37. Calatayud, J, Vinstrop, J, Jakobsen, MD, Sundstrup, E, Colado, JC, and Andersen, LL. Influence of different attentional focus on EMG amplitude and contraction duration during the bench press at different speeds. *J. Sports Sci.* 36: 1162-1166, 2018.
38. Calder, AW, Chilibeck, PD, Webber, CE, and Sale, DG. Comparison of whole and split weight training routines in young women. *Can. J. Appl. Physiol.* 19: 185-199, 1994.
39. Campos, GER, Luecke, TJ, Wendeln, HK, Toma, K, Hagerman, FC, Murray, TE, Ragg, KE, Ratamess, NA, Kraemer, WJ, and Staron, RS. Muscular adaptations in response to three different resistance-training regimens: specificity of repetition maximum training zones. *Eur. J. Appl. Physiol.* 88: 50-60, 2002.
40. Candow, DG, and Burke, DG. Effect of short-term equal-volume resistance training with different workout frequency on muscle mass and strength in untrained men and women. *J. Strength Cond Res.* 21: 204-207, 2007.
41. Cannon, J, and Marino, FE. Early-phase neuromuscular adaptations to high- and low-volume resistance training in untrained young and older women. *J. Sports Sci.* 28: 1505-1514, 2010.
42. Cardozo, DC, DE Salles, BF, Mannarino, P, Vasconcelos, APS, Miranda, H, Willardson, JM, and Simao, R. The Effect of Exercise Order in Circuit Training on Muscular Strength and Functional Fitness in Older Women. *Int. J. Exerc. Sci.* 12: 657-665, 2019.
43. Carlson, L, Jonker, B, Westcott, WL, Steele, J, and Fisher, JP. Neither repetition duration nor number of muscle actions affect strength increases, body composition, muscle size, or fasted blood glucose in trained males and females. *Appl. Physiol. Nutr. Metab.* 44: 200-207, 2019.
44. Carneiro, NH, Ribeiro, AS, Nascimento, MA, Gobbo, LA, Schoenfeld, BJ, Achour Junior, A, Gobbi, S, Oliveira, AR, and Cyrino, ES. Effects of different resistance training frequencies on flexibility in older women. *Clin. Interv. Aging* 10: 531-538, 2015.
45. Carpinelli, RN. The size principle and a critical analysis of the unsubstantiated heavier-is-better recommendation for resistance training. *J Exerc Sci Fit* 6: 67-86, 2008.
46. Carroll, KM, Bazzyler, CD, Bernards, JR, Taber, CB, Stuart, CA, DeWeese, BH, Sato, K, and Stone, MH. Skeletal Muscle Fiber Adaptations Following Resistance Training Using Repetition Maximums or Relative Intensity. *Sports (Basel)* 7: 10.3390/sports7070169, 2019.
47. Chestnut, J, and Docherty, D. The effects of 4 and 10 repetition maximum weight-training protocols on neuromuscular adaptations in untrained men. *J Strength Cond Res* 13: 353-359, 1999.
48. Choi, J, Takahashi, H, and Itai, Y. The difference between effects of 'power-up type' and 'bulk-up type' strength training exercises: with special reference to muscle cross-sectional area. *Jpn J Phys Fitness Sports Med* 47: 119-129, 1998.
49. Claflin, DR, Larkin, LM, Cederna, PS, Horowitz, JF, Alexander, NB, Cole, NM, Galecki, AT, Chen, S, Nyquist, LV, Carlson,

- BM, Faulkner, JA, and Ashton-Miller, JA. Effects of high- and low-velocity resistance training on the contractile properties of skeletal muscle fibers from young and older humans. *J. Appl. Physiol.* (1985) 111: 1021-1030, 2011.
50. Colquhoun, RJ, Gai, CM, Aguilar, D, Bove, D, Dolan, J, Vargas, A, Couvillion, K, Jenkins, NDM, and Campbell, BI. Training Volume, Not Frequency, Indicative of Maximal Strength Adaptations to Resistance Training. *J. Strength Cond Res.* 32: 1207-1213, 2018.
51. Correa, CS, Teixeira, BC, Bittencourt, A, Lemos, L, Marques, NR, Radaelli, R, Kruger, RL, Reischak-Oliveira, A, and Pinto, RS. Effects of high and low volume of strength training on muscle strength, muscle volume and lipid profile in post-menopausal women. *Journal of Exercise Science and Fitness* 12: 62-67, 2014.
52. Cuthbertson, DJ, Babraj, J, Smith, K, Wilkes, E, Fedele, MJ, Esser, K, and Rennie, M. Anabolic signaling and protein synthesis in human skeletal muscle after dynamic shortening or lengthening exercise. *Am. J. Physiol. Endocrinol. Metab.* 290: E731-8, 2006.
53. da Silva, LXN, Teodoro, JL, Menger, E, Lopez, P, Grazioli, R, Farinha, J, Moraes, K, Bottaro, M, Pinto, RS, Izquierdo, M, and Cadore, EL. Repetitions to failure versus not to failure during concurrent training in healthy elderly men: A randomized clinical trial. *Exp. Gerontol.* 108: 18-27, 2018.
54. Damas, F, Phillips, SM, Libardi, CA, Vechin, FC, Lixandrao, ME, Jannig, PR, Costa, LA, Bacurau, AV, Snijders, T, Parise, G, Tricoli, V, Roschel, H, and Ugrinowitsch, C. Resistance training-induced changes in integrated myofibrillar protein synthesis are related to hypertrophy only after attenuation of muscle damage. *J. Physiol.* 594: 5209-5222, 2016.
55. Dankel, SJ, Mattocks, KT, Jessee, MB, Buckner, SL, Mouser, JG, Counts, BR, Laurentino, GC, and Loenneke, JP. Frequency: The Overlooked Resistance Training Variable for Inducing Muscle Hypertrophy? *Sports Med.* 47: 799-805, 2017.
56. de Souza, TPJr, Fleck, SJ, Simao, R, Dubas, JP, Pereira, B, de Brito Pacheco, EM, da Silva, AC, and de Oliveira, PR. Comparison between constant and decreasing rest intervals: influence on maximal strength and hypertrophy. *J. Strength Cond Res.* 24: 1843-1850, 2010.
57. Dimitrova, NA, and Dimitrov, GV. Interpretation of EMG changes with fatigue: facts, pitfalls, and fallacies. *J. Electromyogr. Kinesiol.* 13: 13-36, 2003.
58. Durand, RJ, Castracane, VD, Hollander, DB, Tryniecki, JL, Bamman, MM, O'Neal, S, Hebert, EP, and Kraemer, RR. Hormonal responses from concentric and eccentric muscle contractions. *Med. Sci. Sports Exerc.* 35: 937-943, 2003.
59. Ebben, WP, Feldmann, CR, Dayne, A, Mitsche, D, Alexander, P, and Knetzger, KJ. Muscle activation during lower body resistance training. *Int. J. Sports Med.* 30: 1-8, 2009.
60. Eliasson, J, Elfegoun, T, Nilsson, J, Kohnke, R, Ekblom, B, and Blomstrand, E. Maximal lengthening contractions increase p70 S6 kinase phosphorylation in human skeletal muscle in the absence of nutritional supply. *Am. J. Physiol. Endocrinol. Metab.* 291: 1197-1205, 2006.
61. English, AW, Wolf, SL, and Segal, RL. Compartmentalization of muscles and their motor nuclei: the partitioning hypothesis. *Phys. Ther.* 73: 857-867, 1993.
62. Enoka, RM. Physiological validation of the decomposition of surface EMG signals. *J. Electromyogr. Kinesiol.* 46: 70-83, 2019.
63. Escamilla, RE, Fleisig, GS, Zheng, N, Barrentine, SW, Wilk, KE, and Andrews, JR. Biomechanics of the knee during closed kinetic chain and open kinetic chain exercises. *Med. Sci. Sports Exerc.* 30: 556-569, 1998.
64. Farthing, JP, and Chilibeck, PD. The effects of eccentric and concentric training at different velocities on muscle hypertrophy. *Eur. J. Appl. Physiol.* 89: 578-586, 2003.
65. Farup, J, Rahbek, SK, Riis, S, Vendelbo, MH, Paoli, F, and Vissing, K. Influence of exercise contraction mode and protein supplementation on human skeletal muscle satellite cell content and muscle fiber growth. *J. Appl. Physiol.* (1985) 117: 898-909, 2014.
66. Farup, J, Rahbek, SK, Vendelbo, MH, Matzon, A, Hindhede, J, Bejder, A, Ringgard, S, and Vissing, K. Whey protein hydrolysate augments tendon and muscle hypertrophy independent of resistance exercise contraction mode. *Scand. J. Med. Sci. Sports* 24: 788-798, 2014.
67. Farup, J, de Paoli, F, Bjerg, K, Riis, S, Ringgard, S, and Vissing, K. Blood flow restricted and traditional resistance training performed to fatigue produce equal muscle hypertrophy. *Scand. J. Med. Sci. Sports* 25: 754-763, 2015.
68. Fernandez-Lezaun, E, Schumann, M, Makinen, T, Kyrolainen, H, and Walker, S. Effects of resistance training frequency on cardiorespiratory fitness in older men and women during intervention and follow-up. *Exp. Gerontol.* 95: 44-53, 2017.
69. Fink, JE, Schoenfeld, BJ, Kikuchi, N, and Nakazato, K. Acute and Long-term Responses to Different Rest Intervals in Low-load Resistance Training. *Int. J. Sports Med.* 38: 118-124, 2017.
70. Fisher, JP, Carlson, L, Steele, J, and Smith, D. The effects of pre-exhaustion, exercise order, and rest intervals in a full-body resistance training intervention. *Appl. Physiol. Nutr. Metab.* 39: 1265-1270, 2014.
71. Fonseca, RM, Roschel, H, Tricoli, V, de Souza, EO, Wilson, JM, Laurentino, GC, Aihara, AY, de Souza Leao, AR, and Ugrinowitsch, C. Changes in exercises are more effective than in loading schemes to improve muscle strength. *J. Strength Cond Res.* , 2014.
72. Franchi, MV, Atherton, PJ, Reeves, ND, Fluck, M, Williams, J, Mitchell, WK, Selby, A, Beltran Valls, RM, and Narici, MV. Architectural, functional and molecular responses to concentric and eccentric loading in human skeletal muscle. *Acta Physiol. (Oxf)* 210: 642-654, 2014.
73. Franco, CMC, Carneiro, MADS, Alves, LTH, Junior, GNO, de Sousa, JFR, and Orsatti, FL. Lower-Load is More Effective Than Higher-Load Resistance Training in Increasing Muscle Mass in Young Women. *J. Strength Cond Res.* 33 Suppl 1: S152-S158, 2019.
74. Fry, AC, and Kraemer, WJ. Resistance exercise overtraining and overreaching. Neuroendocrine responses. *Sports Med.* 23: 106-129, 1997.
75. Galvao, DA, and Taaffe, DR. Resistance exercise dosage in older adults: single- versus multiset effects on physical performance and body composition. *J. Am. Geriatr. Soc.* 53: 2090-2097, 2005.
76. Gentil, P, Fischer, B, Martorelli, AS, Lima, RM, and Bottaro, M. Effects of equal-volume resistance training performed one or two times a week in upper body muscle size and strength of untrained young men. *J. Sports Med. Phys. Fitness* 55: 144-149, 2015.
77. Gentil, P, Fisher, J, Steele, J, Campos, MH, Silva, MH, Paoli, A, Giessing, J, and Bottaro, M. Effects of equal-volume resistance training with different training frequencies in muscle size and strength in trained men. *PeerJ* 6: e5020, 2018.
78. Gibala, MJ, Interisano, SA, Tarnopolsky, MA, Roy, BD, MacDonald, JR, Yarasheski, KE, and MacDougall, JD. Myofibrillar disruption following acute concentric and eccentric resistance exercise in strength-trained men. *Can. J. Physiol. Pharmacol.* 78: 656-661, 2000.

79. Giessing, J, Fisher, J, Steele, J, Rothe, F, Raubold, K, and Eichmann, B. The effects of low volume resistance training with and without advanced techniques in trained participants. *J. Sports Med. Phys. Fitness* , 2014.
80. Gollnick, PD, Karlsson, J, Piehl, K, and Saltin, B. Selective glycogen depletion in skeletal muscle fibres of man following sustained contractions. *J. Physiol.* 241: 59-67, 1974.
81. Gomes, GK, Franco, CM, Nunes, PRP, and Orsatti, FL. High-frequency resistance training is not more effective than low-frequency resistance training in increasing muscle mass and strength in well-trained men. *J. Strength Cond Res.* , 2018.
82. Gonzalez, AM, Ghigiarelli, JJ, Sell, KM, Shone, EW, Kelly, CF, and Mangine, GT. Muscle activation during resistance exercise at 70% and 90% 1-repetition maximum in resistance-trained men. *Muscle Nerve* , 2016.
83. Goto, K, Ishii, N, Kizuka, T, and Takamatsu, K. The impact of metabolic stress on hormonal responses and muscular adaptations. *Med. Sci. Sports Exerc.* 37: 955-963, 2005.
84. Goto, M, Maeda, C, Hirayama, T, Terada, S, Nirengi, S, Kurosawa, Y, Nagano, A, and Hamaoka, T. Partial Range of Motion Exercise Is Effective for Facilitating Muscle Hypertrophy and Function Through Sustained Intramuscular Hypoxia in Young Trained Men. *J. Strength Cond Res.* 33: 1286-1294, 2019.
85. Grgic, J, and Schoenfeld, BJ. Are the Hypertrophic Adaptations to High and Low-Load Resistance Training Muscle Fiber Type Specific? *Front. Physiol.* 9: 402, 2018.
86. Gundermann, D. Mechanisms of Blood Flow Restriction Exercise in Skeletal Muscle Adaptations. , 2016.
87. Hackett, DA, Johnson, NA, and Chow, CM. Training practices and ergogenic aids used by male bodybuilders. *J. Strength Cond Res.* 27: 1609-1617, 2013.
88. Hackett, DA, Amirthalingam, T, Mitchell, L, Mavros, Y, Wilson, GC, and Halaki, M. Effects of a 12-Week Modified German Volume Training Program on Muscle Strength and Hypertrophy-A Pilot Study. *Sports (Basel)* 6: 10.3390/sports6010007, 2018.
89. Haddad, F, and Adams, GR. Selected contribution: acute cellular and molecular responses to resistance exercise. *J. Appl. Physiol.* (1985) 93: 394-403, 2002.
90. Häkkinen, K, and Kallinen, M. Distribution of strength training volume into one or two daily sessions and neuromuscular adaptations in female athletes. *Electromyogr. Clin. Neurophysiol.* 34: 117-124, 1994.
91. Häkkinen, K, Pakarinen, A, Kraemer, WJ, Häkkinen, A, Valkeinen, H, and Alen, M. Selective muscle hypertrophy, changes in EMG and force, and serum hormones during strength training in older women. *J. Appl. Physiol.* 91: 569-580, 2001.
92. Hammarstrom, D, Ofsteng, S, Koll, L, Hanestadhaugen, M, Hollan, I, Apro, W, Whist, JE, Blomstrand, E, Ronnestad, BR, and Ellefsen, S. Benefits of higher resistance-training volume are related to ribosome biogenesis. *J. Physiol.* , 2019.
93. Hanssen, KE, Kvamme, NH, Nilsen, TS, Ronnestad, B, Ambjornsen, IK, Norheim, F, Kadi, F, Hallen, J, Drevon, CA, and Raastad, T. The effect of strength training volume on satellite cells, myogenic regulatory factors, and growth factors. *Scand. J. Med. Sci. Sports* 23: 728-739, 2013.
94. Hartman, MJ, Clark, B, Bembens, DA, Kilgore, JL, and Bembens, MG. Comparisons between twice-daily and once-daily training sessions in male weight lifters. *Int. J. Sports Physiol. Perform.* 2: 159-169, 2007.
95. Haun, CT, Vann, CG, Mobley, CB, Roberson, PA, Osburn, SC, Holmes, HM, Mumford, PM, Romero, MA, Young, KC, Moon, JR, Gladden, LB, Arnold, RD, Israetel, MA, Kirby, AN, and Roberts, MD. Effects of Graded Whey Supplementation During Extreme-Volume Resistance Training. *Front. Nutr.* 5: 84, 2018.
96. Hawkins, SA, Schroeder, ET, Wiswell, RA, Jaque, SV, Marcell, TJ, and Costa, K. Eccentric muscle action increases site-specific osteogenic response. *Med. Sci. Sports Exerc.* 31: 1287-1292, 1999.
97. Heaselgrave, SR, Blacker, J, Smeuninx, B, McKendry, J, and Breen, L. Dose-Response Relationship of Weekly Resistance-Training Volume and Frequency on Muscular Adaptations in Trained Men. *Int. J. Sports Physiol. Perform.* 14: 360-368, 2019.
98. Hedayatpour, N, and Falla, D. Non-uniform muscle adaptations to eccentric exercise and the implications for training and sport. *J. Electromyogr. Kinesiol.* 22: 329-333, 2012.
99. Helms, E, Fitschen, PJ, Aragon, A, Cronin, J, and Schoenfeld, BJ. Recommendations for natural bodybuilding contest preparation: resistance and cardiovascular training. *J. Sports Med. Phys. Fitness*, 2014.
100. Herman-Montemayor, JR, Hikida, RS, and Staron, RS. Early-phase satellite cell and myonuclear domain adaptations to slow-speed versus traditional resistance training programs. *J. Strength Cond Res.* , 2015.
101. Heron, MI, and Richmond, FJ. In-series fiber architecture in long human muscles. *J. Morphol.* 216: 35-45, 1993.
102. Higbie, EJ, Cureton, KJ, Warren, GL, 3rd, and Prior, BM. Effects of concentric and eccentric training on muscle strength, cross-sectional area, and neural activation. *J. Appl. Physiol.* (1985) 81: 2173-2181, 1996.
103. Hill-Haas, S, Bishop, D, Dawson, B, Goodman, C, and Edge, J. Effects of rest interval during high-repetition resistance training on strength, aerobic fitness, and repeated-sprint ability. *J. Sports Sci.* 25: 619-628, 2007.
104. Hoeger, WW, Barette, SL, Hale, DF, and Hopkins, DR. Relationship between repetitions and selected percentages of one repetition maximum.. *J. Appl. Sport Sci. Res.* 1: 11-13, 1987.
105. Holm, L, Reitelseder, S, Pedersen, TG, Doessing, S, Petersen, SG, Flyvbjerg, A, Andersen, JL, Aagaard, P, and Kjaer, M. Changes in muscle size and MHC composition in response to resistance exercise with heavy and light loading intensity. *J. Appl. Physiol.* 105: 1454-1461, 2008.
106. Hortobagyi, T, Hill, JP, Houmard, JA, Fraser, DD, Lambert, NJ, and Israel, RG. Adaptive responses to muscle lengthening and shortening in humans. *J. Appl. Physiol.* (1985) 80: 765-772, 1996.
107. Hortobagyi, T, Dempsey, L, Fraser, D, Zheng, D, Hamilton, G, Lambert, J, and Dohm, L. Changes in muscle strength, muscle fibre size and myofibrillar gene expression after immobilization and retraining in humans. *J. Physiol.* 524 Pt 1: 293-304, 2000.
108. Housh, DJ, Housh, TJ, Johnson, GO, and Chu, WK. Hypertrophic response to unilateral concentric isokinetic resistance training. *J. Appl. Physiol.* (1985) 73: 65-70, 1992.
109. Hulmi, JJ, Walker, S, Ahtiainen, JP, Nyman, K, Kraemer, WJ, and Häkkinen, K. Molecular signaling in muscle is affected by the specificity of resistance exercise protocol. *Scand. J. Med. Sci. Sports* 22: 240-248, 2012.
110. Izquierdo, M, Ibanez, J, Gonzalez-Badillo, JJ, Häkkinen, K, Ratamess, NA, Kraemer, WJ, French, DN, Eslava, J, Altadill, A, Asiain, X, and Gorostiaga, EM. Differential effects of strength training leading to failure versus not to failure on hormonal responses, strength, and muscle power gains. *J. Appl. Physiol.* 100: 1647-1656, 2006.

111. Jakobsgaard, JE, Christiansen, M, Sieljacks, P, Wang, J, Groennebaek, T, de Paoli, E, and Vissing, K. Impact of blood flow-restricted bodyweight exercise on skeletal muscle adaptations. *Clin. Physiol. Funct. Imaging*, 2018.
112. Jenkins, ND, Housh, TJ, Bergstrom, HC, Cochrane, KC, Hill, EC, Smith, CM, Johnson, GO, Schmidt, RJ, and Cramer, JT. Muscle activation during three sets to failure at 80 vs. 30% 1RM resistance exercise. *Eur. J. Appl. Physiol.* 115: 2335-2347, 2015.
113. Jessee, MB, Buckner, SL, Mouser, JG, Mattocks, KT, Dankel, SJ, Abe, T, Bell, ZW, Bentley, JP, and Loenneke, JP. Muscle Adaptations to High-Load Training and Very Low-Load Training With and Without Blood Flow Restriction. *Front. Physiol.* 9: 1448, 2018.
114. Johnston, BD. Moving Too Rapidly in Strength Training Will Unload Muscles and Limit Full Range Strength Development Adaptation. *Journal of Exercise Physiology Online* 8: 36-45, 2005.
115. Jones, DA, and Rutherford, OM. Human muscle strength training: the effects of three different regimens and the nature of the resultant changes. *J. Physiol. (Lond.)* 391: 1-11, 1987.
116. Karsten, B, Fu, YL, Larumbe-Zabala, E, Seijo, M, and Naclerio, F. Impact of Two High-Volume Set Configuration Workouts on Resistance Training Outcomes in Recreationally Trained Men. *J. Strength Cond Res.*, 2019.
117. Keeler, LK, Finkelstein, LH, Miller, W, and Fernhall, B. Early-phase adaptations of traditional-speed vs. superslow resistance training on strength and aerobic capacity in sedentary individuals. *J. Strength Cond Res.* 15: 309-314, 2001.
118. Keogh, JWL, Wilson, GJ, and Weatherby, RP. A Cross-Sectional Comparison of Different Resistance Training Techniques in the Bench Press. *Journal of Strength and Conditioning Research* 13: 247-258, 1999.
119. Kerkick, CM, Wilborn, CD, Campbell, BI, Roberts, MD, Rasmussen, CJ, Greenwood, M, and Kreider, RB. Early-phase adaptations to a split-body, linear periodization resistance training program in college-aged and middle-aged men. *J. Strength Cond Res* 23: 962-971, 2009.
120. Kim, PL, Staron, RS, and Phillips, SM. Fasted-state skeletal muscle protein synthesis after resistance exercise is altered with training. *J. Physiol.* 568: 283-290, 2005.
121. Kim, SY, Ko, JB, Farthing, JP, and Butcher, SJ. Investigation of supraspinatus muscle architecture following concentric and eccentric training. *J. Sci. Med. Sport*, 2014.
122. Komi, PV, and Buskirk, ER. Effect of eccentric and concentric muscle conditioning on tension and electrical activity of human muscle. *Ergonomics* 15: 417-434, 1972.
123. Kraemer, WJ. A series of studies: the physiological basis for strength training in American football: fact over philosophy. *J. Strength Cond Res* 11: 131-134, 1997.
124. Kraemer, WJ, Noble, BJ, Clark, MJ, and Culver, BW. Physiologic responses to heavy-resistance exercise with very short rest periods. *Int. J. Sports Med.* 8: 247-252, 1987.
125. Kraemer, WJ, and Ratamess, NA. Fundamentals of resistance training: progression and exercise prescription. *Med. Sci. Sports Exerc.* 36: 674-688, 2004.
126. Krentz, JR, and Farthing, JP. Neural and morphological changes in response to a 20-day intense eccentric training protocol. *Eur. J. Appl. Physiol.* 110: 333-340, 2010.
127. Kubo, K, Ikebukuro, T, and Yata, H. Effects of squat training with different depths on lower limb muscle volumes. *Eur. J. Appl. Physiol.*, 2019.
128. Kumar, V, Selby, A, Rankin, D, Patel, R, Atherton, P, Hildebrandt, W, Williams, J, Smith, K, Seynnes, O, Hiscock, N, and Rennie, MJ. Age-related differences in the dose-response relationship of muscle protein synthesis to resistance exercise in young and old men. *J. Physiol.* 587: 211-217, 2009.
129. Lasevicius, T, Ugrinowitsch, C, Schoenfeld, BJ, Roschel, H, Tavares, LD, De Souza, EO, Laurentino, G, and Tricoli, V. Effects of different intensities of resistance training with equated volume load on muscle strength and hypertrophy. *Eur. J. Sport. Sci.* 18: 772-780, 2018.
130. Lasevicius, T, Schoenfeld, BJ, Grgic, J, Laurentino, G, Tavares, LD, and Tricoli, V. Similar Muscular Adaptations in Resistance Training Performed Two Versus Three Days Per Week. *J. Hum. Kinet* 68: 135-143, 2019.
131. Leger, B, Cartoni, R, Praz, M, Lamon, S, Deriaz, O, Crettenand, A, Gobelet, C, Rohmer, P, Konzelmann, M, Luthi, F, and Russell, AP. Akt signalling through GSK-3 β , mTOR and Foxo1 is involved in human skeletal muscle hypertrophy and atrophy. *J. Physiol.* 576: 923-933, 2006.
132. Lim, C, Kim, HJ, Morton, RW, Harris, R, Phillips, SM, Jeong, TS, and Kim, CK. Resistance Exercise-induced Changes in Muscle Phenotype Are Load Dependent. *Med. Sci. Sports Exerc.* 51: 2578-2585, 2019.
133. Lindman, R, Eriksson, A, and Thornell, LE. Fiber type composition of the human female trapezius muscle: enzyme-histochemical characteristics. *Am. J. Anat.* 190: 385-392, 1991.
134. Lysenko, EA, Popov, DV, Vepkhvadze, TF, Sharova, AP, and Vinogradova, OL. Signaling responses to high and moderate load strength exercise in trained muscle. *Physiol. Rep.* 7: e14100, 2019.
135. MacDougall, JD, Gibala, MJ, Tarnopolsky, MA, MacDonald, JR, Interisano, SA, and Yarasheski, KE. The time course for elevated muscle protein synthesis following heavy resistance exercise. *Can. J. Appl. Physiol.* 20: 480-486, 1995.
136. Machado, M, Koch, AJ, Willardson, JM, Pereira, LS, Cardoso, MI, Motta, MK, Pereira, R, and Monteiro, AN. Effect of varying rest intervals between sets of assistance exercises on creatine kinase and lactate dehydrogenase responses. *J. Strength Cond Res.* 25: 1339-1345, 2011.
137. Maeo, S, Shan, X, Otsuka, S, Kanehisa, H, and Kawakami, Y. Neuromuscular Adaptations to Work-matched Maximal Eccentric versus Concentric Training. *Med. Sci. Sports Exerc.* 50: 1629-1640, 2018.
138. Martineau, LC, and Gardiner, PF. Insight into skeletal muscle mechanotransduction: MAPK activation is quantitatively related to tension. *J. Appl. Physiol.* 91: 693-702, 2001.
139. Martineau, LC, and Gardiner, PF. Skeletal muscle is sensitive to the tension-time integral but not to the rate of change of tension, as assessed by mechanically induced signaling. *J. Biomech.* 35: 657-663, 2002.
140. Martorelli, S, Cadore, EL, Izquierdo, M, Celes, R, Martorelli, A, Cleto, VA, Alvarenga, JG, and Bottaro, M. Strength Training with Repetitions to Failure does not Provide Additional Strength and Muscle Hypertrophy Gains in Young Women. *Eur. J. Transl. Myol* 27: 6339, 2017.
141. Marzolini, S, Oh, PI, Thomas, SG, and Goodman, JM. Aerobic and resistance training in coronary disease: single versus multiple sets. *Med. Sci. Sports Exerc.* 40: 1557-1564, 2008.
142. Massey, CD, Vincent, J, Maneval, M, and Johnson, JT. Influence of range of motion in resistance training in women: early phase adaptations. *J. Strength Cond Res.* 19: 409-411, 2005.
143. Masuda, K, Choi, JY, Shimojo, H, and Katsuta, S. Maintenance of myoglobin concentration in human skeletal muscle after

- heavy resistance training. *Eur. J. Appl. Physiol. Occup. Physiol.* 79: 347-352, 1999.
144. Mayhew, TP, Rothstein, JM, Finucane, SD, and Lamb, RL. Muscular adaptation to concentric and eccentric exercise at equal power levels. *Med. Sci. Sports Exerc.* 27: 868-873, 1995.
 145. Mazzetti, S, Douglass, M, Yocum, A, and Harber, M. Effect of explosive versus slow contractions and exercise intensity on energy expenditure. *Med. Sci. Sports Exerc.* 39: 1291-1301, 2007.
 146. McBride, JM, Blaak, JB, and Triplett-McBride, T. Effect of resistance exercise volume and complexity on EMG, strength, and regional body composition. *Eur. J. Appl. Physiol.* 90: 626-632, 2003.
 147. McHugh, MP, Connolly, DA, Eston, RG, and Gleim, GW. Electromyographic analysis of exercise resulting in symptoms of muscle damage. *J. Sports Sci.* 18: 163-172, 2000.
 148. McHugh, MP. Recent advances in the understanding of the repeated bout effect: the protective effect against muscle damage from a single bout of eccentric exercise. *Scand. J. Med. Sci. Sports* 13: 88-97, 2003.
 149. McLester, JR, Bishop, P, and Williams, ME. Comparison of 1 day and 3 days per week of equal-volume resistance training in experienced subjects. *J Strength Cond Res* 14: 273-281, 2000.
 150. McMahon, G, Morse, CI, Burden, A, Winwood, K, and Onambele, GL. Muscular adaptations and insulin-like growth factor-I (IGF-I) responses to resistance training are stretch-mediated. *Muscle Nerve*, 2013.
 151. McMahon, GE, Morse, CI, Burden, A, Winwood, K, and Onambele, GL. Impact of range of motion during ecologically valid resistance training protocols on muscle size, subcutaneous fat, and strength. *J. Strength Cond Res.* 28: 245-255, 2014.
 152. Medeiros, HS, Jr, Mello, RS, Amorim, MZ, Koch, AJ, and Machado, M. Planned intensity reduction to maintain repetitions within recommended hypertrophy range. *Int. J. Sports Physiol. Perform.* 8: 384-390, 2013.
 153. Mitchell, CJ, Churchward-Venne, TA, West, DD, Burd, NA, Breen, L, Baker, SK, and Phillips, SM. Resistance exercise load does not determine training-mediated hypertrophic gains in young men. *J. Appl. Physiol.*, 2012.
 154. Miyamoto, N, Wakahara, T, Ema, R, and Kawakami, Y. Non-uniform muscle oxygenation despite uniform neuromuscular activity within the vastus lateralis during fatiguing heavy resistance exercise. *Clin. Physiol. Funct. Imaging*, 2013.
 155. Mookerjee, S, and Ratamess, N. Comparison of strength differences and joint action durations between full and partial range-of-motion bench press exercise. *J Strength Cond Res* 13: 76-81, 1999.
 156. Moore, DR, Young, M, and Phillips, SM. Similar increases in muscle size and strength in young men after training with maximal shortening or lengthening contractions when matched for total work. *Eur. J. Appl. Physiol.* 112: 1587-1592, 2012.
 157. Moore, DR, Phillips, SM, Babraj, JA, Smith, K, and Rennie, MJ. Myofibrillar and collagen protein synthesis in human skeletal muscle in young men after maximal shortening and lengthening contractions. *Am. J. Physiol. Endocrinol. Metab.* 288: 1153-1159, 2005.
 158. Morton, RW, Sonne, MW, Farias Zuniga, A, Mohammad, IYZ, Jones, A, McGlory, C, Keir, PJ, Potvin, JR, and Phillips, SM. Muscle fibre activation is unaffected by load and repetition duration when resistance exercise is performed to task failure. *J. Physiol.* 597: 4601-4613, 2019.
 159. Muddle, TWD, Colquhoun, RJ, Magrini, MA, Luera, MJ, DeFreitas, JM, and Jenkins, NDM. Effects of fatiguing, submaximal high- versus low-torque isometric exercise on motor unit recruitment and firing behavior. *Physiol. Rep.* 6: e13675, 2018.
 160. Munn, J, Herbert, RD, Hancock, MJ, and Gandevia, SC. Resistance training for strength: effect of number of sets and contraction speed. *Med. Sci. Sports Exerc.* 37: 1622-1626, 2005.
 161. Murlasits, Z, Reed, J, and Wells, K. Effect of resistance training frequency on physiological adaptations in older adults. *Journal of Exercise Science & Fitness* 10: 28-32, 2012.
 162. Nardone, A, and Schieppati, M. Shift of activity from slow to fast muscle during voluntary lengthening contractions of the triceps surae muscles in humans. *J. Physiol. (Lond.)* 395: 363-381, 1988.
 163. Narici, MV, Hoppeler, H, Kayser, B, Landoni, L, Claassen, H, Gavardi, C, Conti, M, and Cerretelli, P. Human quadriceps cross-sectional area, torque and neural activation during 6 months strength training. *Acta Physiol. Scand.* 157: 175-186, 1996.
 164. Nascimento, MAD, Gerage, AM, Silva, DRPD, Ribeiro, AS, Machado, DGDS, Pina, FLC, Tomeleri, CM, Venturini, D, Barbosa, DS, Mayhew, JL, and Cyrino, ES. Effect of resistance training with different frequencies and subsequent detraining on muscle mass and appendicular lean soft tissue, IGF-1, and testosterone in older women. *Eur. J. Sport. Sci.* : 1-9, 2018.
 165. Neils, CM, Udermann, BE, Brice, GA, Winchester, JB, and McGuigan, MR. Influence of contraction velocity in untrained individuals over the initial early phase of resistance training. *J. Strength Cond Res.* 19: 883-887, 2005.
 166. Nicholson, G, McLoughlin, G, Bissas, A, and Ispoglou, T. Do the acute biochemical and neuromuscular responses justify the classification of strength and hypertrophy-type resistance exercise? *J. Strength Cond Res.*, 2014.
 167. Nickols-Richardson, SM, Miller, LE, Wooten, DF, Ramp, WK, and Herbert, WG. Concentric and eccentric isokinetic resistance training similarly increases muscular strength, fat-free soft tissue mass, and specific bone mineral measurements in young women. *Osteoporos. Int.* 18: 789-796, 2007.
 168. Nobrega, SR, Ugrinowitsch, C, Pintanel, L, Barcelos, C, and Libardi, CA. Effect of Resistance Training to Muscle Failure vs. Volitional Interruption at High- and Low-Intensities on Muscle Mass and Strength. *J. Strength Cond Res.* 32: 162-169, 2018.
 169. Nogueira, W, Gentil, P, Mello, SN, Oliveira, RJ, Bezerra, AJ, and Bottaro, M. Effects of power training on muscle thickness of older men. *Int. J. Sports Med.* 30: 200-204, 2009.
 170. Noorkoiv, M, Nosaka, K, and Blazevich, AJ. Neuromuscular adaptations associated with knee joint angle-specific force change. *Med. Sci. Sports Exerc.* 46: 1525-1537, 2014.
 171. Ochi, E, Maruo, M, Tsuchiya, Y, Ishii, N, Miura, K, and Sasaki, K. Higher Training Frequency Is Important for Gaining Muscular Strength Under Volume-Matched Training. *Front. Physiol.* 9: 744, 2018.
 172. Ogasawara, R, Loenneke, JP, Thiebaud, RS, and Abe, T. Low-Load Bench Press Training to Fatigue Results in Muscle Hypertrophy Similar to High-Load Bench Press Training. *Int J Clin Med.* 4: 114-121, 2013.
 173. Ogborn, D, and Schoenfeld, BJ. The role of fiber types in muscle hypertrophy: Implications for loading strategies. *Strength Cond J* 36: 20-25, 2014.
 174. Ostrowski, K, Wilson, GJ, Weatherby, R, Murphy, PW, and Little, AD. The effect of weight training volume on hormonal output and muscular size and function. *J Strength Cond Res* 11: 149-154, 1997.

175. Pareja-Blanco, F, Rodriguez-Rosell, D, Sanchez-Medina, L, Sanchis-Moysi, J, Dorado, C, Mora-Custodio, R, Yanez-Garcia, JM, Morales-Alamo, D, Perez-Suarez, I, Calbet, JAL, and Gonzalez-Badillo, JJ. Effects of velocity loss during resistance training on athletic performance, strength gains and muscle adaptations. *Scand. J. Med. Sci. Sports* 27: 724-735, 2017.
176. Peterson, MD, Rhea, MR, and Alvar, BA. Applications of the dose-response for muscular strength development: a review of meta-analytic efficacy and reliability for designing training prescription. *J. Strength Cond Res.* 19: 950-958, 2005.
177. Phillips, SM, Tipton, KD, Aarsland, A, Wolf, SE, and Wolfe, RR. Mixed muscle protein synthesis and breakdown after resistance exercise in humans. *Am. J. Physiol.* 273: E99-107, 1997.
178. Piirainen, JM, Tanskanen, M, Nissila, J, Kaarela, J, Vaarala, A, Sippola, N, and Linnamo, V. Effects of a heart rate-based recovery period on hormonal, neuromuscular, and aerobic performance responses during 7 weeks of strength training in men. *J. Strength Cond Res.* 25: 2265-2273, 2011.
179. Pina, FLC, Nunes, JP, Nascimento, MA, Ribeiro, AS, Mayhew, JL, and Cyrino, ES. Similar Effects of 24 Weeks of Resistance Training Performed with Different Frequencies on Muscle Strength, Muscle Mass, and Muscle Quality in Older Women. *Int. J. Exerc. Sci.* 12: 623-635, 2019.
180. Pinto, RS, Gomes, N, Radaelli, R, Botton, CE, Brown, LE, and Bottaro, M. Effect of range of motion on muscle strength and thickness. *J. Strength Cond Res.* 26: 2140-2145, 2012.
181. Popov, DV, Tsvirkun, DV, Netreba, AI, Tarasova, OS, Prostova, AB, Larina, IM, Borovik, AS, and Vinogradova, OL. Hormonal adaptation determines the increase in muscle mass and strength during low-intensity strength training without relaxation. *Fiziol. Cheloveka* 32: 121-127, 2006.
182. Popov, DV, Lysenko, EA, Bachinin, AV, Miller, TF, Kurochkina, NS, Kravchenko, IV, Furalyov, VA, and Vinogradova, OL. Influence of resistance exercise intensity and metabolic stress on anabolic signaling and expression of myogenic genes in skeletal muscle. *Muscle Nerve* 51: 434-442, 2015.
183. Raastad, T, Kirketeig, A, Wolf, D, and Paulsen, G. Powerlifters improved strength and muscular adaptations to a greater extent when equal total training volume was divided into 6 compared to 3 training sessions per week. *17th annual conference of the ECSS*, 2012.
184. Radaelli, R, Botton, CE, Wilhelm, EN, Bottaro, M, Lacerda, F, Gaya, A, Moraes, K, Peruzzolo, A, Brown, LE, and Pinto, RS. Low- and high-volume strength training induces similar neuromuscular improvements in muscle quality in elderly women. *Exp. Gerontol.* 48: 710-716, 2013.
185. Radaelli, R, Botton, CE, Wilhelm, EN, Bottaro, M, Brown, LE, Lacerda, F, Gaya, A, Moraes, K, Peruzzolo, A, and Pinto, RS. Time course of low- and high-volume strength training on neuromuscular adaptations and muscle quality in older women. *Age (Dordr)* 36: 881-892, 2014.
186. Radaelli, R, Fleck, SJ, Leite, T, Leite, RD, Pinto, RS, Fernandes, L, and Simao, R. Dose Response of 1, 3 and 5 Sets of Resistance Exercise on Strength, Local Muscular Endurance and Hypertrophy. *J. Strength Cond Res.* 29: 1349-1358, 2014.
187. Radaelli, R, Wilhelm, EN, Botton, CE, Rech, A, Bottaro, M, Brown, LE, and Pinto, RS. Effects of single vs. multiple-set short-term strength training in elderly women. *Age (Dordr)* 36: 9720-014-9720-6. Epub 2014 Oct 31, 2014.
188. Rana, SR, Chleboun, GS, Gilders, RM, Hagerman, FC, Herman, JR, Hikida, RS, Kushnick, MR, Staron, RS, and Toma, K. Comparison of early phase adaptations for traditional strength and endurance, and low velocity resistance training programs in college-aged women. *J. Strength Cond Res.* 22: 119-127, 2008.
189. Ratamess, NA, Falvo, MJ, Mangine, GT, Hoffman, JR, Faigenbaum, AD, and Kang, J. The effect of rest interval length on metabolic responses to the bench press exercise. *Eur. J. Appl. Physiol.* 100: 1-17, 2007.
190. Reeves, ND, Maganaris, CN, Longo, S, and Narici, MV. Differential adaptations to eccentric versus conventional resistance training in older humans. *Exp. Physiol.* 94: 825-833, 2009.
191. Rhea, MR, Alvar, BA, Ball, SD, and Burkett, LN. Three sets of weight training superior to 1 set with equal intensity for eliciting strength. *J. Strength Cond Res.* 16: 525-529, 2002.
192. Ribeiro, AS, Souza, ME, Pina, FLC, Nascimento, MA, dos Santos, L, Antunes, M, Schoenfeld, BJ, and Cyrino, ES. Resistance training in older women: comparison of single vs. multiple sets on muscle strength and body composition. *Isokinetics and Exercise Science* 23: 53-60, 2015.
193. Ribeiro, AS, Schoenfeld, BJ, Silva, DR, Pina, FL, Guariglia, DA, Porto, M, Maesta, N, Burini, RC, and Cyrino, ES. Effect of Two- Versus Three-Way Split Resistance Training Routines on Body Composition and Muscular Strength in Bodybuilders: A Pilot Study. *Int. J. Sport Nutr. Exerc. Metab.*, 2015.
194. Richardson, DL, Duncan, MJ, Jimenez, A, Juris, PM, and Clarke, ND. Effects of movement velocity and training frequency of resistance exercise on functional performance in older adults: a randomised controlled trial. *Eur. J. Sport. Sci.* : 1-13, 2018.
195. Robbins, DW, Goodale, TL, Docherty, D, Behm, DG, and Tran, QT. The effects of load and training pattern on acute neuromuscular responses in the upper body. *J. Strength Cond Res.* 24: 2996-3007, 2010.
196. Ronnestad, BR, Egeland, W, Kvamme, NH, Refsnes, PE, Kadi, F, and Raastad, T. Dissimilar effects of one- and three-set strength training on strength and muscle mass gains in upper and lower body in untrained subjects. *J. Strength Cond Res.* 21: 157-163, 2007.
197. Ronnestad, BR, Nygaard, H, and Raastad, T. Physiological elevation of endogenous hormones results in superior strength training adaptation. *Eur. J. Appl. Physiol.* 111: 2249-2259, 2011.
198. Roschel, H, Ugrinowistch, C, Barroso, R, Batista, MA, Souza, EO, Aoki, MS, Siqueira-Filho, MA, Zanuto, R, Carvalho, CR, Neves, M, Mello, MT, and Tricoli, V. Effect of eccentric exercise velocity on akt/mTOR/p70(s6k) signaling in human skeletal muscle. *Appl. Physiol. Nutr. Metab.* 36: 283-290, 2011.
199. Ruas, CV, Brown, LE, Lima, CD, Gregory Haff, G, and Pinto, RS. Different Muscle Action Training Protocols on Quadriceps-Hamstrings Neuromuscular Adaptations. *Int. J. Sports Med.* 39: 355-365, 2018.
200. Sampson, JA, and Groeller, H. Is repetition failure critical for the development of muscle hypertrophy and strength? *Scand. J. Med. Sci. Sports*, 2015.
201. Saric, J, Lisica, D, Orlic, I, Grgic, J, Krieger, JW, Vuk, S, and Schoenfeld, BJ. Resistance training frequencies of 3 and 6 times per week produce similar muscular adaptations in resistance-trained men. *Journal of Strength and Conditioning Research* doi: 10.1519/JSC.0000000000002909. [Epub ahead of print], 2018.
202. Schoenfeld, B. The use of specialized training techniques to maximize muscle hypertrophy. *Strength Cond J* 33: 60-65, 2011.
203. Schoenfeld, BJ, Contreras, B, Vigotsky, A, Sonmez, GT, and Fontana, F. Upper body muscle activation during low- versus high-load resistance exercise in the bench press. *Isokinetics and Exercise Science* 24: 217-224, 2016.

204. Schoenfeld, BJ, Pope, ZK, Benik, FM, Hester, GM, Sellers, J, Nooner, JL, Schnaiter, JA, Bond-Williams, KE, Carter, AS, Ross, CL, Just, BL, Henselmans, M, and Krieger, JW. Longer inter-set rest periods enhance muscle strength and hypertrophy in resistance-trained men. *Journal of Strength and Conditioning Research* 30: 1805-1812, 2016.
205. Schoenfeld, BJ. The mechanisms of muscle hypertrophy and their application to resistance training. *J. Strength Cond Res.* 24: 2857-2872, 2010.
206. Schoenfeld, BJ. Does exercise-induced muscle damage play a role in skeletal muscle hypertrophy? *J. Strength Cond Res.* 26: 1441-1453, 2012.
207. Schoenfeld, BJ, Contreras, B, Willardson, JM, Fontana, F, and Tiriyaki-Sonmez, G. Muscle activation during low- versus high-load resistance training in well-trained men. *Eur. J. Appl. Physiol.* 114: 2491-2497, 2014.
208. Schoenfeld, BJ, Ratamess, NA, Peterson, MD, Contreras, B, Tiriyaki-Sonmez, G, and Alvar, BA. Effects of different volume-equated resistance training loading strategies on muscular adaptations in well-trained men. *J. Strength Cond Res.* 28: 2909-2918, 2014.
209. Schoenfeld, BJ, Ogborn, DI, and Krieger, JW. Effect of Repetition Duration During Resistance Training on Muscle Hypertrophy: A Systematic Review and Meta-Analysis. *Sports Med.*, 2015.
210. Schoenfeld, BJ, Peterson, MD, Ogborn, D, Contreras, B, and Sonmez, GT. Effects of Low- Versus High-Load Resistance Training on Muscle Strength and Hypertrophy in Well-Trained Men. *J. Strength Cond Res.* 29: 2954-2963, 2015.
211. Schoenfeld, BJ, Pope, ZK, Benik, FM, Hester, GM, Sellers, J, Nooner, JL, Schnaiter, JA, Bond-Williams, KE, Carter, AS, Ross, CL, Just, BL, Henselmans, M, and Krieger, JW. Longer inter-set rest periods enhance muscle strength and hypertrophy in resistance-trained men. *J. Strength Cond Res.*, 2015.
212. Schoenfeld, BJ, Ratamess, NA, Peterson, MD, Contreras, B, and Tiriyaki-Sonmez, G. Influence of Resistance Training Frequency on Muscular Adaptations in Well-Trained Men. *J. Strength Cond Res.* 29: 1821-1829, 2015.
213. Schoenfeld, BJ, Contreras, B, Vigotsky, AD, and Peterson, M. Differential Effects of Heavy Versus Moderate Loads on Measures of Strength and Hypertrophy in Resistance-Trained Men. *J. Sports Sci. Med.* 15: 715-722, 2016.
214. Schoenfeld, BJ, Grgic, J, Ogborn, D, and Krieger, JW. Strength and Hypertrophy Adaptations Between Low- vs. High-Load Resistance Training: A Systematic Review and Meta-analysis. *J. Strength Cond Res.* 31: 3508-3523, 2017.
215. Schoenfeld, BJ, Ogborn, D, and Krieger, JW. Dose-response relationship between weekly resistance training volume and increases in muscle mass: A systematic review and meta-analysis. *J. Sports Sci.* 35: 1073-1082, 2017.
216. Schoenfeld, BJ, Ogborn, DI, Vigotsky, AD, Franchi, MV, and Krieger, JW. Hypertrophic Effects of Concentric vs. Eccentric Muscle Actions: A Systematic Review and Meta-analysis. *J. Strength Cond Res.* 31: 2599-2608, 2017.
217. Schoenfeld, BJ, Vigotsky, A, Contreras, B, Golden, S, Alto, A, Larson, R, Winkelman, N, and Paoli, A. Differential effects of attentional focus strategies during long-term resistance training. *Eur. J. Sport. Sci.* 18: 705-712, 2018.
218. Schoenfeld, BJ, Contreras, B, Krieger, J, Grgic, J, Delcastillo, K, Belliard, R, and Alto, A. Resistance Training Volume Enhances Muscle Hypertrophy but Not Strength in Trained Men. *Med. Sci. Sports Exerc.* 51: 94-103, 2019.
219. Schoenfeld, BJ, Grgic, J, and Krieger, J. How many times per week should a muscle be trained to maximize muscle hypertrophy? A systematic review and meta-analysis of studies examining the effects of resistance training frequency. *J. Sports Sci.* 37: 1286-1295, 2019.
220. Schott, J, McCully, K, and Rutherford, OM. The role of metabolites in strength training. II. Short versus long isometric contractions. *Eur. J. Appl. Physiol. Occup. Physiol.* 71: 337-341, 1995.
221. Schuenke, MD, Herman, JR, Gliders, RM, Hagerman, FC, Hikida, RS, Rana, SR, Ragg, KE, and Staron, RS. Early-phase muscular adaptations in response to slow-speed versus traditional resistance-training regimens. *Eur. J. Appl. Physiol.* 112: 3585-3595, 2012.
222. Scott, BR, Slattery, KM, and Dascombe, BJ. Intermittent hypoxic resistance training: Is metabolic stress the key moderator? *Med. Hypotheses* 84: 145-149, 2015.
223. Segal, RL. Neuromuscular compartments in the human biceps brachii muscle. *Neurosci. Lett.* 140: 98-102, 1992.
224. Seger, JY, Arvidsson, B, and Thorstensson, A. Specific effects of eccentric and concentric training on muscle strength and morphology in humans. *Eur. J. Appl. Physiol. Occup. Physiol.* 79: 49-57, 1998.
225. Senna, GW, Figueiredo, T, Scudese, E, Baffi, M, Carneiro, F, Moraes, E, Miranda, H, and Simão, R. Influence of different rest interval lengths in multi-joint and single-joint exercises on repetition performance, perceived exertion, and blood lactate. *Journal of Exercise Physiology* 15: 96-106, 2012.
226. Serra, R, Saavedra, E, Jotta, B, Novaes, J, Cardozo, D, Alves, H, Dias, M, and Simao, R. The Influence Weekly Resistance Training Frequency on Strength and Body Composition. *International Journal of Sports Science* 8: 19-24, 2018.
227. Seynnes, OR, de Boer, M, and Narici, MV. Early skeletal muscle hypertrophy and architectural changes in response to high-intensity resistance training. *J. Appl. Physiol.* 102: 368-373, 2007.
228. Shephstone, TN, Tang, JE, Dallaire, S, Schuenke, MD, Staron, RS, and Phillips, SM. Short-term high- vs. low-velocity isokinetic lengthening training results in greater hypertrophy of the elbow flexors in young men. *J. Appl. Physiol.* (1985) 98: 1768-1776, 2005.
229. Shibata, K, Takizawa, K, Nosaka, K, and Mizuno, M. Effects of Prolonging Eccentric Phase Duration in Parallel Back-Squat Training to Momentary Failure on Muscle Cross-Sectional Area, Squat One Repetition Maximum, and Performance Tests in University Soccer Players. *J. Strength Cond Res.*, 2018.
230. Signorile, JF, Lew, KM, Stoutenberg, M, Pluchino, A, Lewis, JE, and Gao, J. Range of motion and leg rotation affect electromyography activation levels of the superficial quadriceps muscles during leg extension. *J. Strength Cond Res.* 28: 2536-2545, 2014.
231. Simao, R, Spinetti, J, de Salles, BE, Oliveira, LE, Matta, T, Miranda, F, Miranda, H, and Costa, PB. Influence of exercise order on maximum strength and muscle thickness in untrained men. *J. Sports Sci. Med.* 9: 1-7, 2010.
232. Simao, R, de Salles, BE, Figueiredo, T, Dias, I, and Willardson, JM. Exercise order in resistance training. *Sports Med.* 42: 251-265, 2012.
233. Smith, RC, and Rutherford, OM. The role of metabolites in strength training. I. A comparison of eccentric and concentric contractions. *Eur. J. Appl. Physiol. Occup. Physiol.* 71: 332-336, 1995.
234. Snyder, BJ, and Fry, WR. Effect of verbal instruction on muscle activity during the bench press exercise. *J. Strength Cond Res.* 26: 2394-2400, 2012.

235. Sooneste, H, Tanimoto, M, Kakigi, R, Saga, N, and Katamoto, S. Effects of training volume on strength and hypertrophy in young men. *J. Strength Cond Res.* 27: 8-13, 2013.
236. Souza-Junior, TP, Willardson, JM, Bloomer, R, Leite, RD, Fleck, SJ, Oliveira, PR, and Simao, R. Strength and hypertrophy responses to constant and decreasing rest intervals in trained men using creatine supplementation. *J. Int. Soc. Sports Nutr.* 8: 17-2783-8-17, 2011.
237. Spiering, BA, Kraemer, WJ, Anderson, JM, Armstrong, LE, Nindl, BC, Volek, JS, and Maresh, CM. Resistance exercise biology: manipulation of resistance exercise programme variables determines the responses of cellular and molecular signalling pathways. *Sports Med.* 38: 527-540, 2008.
238. Spinetti, J, de Salles, BF, Rhea, MR, Lavigne, D, Matta, T, Miranda, F, Fernandes, L, and Simao, R. Influence of exercise order on maximum strength and muscle volume in nonlinear periodized resistance training. *J. Strength Cond Res.* 24: 2962-2969, 2010.
239. Starkey, DB, Pollock, ML, Ishida, Y, Welsch, MA, Brechue, WF, Graves, JE, and Feigenbaum, MS. Effect of resistance training volume on strength and muscle thickness. *Med. Sci. Sports Exerc.* 28: 1311-1320, 1996.
240. Stefanaki, DGA, Dzulkarnain, A, and Gray, SR. Comparing the effects of low and high load resistance exercise to failure on adaptive responses to resistance exercise in young women. *J. Sports Sci.* 37: 1375-1380, 2019.
241. Taaffe, DR, Duret, C, Wheeler, S, and Marcus, R. Once-weekly resistance exercise improves muscle strength and neuromuscular performance in older adults. *J. Am. Geriatr. Soc.* 47: 1208-1214, 1999.
242. Takarada, Y, Takazawa, H, Sato, Y, Takebayashi, S, Tanaka, Y, and Ishii, N. Effects of resistance exercise combined with moderate vascular occlusion on muscular function in humans. *J. Appl. Physiol.* 88: 2097-2106, 2000.
243. Tang, JE, Perco, JG, Moore, DR, Wilkinson, SB, and Phillips, SM. Resistance training alters the response of fed state mixed muscle protein synthesis in young men. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 294: R172-8, 2008.
244. Tanimoto, M, and Ishii, N. Effects of low-intensity resistance exercise with slow movement and tonic force generation on muscular function in young men. *J. Appl. Physiol.* 100: 1150-1157, 2006.
245. Tanimoto, M, Sanada, K, Yamamoto, K, Kawano, H, Gando, Y, Tabata, I, Ishii, N, and Miyachi, M. Effects of whole-body low-intensity resistance training with slow movement and tonic force generation on muscular size and strength in young men. *J. Strength Cond Res.* 22: 1926-1938, 2008.
246. Tavares, LD, de Souza, EO, Ugrinowitsch, C, Laurentino, GC, Roschel, H, Aihara, AY, Cardoso, FN, and Tricoli, V. Effects of different strength training frequencies during reduced training period on strength and muscle cross-sectional area. *Eur. J. Sport. Sci.* 17: 665-672, 2017.
247. Taylor, LW, Wilborn, CD, Kreider, RB, and Willoughby, DS. Effects of resistance exercise intensity on extracellular signal-regulated kinase 1/2 mitogen-activated protein kinase activation in men. *J. Strength Cond Res.* 26: 599-607, 2012.
248. Tee, JC, Bosch, AN, and Lambert, MI. Metabolic consequences of exercise-induced muscle damage. *Sports Med.* 37: 827-836, 2007.
249. Teodoro, JL, da Silva, LXN, Fritsch, CG, Baroni, BM, Graziloli, R, Boeno, FP, Lopez, P, Gentil, P, Bottaro, M, Pinto, RS, Izquierdo, M, and Cadore, EL. Concurrent training performed with and without repetitions to failure in older men: A randomized clinical trial. *Scand. J. Med. Sci. Sports* 29: 1141-1152, 2019.
250. ter Haar Romeny, BM, Denier van der Gon, JJ, and Gielen, CC. Changes in recruitment order of motor units in the human biceps muscle. *Exp. Neurol.* 78: 360-368, 1982.
251. ter Haar Romeny, BM, van der Gon, JJ, and Gielen, CC. Relation between location of a motor unit in the human biceps brachii and its critical firing levels for different tasks. *Exp. Neurol.* 85: 631-650, 1984.
252. Terzis, G, Spengos, K, Mascher, H, Georgiadis, G, Manta, P, and Blomstrand, E. The degree of p70 S6k and S6 phosphorylation in human skeletal muscle in response to resistance exercise depends on the training volume. *Eur. J. Appl. Physiol.* 110: 835-843, 2010.
253. Tesch, PA, Ploutz-Snyder, LL, Ystrom, L, Castro, MJ, and Dudley, GA. Skeletal muscle glycogen loss evoked by resistance exercise. *Journal of Strength and Conditioning Research* : 67-73, 1998.
254. Thomas, MH, and Burns, SP. Increasing Lean Mass and Strength: A Comparison of High Frequency Strength Training to Lower Frequency Strength Training. *Int. J. Exerc. Sci.* 9: 159-167, 2016.
255. Tomeleri, CM, Ribeiro, AS, Nunes, JP, Schoenfeld, BJ, Souza, ME, Schiavoni, D, Junior, PS, Cavaglieri, CR, Cunha, PM, Venturini, D, and Barbosa, DS. Influence of Resistance Training Exercise Order on Muscle Strength, Hypertrophy, and Anabolic Hormones in Older Women: A Randomized Controlled Trial. *Journal of Strength and Conditioning Research* doi: 10.1519/JSC.0000000000003147, 2019.
256. Turpela, M, Häkkinen, K, Haff, GG, and Walker, S. Effects of different strength training frequencies on maximum strength, body composition and functional capacity in healthy older individuals. *Exp. Gerontol.* 98: 13-21, 2017.
257. Valamatos, MJ, Tavares, E, Santos, RM, Veloso, AP, and Mil-Homens, P. Influence of full range of motion vs. equalized partial range of motion training on muscle architecture and mechanical properties. *Eur. J. Appl. Physiol.* 118: 1969-1983, 2018.
258. Van Roie, E, Delecluse, C, Coudyzer, W, Boonen, S, and Bautmans, I. Strength training at high versus low external resistance in older adults: effects on muscle volume, muscle strength, and force-velocity characteristics. *Exp. Gerontol.* 48: 1351-1361, 2013.
259. Vikne, H, Refsnes, PE, Ekmark, M, Medbo, JJ, Gundersen, V, and Gundersen, K. Muscular performance after concentric and eccentric exercise in trained men. *Med. Sci. Sports Exerc.* 38: 1770-1781, 2006.
260. Villanueva, MG, Lane, CJ, and Schroeder, ET. Short rest interval lengths between sets optimally enhance body composition and performance with 8 weeks of strength resistance training in older men. *Eur. J. Appl. Physiol.* 115: 295-308, 2015.
261. Vissing, K, Rahbek, SK, Lamon, S, Farup, J, Stefanetti, RJ, Wallace, MA, Vendelbo, MH, and Russell, A. Effect of resistance exercise contraction mode and protein supplementation on members of the STARS signalling pathway. *J. Physiol.* 591: 3749-3763, 2013.
262. Wakahara, T, Miyamoto, N, Sugisaki, N, Murata, K, Kanehisa, H, Kawakami, Y, Fukunaga, T, and Yanai, T. Association between regional differences in muscle activation in one session of resistance exercise and in muscle hypertrophy after resistance training. *Eur. J. Appl. Physiol.* 112: 1569-1576, 2012.
263. Wakahara, T, Fukutani, A, Kawakami, Y, and Yanai, T. Nonuniform muscle hypertrophy: its relation to muscle activation in training session. *Med. Sci. Sports Exerc.* 45: 2158-2165, 2013.

264. Watanabe, Y, Tanimoto, M, Ohgane, A, Sanada, K, Miyachi, M, and Ishii, N. Increased muscle size and strength from slow-movement, low-intensity resistance exercise and tonic force generation. *J. Aging Phys. Act.* 21: 71-84, 2013.
265. Watanabe, Y, Madarame, H, Ogasawara, R, Nakazato, K, and Ishii, N. Effect of very low-intensity resistance training with slow movement on muscle size and strength in healthy older adults. *Clin. Physiol. Funct. Imaging* 34: 463-470, 2014.
266. Weiss, LW, Coney, HD, and Clark, FC. Gross measures of exercise-induced muscular hypertrophy. *J. Orthop. Sports Phys. Ther.* 30: 143-148, 2000.
267. Wernbom, M, Augustsson, J, and Thomee, R. The influence of frequency, intensity, volume and mode of strength training on whole muscle cross-sectional area in humans. *Sports Med.* 37: 225-264, 2007.
268. West, DW, Burd, NA, Tang, JE, Moore, DR, Staples, AW, Holwerda, AM, Baker, SK, and Phillips, SM. Elevations in ostensibly anabolic hormones with resistance exercise enhance neither training-induced muscle hypertrophy nor strength of the elbow flexors. *J. Appl. Physiol.* 108: 60-67, 2010.
269. West, DW, Cotie, LM, Mitchell, CJ, Churchward-Venne, TA, MacDonald, MJ, and Phillips, SM. Resistance exercise order does not determine postexercise delivery of testosterone, growth hormone, and IGF-1 to skeletal muscle. *Appl. Physiol. Nutr. Metab.* 38: 220-226, 2013.
270. Westcott, WL, Winett, RA, Anderson, ES, Wojcik, JR, Loud, RL, Cleggett, E, and Glover, S. Effects of regular and slow speed resistance training on muscle strength. *J. Sports Med. Phys. Fitness* 41: 154-158, 2001.
271. Wickiewicz, TL, Roy, RR, Powell, PL, and Edgerton, VR. Muscle architecture of the human lower limb. *Clin. Orthop. Relat. Res.* (179): 275-283, 1983.
272. Willardson, JM, Norton, L, and Wilson, G. Training to failure and beyond in mainstream resistance exercise programs. *Strength Cond J* 32: 21-29, 2010.
273. Woodley, SJ, and Mercer, SR. Hamstring muscles: architecture and innervation. *Cells Tissues Organs* 179: 125-141, 2005.
274. Young, WB, and Bilby, GE. The effect of voluntary effort to influence speed of contraction on strength, muscular power, and hypertrophy development. *Journal of Strength and Conditioning Research* 7: 172-178, 1993.
275. Yue, FL, Karsten, B, Larumbe-Zabala, E, Seijo, M, and Naclerio, F. Comparison of 2 weekly-equalized volume resistance-training routines using different frequencies on body composition and performance in trained males. *Appl. Physiol. Nutr. Metab.* 43: 475-481, 2018.
276. Zaroni, RS, Brigatto, FA, Schoenfeld, BJ, Braz, TV, Benvenutti, JC, Germano, MD, Marchetti, PH, Aoki, MS, and Lopes, CR. High Resistance-Training Frequency Enhances Muscle Thickness in Resistance-Trained Men. *Journal of Strength and Conditioning Research* Published Ahead-of-Print, 2018.
277. Zourdos, MC, Klemp, A, Dolan, C, Quiles, JM, Schau, KA, Jo, E, Helms, E, Esgro, B, Duncan, S, Garcia Merino, S, and Blanco, R. Novel Resistance Training-Specific Rating of Perceived Exertion Scale Measuring Repetitions in Reserve. *J. Strength Cond Res.* 30: 267-275, 2016.
- in stretch-induced enlargement. *J. Appl. Physiol.* (1985) 66: 771-781, 1989.
3. Andersen, LL, Magnusson, SP, Nielsen, M, Haleem, J, Poulsen, K, and Aagaard, P. Neuromuscular activation in conventional therapeutic exercises and heavy resistance exercises: implications for rehabilitation. *Phys. Ther.* 86: 683-697, 2006.
4. Angleri, V, Ugrinowitsch, C, and Libardi, CA. Crescent pyramid and drop-set systems do not promote greater strength gains, muscle hypertrophy, and changes on muscle architecture compared with traditional resistance training in well-trained men. *Eur. J. Appl. Physiol.* 117: 359-369, 2017.
5. Antonio, J, and Gonyea, WJ. Progressive stretch overload of skeletal muscle results in hypertrophy before hyperplasia. *J. Appl. Physiol.* (1985) 75: 1263-1271, 1993.
6. Assis-Pereira, PE, Motoyama, YL, Esteves, GJ, Quinelato, WC, Botter, L, Tanaka, KH, and Azevedo, P. Resistance training with slow speed of movement is better for hypertrophy and muscle strength gains than fast speed of movement. *Int J Appl Exerc Physiol* 5: 37-43, 2016.
7. Augustsson, J, Thomee, R, Hornstedt, P, Lindblom, J, Karlsson, J, and Grimby, G. Effect of pre-exhaustion exercise on lower-extremity muscle activation during a leg press exercise. *J. Strength Cond Res.* 17: 411-416, 2003.
8. Bleakley, C, McDonough, S, Gardner, E, Baxter, GD, Hopkins, JT, and Davison, GW. Cold-water immersion (cryotherapy) for preventing and treating muscle soreness after exercise. *Cochrane Database Syst. Rev.* (2):CD008262. doi: CD008262, 2012.
9. Bleakley, CM, and Davison, GW. What is the biochemical and physiological rationale for using cold-water immersion in sports recovery? A systematic review. *Br. J. Sports Med.* 44: 179-187, 2010.
10. Brennecke, A, Guimaraes, TM, Leone, R, Cadarci, M, Mochizuki, L, Simao, R, Amadio, AC, and Serrao, JC. Neuromuscular activity during bench press exercise performed with and without the preexhaustion method. *J. Strength Cond Res.* 23: 1933-1940, 2009.
11. Brentano, MA, Umpierre, D, Santos, LP, Lopes, AL, Radaelli, R, Pinto, RS, and Kruel, LFM. Muscle Damage and Muscle Activity Induced by Strength Training Super-Sets in Physically Active Men. *J. Strength Cond Res.* 31: 1847-1858, 2017.
12. Broatch, JR, Petersen, A, and Bishop, DJ. The Influence of Post-Exercise Cold-Water Immersion on Adaptive Responses to Exercise: A Review of the Literature. *Sports Med.* 48: 1369-1387, 2018.
13. Cheng, AJ. Cooling down the use of cryotherapy for post-exercise skeletal muscle recovery. *Temperature (Austin)* 5: 103-105, 2018.
14. Drinkwater, EJ, Latella, C, Wilshire, C, Bird, SP, and Skein, M. Foam Rolling as a Recovery Tool Following Eccentric Exercise: Potential Mechanisms Underpinning Changes in Jump Performance. *Front. Physiol.* 10: 768, 2019.
15. Dupuy, O, Douzi, W, Theurot, D, Bosquet, L, and Dugue, B. An Evidence-Based Approach for Choosing Post-exercise Recovery Techniques to Reduce Markers of Muscle Damage, Soreness, Fatigue, and Inflammation: A Systematic Review With Meta-Analysis. *Front. Physiol.* 9: 403, 2018.
16. Lima, KM, Carneiro, SP, de Souza Alves, D, Peixinho, CC, and de Oliveira, LF. Assessment of muscle architecture of the biceps femoris and vastus lateralis by ultrasound after a chronic stretching program. *Clin. J. Sport Med.* 25: 55-60, 2015.
17. Eliasson, J, Elfegoun, T, Nilsson, J, Kohnke, R, Ekblom, B, and Blomstrand, E. Maximal lengthening contractions increase

Chapter 5

1. Akagi, R, and Takahashi, H. Effect of a 5-week static stretching program on hardness of the gastrocnemius muscle. *Scand. J. Med. Sci. Sports* 24: 950-957, 2014.
2. Alway, SE, Winchester, PK, Davis, ME, and Gonyea, WJ. Regionalized adaptations and muscle fiber proliferation

- p70 S6 kinase phosphorylation in human skeletal muscle in the absence of nutritional supply. *Am. J. Physiol. Endocrinol. Metab.* 291: 1197-1205, 2006.
18. Ema, R, Wakahara, T, Miyamoto, N, Kanehisa, H, and Kawakami, Y. Inhomogeneous architectural changes of the quadriceps femoris induced by resistance training. *Eur. J. Appl. Physiol.* 113: 2691-2703, 2013.
 19. Ema, R, Sakaguchi, M, Akagi, R, and Kawakami, Y. Unique activation of the quadriceps femoris during single- and multi-joint exercises. *Eur. J. Appl. Physiol.* 116: 1031-1041, 2016.
 20. Evangelista, AL, De Souza, EO, Moreira, DCB, Alonso, AC, Teixeira, CVS, Wadhi, T, Rauch, J, Bocalini, DS, Pereira, PEA, and Greve, JMD. Interset Stretching vs. Traditional Strength Training: Effects on Muscle Strength and Size in Untrained Individuals. *J. Strength Cond Res.*, 2019.
 21. Figueiredo, VC, Roberts, LA, Markworth, JF, Barnett, MP, Coombes, JS, Raastad, T, Peake, JM, and Cameron-Smith, D. Impact of resistance exercise on ribosome biogenesis is acutely regulated by post-exercise recovery strategies. *Physiol. Rep.* 4: 10.14814/phy2.12670, 2016.
 22. Figueiredo, VC, and McCarthy, JJ. Regulation of Ribosome Biogenesis in Skeletal Muscle Hypertrophy. *Physiology (Bethesda)* 34: 30-42, 2019.
 23. Fink, J, Schoenfeld, BJ, Kikuchi, N, and Nakazato, K. Effects of drop set resistance training on acute stress indicators and long-term muscle hypertrophy and strength. *J. Sports Med. Phys. Fitness*, 2017.
 24. Fisher, JP, Carlson, L, Steele, J, and Smith, D. The effects of pre-exhaustion, exercise order, and rest intervals in a full-body resistance training intervention. *Appl. Physiol. Nutr. Metab.* 39: 1265-1270, 2014.
 25. Fisher, JP, Carlson, L, and Steele, J. The Effects of Breakdown Set Resistance Training on Muscular Performance and Body Composition in Young Men and Women. *J. Strength Cond Res.* 30: 1425-1432, 2016.
 26. Freitas, SR, and Mil-Homens, P. Effect of 8-week high-intensity stretching training on biceps femoris architecture. *J. Strength Cond Res.* 29: 1737-1740, 2015.
 27. Friedmann, B, Kinscherf, R, Vorwald, S, Muller, H, Kucera, K, Borisch, S, Richter, G, Bartsch, P, and Billeter, R. Muscular adaptations to computer-guided strength training with eccentric overload. *Acta Physiol. Scand.* 182: 77-88, 2004.
 28. Friedmann-Bette, B, Bauer, T, Kinscherf, R, Vorwald, S, Klute, K, Bischoff, D, Muller, H, Weber, MA, Metz, J, Kauczor, HU, Bartsch, P, and Billeter, R. Effects of strength training with eccentric overload on muscle adaptation in male athletes. *Eur. J. Appl. Physiol.* 108: 821-836, 2010.
 29. Fujita, S, Rasmussen, BB, Cadenas, JG, Grady, JJ, and Volpi, E. Effect of insulin on human skeletal muscle protein synthesis is modulated by insulin-induced changes in muscle blood flow and amino acid availability. *Am. J. Physiol. Endocrinol. Metab.* 291: E745-54, 2006.
 30. Gentil, P, Oliveira, E, de Araujo Rocha Junior, V, do Carmo, J, and Bottaro, M. Effects of exercise order on upper-body muscle activation and exercise performance. *J. Strength Cond Res.* 21: 1082-1086, 2007.
 31. Golas, A, Maszczyk, A, Pietraszewski, P, Stastny, P, Tufano, JJ, and Zajac, A. Effects of Pre-exhaustion on the Patterns of Muscular Activity in the Flat Bench Press. *J. Strength Cond Res.* 31: 1919-1924, 2017.
 32. Goto, K, Sato, K, and Takamatsu, K. A single set of low intensity resistance exercise immediately following high intensity resistance exercise stimulates growth hormone secretion in men. *J. Sports Med. Phys. Fitness* 43: 243-249, 2003.
 33. Goto, K, Nagasawa, M, Yanagisawa, O, Kizuka, T, Ishii, N, and Takamatsu, K. Muscular adaptations to combinations of high- and low-intensity resistance exercises. *J. Strength Cond Res.* 18: 730-737, 2004.
 34. Goto, K, Oda, H, Kondo, H, Igaki, M, Suzuki, A, Tsuchiya, S, Murase, T, Hase, T, Fujiya, H, Matsumoto, I, Naito, H, Sugiura, T, Ohira, Y, and Yoshioka, T. Responses of muscle mass, strength and gene transcripts to long-term heat stress in healthy human subjects. *Eur. J. Appl. Physiol.* 111: 17-27, 2011.
 35. Goto, M, Nirengi, S, Kurosawa, Y, Nagano, A, and Hamaoka, T. Effects of the Drop-set and Reverse Drop-set Methods on the Muscle Activity and Intramuscular Oxygenation of the Triceps Brachii among Trained and Untrained Individuals. *J. Sports Sci. Med.* 15: 562-568, 2016.
 36. Goto, M, Maeda, C, Hirayama, T, Terada, S, Nirengi, S, Kurosawa, Y, Nagano, A, and Hamaoka, T. Partial Range of Motion Exercise Is Effective for Facilitating Muscle Hypertrophy and Function Through Sustained Intramuscular Hypoxia in Young Trained Men. *J. Strength Cond Res.* 33: 1286-1294, 2019.
 37. Guarascio, M, Penn, C, and Sparks, C. The Effects of Pre-exhaustion of a Secondary Synergist On a Primary Mover of a Compound Exercise. *Journal of Orthopaedic & Sports Physical* 46: A178, 2016.
 38. Hafen, PS, Abbott, K, Bowden, J, Lopiano, R, Hancock, CR, and Hyldahl, RD. Daily heat treatment maintains mitochondrial function and attenuates atrophy in human skeletal muscle subjected to immobilization. *J. Appl. Physiol.* (1985) 127: 47-57, 2019.
 39. Haff, GG, Hobbs, RT, Haff, EE, Sands, WA, Pierce, KC, and Stone, MH. Cluster training: A novel method for introducing training program variation. *Strength Cond J* 1: 67-76, 2008.
 40. Horwath, O, Paulsen, G, Esping, T, Seynnes, O, and Olsson, MC. Isokinetic resistance training combined with eccentric overload improves athletic performance and induces muscle hypertrophy in young ice hockey players. *J. Sci. Med. Sport*, 2019.
 41. Iglesias-Soler, E, Carballeira, E, Sanchez-Otero, T, Mayo, X, and Fernandez-del-Olmo, M. Performance of maximum number of repetitions with cluster-set configuration. *Int. J. Sports Physiol. Perform.* 9: 637-642, 2014.
 42. Jakobi, JM, Simpson, CL, Smart, RR, and O'Connor, B. Response to Nunes and colleagues letter: The data do not seem to support the effect of stretch training in increasing MT. *Scand. J. Med. Sci. Sports* 28: 2769-2771, 2018.
 43. Junior, VA, Bottaro, M, Pereira, MC, Andrade, MM, P Junior, PR, and Carmo, JC. Electromyographic analyses of muscle pre-activation induced by single joint exercise. *Rev. Bras. Fisioter* 14: 158-165, 2010.
 44. Korak, JA, Paquette, MR, Fuller, DK, Caputo, JL, and Coons, JM. Effect of a rest-pause vs. traditional squat on electromyography and lifting volume in trained women. *Eur. J. Appl. Physiol.* 118: 1309-1314, 2018.
 45. Kubo, K, Ikebukuro, T, and Yata, H. Effects of squat training with different depths on lower limb muscle volumes. *Eur. J. Appl. Physiol.*, 2019.
 46. Leeder, J, Gissane, C, van Someren, K, Gregson, W, and Howatson, G. Cold water immersion and recovery from strenuous exercise: a meta-analysis. *Br. J. Sports Med.* 46: 233-240, 2012.
 47. Lundberg, TR, Garcia-Gutierrez, MT, Mandic, M, Lilja, M, and Fernandez-Gonzalo, R. Regional and muscle-specific adaptations in knee extensor hypertrophy using flywheel versus conventional weight-stack resistance exercise. *Appl. Physiol. Nutr. Metab.* 44: 827-833, 2019.

48. Maia, MF, Willardson, JM, Paz, GA, and Miranda, H. Effects of different rest intervals between antagonist paired sets on repetition performance and muscle activation. *J. Strength Cond Res.* 28: 2529-2535, 2014.
49. Maroto-Izquierdo, S, Garcia-Lopez, D, and de Paz, JA. Functional and Muscle-Size Effects of Flywheel Resistance Training with Eccentric-Overload in Professional Handball Players. *J. Hum. Kinet* 60: 133-143, 2017.
50. Mawhinney, C, Jones, H, Low, DA, Green, DJ, Howatson, G, and Gregson, W. Influence of cold-water immersion on limb blood flow after resistance exercise. *Eur. J. Sport. Sci.* 17: 519-529, 2017.
51. McHugh, MP. Recent advances in the understanding of the repeated bout effect: the protective effect against muscle damage from a single bout of eccentric exercise. *Scand. J. Med. Sci. Sports* 13: 88-97, 2003.
52. McMahon, G, Morse, CI, Burden, A, Winwood, K, and Onambele, GL. Muscular adaptations and insulin-like growth factor-I (IGF-I) responses to resistance training are stretch-mediated. *Muscle Nerve*, 2013.
53. McMahon, GE, Morse, CI, Burden, A, Winwood, K, and Onambele, GL. Impact of range of motion during ecologically valid resistance training protocols on muscle size, subcutaneous fat, and strength. *J. Strength Cond Res.* 28: 245-255, 2014.
54. Menetrier, A, Beliard, S, Ravier, G, Mourot, L, Bouhaddi, M, Regnard, J, and Tordi, N. Changes in femoral artery blood flow during thermoneutral, cold, and contrast-water therapy. *J. Sports Med. Phys. Fitness* 55: 768-775, 2015.
55. Merrigan, JJ, Jones, MT, and White, JB. A Comparison of Compound Set and Traditional Set Resistance Training in Women: Changes in Muscle Strength, Endurance, Quantity, and Architecture. *Journal of Science in Sport and Exercise* : 1-9, 2019.
56. Miyazaki, M, and Esser, KA. Cellular mechanisms regulating protein synthesis and skeletal muscle hypertrophy in animals. *J. Appl. Physiol.* 106: 1367-1373, 2009.
57. Moore, DR, Phillips, SM, Babraj, JA, Smith, K, and Rennie, MJ. Myofibrillar and collagen protein synthesis in human skeletal muscle in young men after maximal shortening and lengthening contractions. *Am. J. Physiol. Endocrinol. Metab.* 288: 1153-1159, 2005.
58. Nardone, A, Romano, C, and Schieppati, M. Selective recruitment of high-threshold human motor units during voluntary isotonic lengthening of active muscles. *J. Physiol. (Lond.)* 409: 451-471, 1989.
59. Nielsen, AR, and Pedersen, BK. The biological roles of exercise-induced cytokines: IL-6, IL-8, and IL-15. *Appl. Physiol. Nutr. Metab.* 32: 833-839, 2007.
60. Norrbrand, L, Fluckey, JD, Pozzo, M, and Tesch, PA. Resistance training using eccentric overload induces early adaptations in skeletal muscle size. *Eur. J. Appl. Physiol.* 102: 271-281, 2008.
61. Oliver, JM, Jagim, AR, Sanchez, AC, Mardock, MA, Kelly, KA, Meredith, HJ, Smith, GL, Greenwood, M, Parker, JL, Riechman, SE, Fluckey, JD, Crouse, SE, and Kreider, RB. Greater gains in strength and power with intraset rest intervals in hypertrophic training. *J. Strength Cond Res.* 27: 3116-3131, 2013.
62. Oliver, JM, Kreutzer, A, Jenke, S, Phillips, MD, Mitchell, JB, and Jones, MT. Acute response to cluster sets in trained and untrained men. *Eur. J. Appl. Physiol.* 115: 2383-2393, 2015.
63. Oliver, JM, Jenke, SC, Mata, JD, Kreutzer, A, and Jones, MT. Acute Effect of Cluster and Traditional Set Configurations on Myokines Associated with Hypertrophy. *Int. J. Sports Med.* 37: 1019-1024, 2016.
64. Ozaki, H, Kubota, A, Natsume, T, Loenneke, JP, Abe, T, Machida, S, and Naito, H. Effects of drop sets with resistance training on increases in muscle CSA, strength, and endurance: a pilot study. *J. Sports Sci.* : 1-6, 2017.
65. Paz, GA, Robbins, DW, de Oliveira, CG, Bottaro, M, and Miranda, H. Volume Load and Neuromuscular Fatigue During an Acute Bout of Agonist-Antagonist Paired-Set vs. Traditional-Set Training. *J. Strength Cond Res.* 31: 2777-2784, 2017.
66. Paz, GA, Maia, MF, Salerno, VP, Coburn, J, Willardson, JM, and Miranda, H. Neuromuscular responses for resistance training sessions adopting traditional, superset, paired set and circuit methods. *J. Sports Med. Phys. Fitness* , 2019.
67. Peake, JM, Roberts, LA, Figueiredo, VC, Egner, I, Krog, S, Aas, SN, Suzuki, K, Markworth, JF, Coombes, JS, Cameron-Smith, D, and Raastad, T. The effects of cold water immersion and active recovery on inflammation and cell stress responses in human skeletal muscle after resistance exercise. *J. Physiol.* 595: 695-711, 2017.
68. Pournot, H, Bieuzen, F, Louis, J, Mounier, R, Fillard, JR, Barbiche, E, and Hausswirth, C. Time-course of changes in inflammatory response after whole-body cryotherapy multi exposures following severe exercise. *PLoS One* 6: e22748, 2011.
69. Prestes, J, Tibana, RA, de Araujo Sousa, E, da Cunha Nascimento, D, de Oliveira Rocha, P, Camarco, NF, Frade de Sousa, NM, and Willardson, JM. Strength And Muscular Adaptations Following 6 Weeks Of Rest-Pause Versus Traditional Multiple-Sets Resistance Training In Trained Subjects. *J. Strength Cond Res.* , 2017.
70. Robbins, DW, Young, WB, and Behm, DG. The effect of an upper-body agonist-antagonist resistance training protocol on volume load and efficiency. *J. Strength Cond Res.* 24: 2632-2640, 2010.
71. Robbins, DW, Young, WB, Behm, DG, Payne, WR, and Klimstra, MD. Physical performance and electromyographic responses to an acute bout of paired set strength training versus traditional strength training. *J. Strength Cond Res.* 24: 1237-1245, 2010.
72. Roberts, LA, Raastad, T, Markworth, JF, Figueiredo, VC, Egner, IM, Shield, A, Cameron-Smith, D, Coombes, JS, and Peake, JM. Post-exercise cold water immersion attenuates acute anabolic signalling and long-term adaptations in muscle to strength training. *J. Physiol.* 593: 4285-4301, 2015.
73. Schoenfeld, B. The use of specialized training techniques to maximize muscle hypertrophy. *Strength Cond J* 33: 60-65, 2011.
74. Schoenfeld, BJ, Grgic, J, Contreras, B, Delcastillo, K, Alto, A, Haun, CT, De Souza, EO, and Vigotsky, AD. To flex or rest: Does adding no-load isometric actions to the inter-set rest period in resistance training enhance muscular adaptations? *Frontiers in Physiology* doi: 10.3389/fphys.2019.01571, 2019.
75. Schoenfeld, BJ. Does exercise-induced muscle damage play a role in skeletal muscle hypertrophy? *J. Strength Cond Res.* 26: 1441-1453, 2012.
76. Schoenfeld, BJ, Ogborn, D, and Krieger, JW. Dose-response relationship between weekly resistance training volume and increases in muscle mass: A systematic review and meta-analysis. *J. Sports Sci.* : 1-10, 2016.
77. Schoenfeld, BJ, Ogborn, DI, Vigotsky, AD, Franchi, MV, and Krieger, JW. Hypertrophic Effects of Concentric vs. Eccentric

- Muscle Actions: A Systematic Review and Meta-analysis. *J. Strength Cond Res.* 31: 2599-2608, 2017.
78. Shepstone, TN, Tang, JE, Dallaire, S, Schuenke, MD, Staron, RS, and Phillips, SM. Short-term high- vs. low-velocity isokinetic lengthening training results in greater hypertrophy of the elbow flexors in young men. *J. Appl. Physiol.* (1985) 98: 1768-1776, 2005.
 79. Shibata, K, Takizawa, K, Nosaka, K, and Mizuno, M. Effects of Prolonging Eccentric Phase Duration in Parallel Back-Squat Training to Momentary Failure on Muscle Cross-Sectional Area, Squat One Repetition Maximum, and Performance Tests in University Soccer Players. *J. Strength Cond Res.*, 2018.
 80. Silva, JE, Lowery, RP, Antonio, J, McCleary, S, Rauch, J, Ormes, J, Shields, K, Sharp, M, Georges, J, Weiner, S, Joy, J, and Wilson, JM. Weighted post-set stretching increases skeletal muscle hypertrophy (NSCA 2014 annual meeting. *J. Strength Cond Res* 28: 65, 2014.
 81. Simpson, CL, Kim, BDH, Bourcet, MR, Jones, GR, and Jakobi, JM. Stretch training induces unequal adaptation in muscle fascicles and thickness in medial and lateral gastrocnemii. *Scand. J. Med. Sci. Sports* 27: 1597-1604, 2017.
 82. Soares, EG, Brown, LE, Gomes, WA, Correa, DA, Serpa, EP, da Silva, JJ, Junior Gde, B, Fioravanti, GZ, Aoki, MS, Lopes, CR, and Marchetti, PH. Comparison Between Pre-Exhaustion and Traditional Exercise Order on Muscle Activation and Performance in Trained Men. *J. Sports Sci. Med.* 15: 111-117, 2016.
 83. Torres Pirauá, AL, Barros Beltrão, N, Ximenes Santos, C, Pitanguí, R, Carolina, A, and Cappato de Araújo, R. Analysis of muscle activity during the bench press exercise performed with the pre-activation method on stable and unstable surfaces. *Kinesiology: International journal of fundamental and applied kinesiology* 49: 161-168, 2017.
 84. Tufano, JJ, Brown, LE, and Haff, GG. Theoretical and Practical Aspects of Different Cluster Set Structures: A Systematic Review. *J. Strength Cond Res.* 31: 848-867, 2017.
 85. Vicens-Bordas, J, Esteve, E, Fort-Vanmeerhaeghe, A, Bandholm, T, and Thorborg, K. Is inertial flywheel resistance training superior to gravity-dependent resistance training in improving muscle strength? A systematic review with meta-analyses. *J. Sci. Med. Sport* 21: 75-83, 2018.
 86. Wackerhage, H, Schoenfeld, BJ, Hamilton, DL, Lehti, M, and Hulmi, JJ. Stimuli and sensors that initiate skeletal muscle hypertrophy following resistance exercise. *J. Appl. Physiol.* (1985) 126: 30-43, 2019.
 87. Walker, S, Blazevich, AJ, Haff, GG, Tufano, JJ, Newton, RU, and Häkkinen, K. Greater Strength Gains after Training with Accentuated Eccentric than Traditional Isoinertial Loads in Already Strength-Trained Men. *Front. Physiol.* 7: 149, 2016.
 88. Wallace, W, Ugrinowitsch, C, Stefan, M, Rauch, J, Barakat, C, Shields, K, Barninger, A, Barroso, R, and De Souza, EO. Repeated Bouts of Advanced Strength Training Techniques: Effects on Volume Load, Metabolic Responses, and Muscle Activation in Trained Individuals. *Sports (Basel)* 7: 10.3390/sports7010014, 2019.
 89. Weakley, JJS, Till, K, Read, DB, Roe, GAB, Darrall-Jones, J, Phibbs, PJ, and Jones, B. The effects of traditional, superset, and tri-set resistance training structures on perceived intensity and physiological responses. *Eur. J. Appl. Physiol.* 117: 1877-1889, 2017.
 90. Yamane, M, Ohnishi, N, and Matsumoto, T. Does Regular Post-exercise Cold Application Attenuate Trained Muscle Adaptation? *Int. J. Sports Med.* 36: 647-653, 2015.
- ## Chapter 6
1. Ackel-D'Elia, C, Carnier, J, Bueno, CR,Jr, Campos, RM, Sanches, PL, Clemente, AP, Tufik, S, de Mello, MT, and Damaso, AR. Effects of different physical exercises on leptin concentration in obese adolescents. *Int. J. Sports Med.* 35: 164-171, 2014.
 2. Adams, G, and Bamman, MM. Characterization and regulation of mechanical loading-induced compensatory muscle hypertrophy. *Comprehensive Physiology* 2829, 2012.
 3. Ahtiainen, JP, Hulmi, JJ, Kraemer, WJ, Lehti, M, Pakarinen, A, Mero, AA, Karavirta, L, Sillanpaa, E, Selanne, H, Alen, M, Komulainen, J, Kovanen, V, Nyman, K, and Häkkinen, K. Strength, [corrected] endurance or combined training elicit diverse skeletal muscle myosin heavy chain isoform proportion but unaltered androgen receptor concentration in older men. *Int. J. Sports Med.* 30: 879-887, 2009.
 4. Andersen, P, and Henriksson, J. Capillary supply of the quadriceps femoris muscle of man: adaptive response to exercise. *J. Physiol.* 270: 677-690, 1977.
 5. Apro, W, Wang, L, Ponten, M, Blomstrand, E, and Sahlin, K. Resistance exercise induced mTORC1 signaling is not impaired by subsequent endurance exercise in human skeletal muscle. *Am. J. Physiol. Endocrinol. Metab.* 305: E22-32, 2013.
 6. Atherton, PJ, Babraj, J, Smith, K, Singh, J, Rennie, MJ, and Wackerhage, H. Selective activation of AMPK-PGC-1alpha or PKB-TSC2-mTOR signaling can explain specific adaptive responses to endurance or resistance training-like electrical muscle stimulation. *FASEB J.* 19: 786-788, 2005.
 7. Babcock, L, Escano, M, D'Lugos, A, Todd, K, Murach, K, and Luden, N. Concurrent aerobic exercise interferes with the satellite cell response to acute resistance exercise. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 302: R1458-65, 2012.
 8. Bell, GJ, Syrotuik, D, Martin, TP, Burnham, R, and Quinney, HA. Effect of concurrent strength and endurance training on skeletal muscle properties and hormone concentrations in humans. *Eur. J. Appl. Physiol.* 81: 418-427, 2000.
 9. Benziane, B, Burton, TJ, Scanlan, B, Galuska, D, Canny, BJ, Chibalin, AV, Zierath, JR, and Stepto, NK. Divergent cell signaling after short-term intensified endurance training in human skeletal muscle. *Am. J. Physiol. Endocrinol. Metab.* 295: E1427-38, 2008.
 10. Bloor, CM. Angiogenesis during exercise and training. *Angiogenesis* 8: 263-271, 2005.
 11. Cadore, EL, Izquierdo, M, Pinto, SS, Alberton, CL, Pinto, RS, Baroni, BM, Vaz, MA, Lanferdini, FJ, Radaelli, R, Gonzalez-Izal, M, Bottaro, M, and Krusel, LF. Neuromuscular adaptations to concurrent training in the elderly: effects of intrasession exercise sequence. *Age (Dordr)* 35: 891-903, 2013.
 12. Carrithers, JA, Carroll, CC, Coker, RH, Sullivan, DH, and Trappe, TA. Concurrent exercise and muscle protein synthesis: implications for exercise countermeasures in space. *Aviat. Space Environ. Med.* 78: 457-462, 2007.
 13. Charifi, N, Kadi, F, Feasson, L, and Denis, C. Effects of endurance training on satellite cell frequency in skeletal muscle of old men. *Muscle Nerve* 28: 87-92, 2003.
 14. Chtara, M, Chaouachi, A, Levin, GT, Chaouachi, M, Chamari, K, Amri, M, and Laursen, PB. Effect of concurrent endurance and circuit resistance training sequence on muscular strength and power development. *J. Strength Cond Res.* 22: 1037-1045, 2008.
 15. Cogley, JN, Bartlett, JD, Kayani, A, Murray, SW, Louhelainen, J, Donovan, T, Waldron, S, Gregson, W, Burniston, JG,

- Morton, JP, and Close, GL. PGC-1 α transcriptional response and mitochondrial adaptation to acute exercise is maintained in skeletal muscle of sedentary elderly males. *Biogerontology* 13: 621-631, 2012.
16. Coffey, VG, Zhong, Z, Shield, A, Canny, BJ, Chibalin, AV, Zierath, JR, and Hawley, JA. Early signaling responses to divergent exercise stimuli in skeletal muscle from well-trained humans. *FASEB J.* 20: 190-192, 2006.
 17. Coffey, VG, Jemio, B, Edge, J, Garnham, AP, Trappe, SW, and Hawley, JA. Effect of consecutive repeated sprint and resistance exercise bouts on acute adaptive responses in human skeletal muscle. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 297: R1441-51, 2009.
 18. Coffey, VG, Pilegaard, H, Garnham, AP, O'Brien, BJ, and Hawley, JA. Consecutive bouts of diverse contractile activity alter acute responses in human skeletal muscle. *J. Appl. Physiol.* (1985) 106: 1187-1197, 2009.
 19. Coggan, AR, Spina, RJ, King, DS, Rogers, MA, Brown, M, Nemeth, PM, and Holloszy, JO. Skeletal muscle adaptations to endurance training in 60- to 70-yr-old men and women. *J. Appl. Physiol.* (1985) 72: 1780-1786, 1992.
 20. Collins, MA, and Snow, TK. Are adaptations to combined endurance and strength training affected by the sequence of training? *J. Sports Sci.* 11: 485-491, 1993.
 21. Creer, A, Gallagher, P, Slivka, D, Jemio, B, Fink, W, and Trappe, S. Influence of muscle glycogen availability on ERK1/2 and Akt signaling after resistance exercise in human skeletal muscle. *J. Appl. Physiol.* 99: 950-956, 2005.
 22. Davitt, PM, Pellegrino, JK, Schanzer, JR, Tjionas, H, and Arent, SM. The effects of a combined resistance training and endurance exercise program in inactive college female subjects: does order matter? *J. Strength Cond Res.* 28: 1937-1945, 2014.
 23. de Souza, EO, Tricoli, V, Roschel, H, Brum, PC, Bacurau, AV, Ferreira, JC, Aoki, MS, Neves, M, Jr, Aihara, AY, da Rocha Correa Fernandes, A, and Ugrinowitsch, C. Molecular adaptations to concurrent training. *Int. J. Sports Med.* 34: 207-213, 2013.
 24. de Souza, EO, Tricoli, V, Aoki, MS, Roschel, H, Brum, PC, Bacurau, AV, Silva-Batista, C, Wilson, JM, Neves, M, Jr, Soares, AG, and Ugrinowitsch, C. Effects of concurrent strength and endurance training on genes related to myostatin signaling pathway and muscle fiber responses. *J. Strength Cond Res.* 28: 3215-3223, 2014.
 25. Di Donato, DM, West, DW, Churchward-Venne, TA, Breen, L, Baker, SK, and Phillips, SM. Influence of aerobic exercise intensity on myofibrillar and mitochondrial protein synthesis in young men during early and late postexercise recovery. *Am. J. Physiol. Endocrinol. Metab.* 306: E1025-32, 2014.
 26. Donges, CE, Burd, NA, Duffield, R, Smith, GC, West, DW, Short, MJ, Mackenzie, R, Plank, LD, Shepherd, PR, Phillips, SM, and Edge, JA. Concurrent resistance and aerobic exercise stimulates both myofibrillar and mitochondrial protein synthesis in sedentary middle-aged men. *J. Appl. Physiol.* (1985) 112: 1992-2001, 2012.
 27. Dreyer, HC, Fujita, S, Cadenas, JG, Chinkes, DL, Volpi, E, and Rasmussen, BB. Resistance exercise increases AMPK activity and reduces 4E-BP1 phosphorylation and protein synthesis in human skeletal muscle. *J. Physiol.* 576: 613-624, 2006.
 28. Edstrom, L, and Ekblom, B. Differences in sizes of red and white muscle fibres in vastus lateralis of musculus quadriceps femoris of normal individuals and athletes. Relation to physical performance. *Scand. J. Clin. Lab. Invest.* 30: 175-181, 1972.
 29. Farup, J, Kjolhede, T, Sorensen, H, Dalgas, U, Moller, AB, Vestergaard, PE, Ringgaard, S, Bojsen-Moller, J, and Vissing, K. Muscle morphological and strength adaptations to endurance vs. resistance training. *J. Strength Cond Res.* 26: 398-407, 2012.
 30. Ferrara, CM, Goldberg, AP, Ortmeier, HK, and Ryan, AS. Effects of aerobic and resistive exercise training on glucose disposal and skeletal muscle metabolism in older men. *J. Gerontol. A Biol. Sci. Med. Sci.* 61: 480-487, 2006.
 31. Fyfe, JJ, Bishop, DJ, and Stepto, NK. Interference between concurrent resistance and endurance exercise: molecular bases and the role of individual training variables. *Sports Med.* 44: 743-762, 2014.
 32. Fyfe, JJ, Bishop, DJ, Bartlett, JD, Hanson, ED, Anderson, MJ, Garnham, AP, and Stepto, NK. Enhanced skeletal muscle ribosome biogenesis, yet attenuated mTORC1 and ribosome biogenesis-related signalling, following short-term concurrent versus single-mode resistance training. *Sci. Rep.* 8: 560-017-18887-6, 2018.
 33. Gollnick, PD, and Saltin, B. Significance of skeletal muscle oxidative enzyme enhancement with endurance training. *Clin. Physiol.* 2: 1-12, 1982.
 34. Goodman, CA, Mayhew, DL, and Hornberger, TA. Recent progress toward understanding the molecular mechanisms that regulate skeletal muscle mass. *Cell. Signal.* 23: 1896-1906, 2011.
 35. Gravelle, BL, and Blessing, DL. Physiological adaptation in women concurrently training for strength and endurance. *J. Strength Cond Res* 14: 5-13, 2000.
 36. Grgic, J, Mclivenna, LC, Fyfe, JJ, Sabol, E, Bishop, DJ, Schoenfeld, BJ, and Pedisic, Z. Does Aerobic Training Promote the Same Skeletal Muscle Hypertrophy as Resistance Training? A Systematic Review and Meta-Analysis. *Sports Med.* 49: 233-254, 2019.
 37. Harber, MP, Konopka, AR, Douglass, MD, Minchev, K, Kaminsky, LA, Trappe, TA, and Trappe, S. Aerobic exercise training improves whole muscle and single myofiber size and function in older women. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 297: R1452-9, 2009.
 38. Harber, MP, Konopka, AR, Undem, MK, Hinkley, JM, Minchev, K, Kaminsky, LA, Trappe, TA, and Trappe, S. Aerobic exercise training induces skeletal muscle hypertrophy and age-dependent adaptations in myofiber function in young and older men. *J. Appl. Physiol.* (1985) 113: 1495-1504, 2012.
 39. Hepple, RT, Mackinnon, SL, Goodman, JM, Thomas, SG, and Plyley, MJ. Resistance and aerobic training in older men: effects on VO₂peak and the capillary supply to skeletal muscle. *J. Appl. Physiol.* (1985) 82: 1305-1310, 1997.
 40. Hickson, RC. Interference of strength development by simultaneously training for strength and endurance. *Eur. J. Appl. Physiol. Occup. Physiol.* 45: 255-263, 1980.
 41. Hoppeler, H. Exercise-induced ultrastructural changes in skeletal muscle. *Int. J. Sports Med.* 7: 187-204, 1986.
 42. Hudelmaier, M, Wirth, W, Himmer, M, Ring-Dimitriou, S, Sanger, A, and Eckstein, F. Effect of exercise intervention on thigh muscle volume and anatomical cross-sectional areas--quantitative assessment using MRI. *Magn. Reson. Med.* 64: 1713-1720, 2010.
 43. Izquierdo, M, Ibanez, J, Häkkinen, K, Kraemer, WJ, Larrion, JL, and Gorostiaga, EM. Once weekly combined resistance and cardiovascular training in healthy older men. *Med. Sci. Sports Exerc.* 36: 435-443, 2004.
 44. Izquierdo, M, Häkkinen, K, Ibanez, J, Kraemer, WJ, and Gorostiaga, EM. Effects of combined resistance and cardiovascular training on strength, power, muscle cross-sectional area, and endurance markers in middle-aged men. *Eur. J. Appl. Physiol.* 94: 70-75, 2005.

45. Jagatheesan, A. Acute Effect of Continuous and Intermittent Cycling on Maximum Strength in Non-Athlete Females. *JPBMS* 8: 1-5, 2011.
46. Jones, TW, Howatson, G, Russell, M, and French, DN. Performance and neuromuscular adaptations following differing ratios of concurrent strength and endurance training. *J. Strength Cond Res.* 27: 3342-3351, 2013.
47. Jubrias, SA, Esselman, PC, Price, LB, Cress, ME, and Conley, KE. Large energetic adaptations of elderly muscle to resistance and endurance training. *J. Appl. Physiol.* (1985) 90: 1663-1670, 2001.
48. Karavirta, L, Häkkinen, A, Sillanpaa, E, Garcia-Lopez, D, Kauhanen, A, Haapasaari, A, Alen, M, Pakarinen, A, Kraemer, WJ, Izquierdo, M, Gorostiaga, E, and Häkkinen, K. Effects of combined endurance and strength training on muscle strength, power and hypertrophy in 40-67-year-old men. *Scand. J. Med. Sci. Sports* 21: 402-411, 2011.
49. Kohn, TA, Essen-Gustavsson, B, and Myburgh, KH. Specific muscle adaptations in type II fibers after high-intensity interval training of well-trained runners. *Scand. J. Med. Sci. Sports* 21: 765-772, 2011.
50. Konopka, AR, and Harber, MP. Skeletal Muscle Hypertrophy after Aerobic Exercise Training. *Exerc. Sport Sci. Rev.*, 2014.
51. Koopman, R, Zorenc, AH, Gransier, RJ, Cameron-Smith, D, and van Loon, LJ. Increase in S6K1 phosphorylation in human skeletal muscle following resistance exercise occurs mainly in type II muscle fibers. *Am. J. Physiol. Endocrinol. Metab.* 290: E1245-52, 2006.
52. Kraemer, WJ, Patton, JF, Gordon, SE, Harman, EA, Deschenes, MR, Reynolds, K, Newton, RU, Triplett, NT, and Dziados, JE. Compatibility of high-intensity strength and endurance training on hormonal and skeletal muscle adaptations. *J. Appl. Physiol.* (1985) 78: 976-989, 1995.
53. Lundberg, TR, Fernandez-Gonzalo, R, Gustafsson, T, and Tesch, PA. Aerobic exercise alters skeletal muscle molecular responses to resistance exercise. *Med. Sci. Sports Exerc.* 44: 1680-1688, 2012.
54. Lundberg, TR, Fernandez-Gonzalo, R, and Tesch, PA. Exercise-induced AMPK activation does not interfere with muscle hypertrophy in response to resistance training in men. *J. Appl. Physiol.* (1985) 116: 611-620, 2014.
55. Mascher, H, Andersson, H, Nilsson, PA, Ekblom, B, and Blomstrand, E. Changes in signalling pathways regulating protein synthesis in human muscle in the recovery period after endurance exercise. *Acta Physiol. (Oxf)* 191: 67-75, 2007.
56. Mascher, H, Ekblom, B, Rooyackers, O, and Blomstrand, E. Enhanced rates of muscle protein synthesis and elevated mTOR signalling following endurance exercise in human subjects. *Acta Physiol. (Oxf)* 202: 175-184, 2011.
57. McCarthy, JP, Pozniak, MA, and Agre, JC. Neuromuscular adaptations to concurrent strength and endurance training. *Med. Sci. Sports Exerc.* 34: 511-519, 2002.
58. Mikkola, J, Rusko, H, Izquierdo, M, Gorostiaga, EM, and Häkkinen, K. Neuromuscular and cardiovascular adaptations during concurrent strength and endurance training in untrained men. *Int. J. Sports Med.* 33: 702-710, 2012.
59. Mora-Rodriguez, R, Sanchez-Roncero, A, Fernandez-Elias, VE, Guadalupe-Grau, A, Ortega, JF, Dela, F, and Helge, JW. Aerobic Exercise Training Increases Muscle Water Content in Obese Middle-Age Men. *Med. Sci. Sports Exerc.* 48: 822-828, 2016.
60. Nelson, AG, Arnall, DA, Loy, SE, Silvester, LJ, and Conlee, RK. Consequences of combining strength and endurance training regimens. *Phys. Ther.* 70: 287-294, 1990.
61. Panissa, VL, Tricoli, VA, Julio, UF, Da Silva, NR, Neto, RM, Carmo, EC, and Franchini, E. Acute effect of high-intensity aerobic exercise performed on treadmill and cycle ergometer on strength performance. *J. Strength Cond Res.*, 2014.
62. Poehlman, ET, Dvorak, RV, DeNino, WF, Brochu, M, and Ades, PA. Effects of resistance training and endurance training on insulin sensitivity in nonobese, young women: a controlled randomized trial. *J. Clin. Endocrinol. Metab.* 85: 2463-2468, 2000.
63. Schwartz, RS, Shuman, WP, Larson, V, Cain, KC, Fellingham, GW, Beard, JC, Kahn, SE, Stratton, JR, Cerqueira, MD, and Abrass, IB. The effect of intensive endurance exercise training on body fat distribution in young and older men. *Metabolism* 40: 545-551, 1991.
64. Sillanpaa, E, Häkkinen, A, Nyman, K, Mattila, M, Cheng, S, Karavirta, L, Laaksonen, DE, Huuhka, N, Kraemer, WJ, and Häkkinen, K. Body composition and fitness during strength and/or endurance training in older men. *Med. Sci. Sports Exerc.* 40: 950-958, 2008.
65. Silva, RF, Cadore, EL, Kothe, G, Guedes, M, Alberton, CL, Pinto, SS, Pinto, RS, Trindade, G, and Kruel, LF. Concurrent training with different aerobic exercises. *Int. J. Sports Med.* 33: 627-634, 2012.
66. Sipila, S, and Suominen, H. Effects of strength and endurance training on thigh and leg muscle mass and composition in elderly women. *J. Appl. Physiol.* (1985) 78: 334-340, 1995.
67. Sipila, S, and Suominen, H. Effects of strength and endurance training on thigh and leg muscle mass and composition in elderly women. *J. Appl. Physiol.* (1985) 78: 334-340, 1995.
68. Sipila, S, Elorinne, M, Alen, M, Suominen, H, and Kovanen, V. Effects of strength and endurance training on muscle fibre characteristics in elderly women. *Clin. Physiol.* 17: 459-474, 1997.
69. Stepto, NK, Coffey, VG, Carey, AL, Ponnampalam, AP, Canny, BJ, Powell, D, and Hawley, JA. Global gene expression in skeletal muscle from well-trained strength and endurance athletes. *Med. Sci. Sports Exerc.* 41: 546-565, 2009.
70. Timmons, JA. Variability in training-induced skeletal muscle adaptation. *J. Appl. Physiol.* 110: 846-853, 2011.
71. Tomiya, S, Kikuchi, N, and Nakazato, K. Moderate Intensity Cycling Exercise after Upper Extremity Resistance Training Interferes Response to Muscle Hypertrophy but Not Strength Gains. *J. Sports Sci. Med.* 16: 391-395, 2017.
72. Trappe, S, Harber, M, Creer, A, Gallagher, P, Slivka, D, Minchev, K, and Whitsett, D. Single muscle fiber adaptations with marathon training. *J. Appl. Physiol.* (1985) 101: 721-727, 2006.
73. Tsitkanou, S, Spengos, K, Stasinaki, AN, Zaras, N, Bogdanis, G, Papadimas, G, and Terzis, G. Effects of high-intensity interval cycling performed after resistance training on muscle strength and hypertrophy. *Scand. J. Med. Sci. Sports* 27: 1317-1327, 2017.
74. Turner, DL, Hoppeler, H, Claassen, H, Vock, P, Kayser, B, Schena, F, and Ferretti, G. Effects of endurance training on oxidative capacity and structural composition of human arm and leg muscles. *Acta Physiol. Scand.* 161: 459-464, 1997.
75. van Wessel, T, de Haan, A, van der Laarse, WJ, and Jaspers, RT. The muscle fiber type-fiber size paradox: hypertrophy or oxidative metabolism? *Eur. J. Appl. Physiol.* 110: 665-694, 2010.
76. Vissing, K, McGee, S, Farup, J, Kjolhede, T, Vendelbo, M, and Jessen, N. Differentiated mTOR but not AMPK signaling after strength vs endurance exercise in training-accustomed individuals. *Scand. J. Med. Sci. Sports* 23: 355-366, 2013.

77. Wilkinson, SB, Phillips, SM, Atherton, PJ, Patel, R, Yarasheski, KE, Tarnopolsky, MA, and Rennie, MJ. Differential effects of resistance and endurance exercise in the fed state on signalling molecule phosphorylation and protein synthesis in human muscle. *J. Physiol.* 586: 3701-3717, 2008.
 78. Willis, LH, Slentz, CA, Bateman, LA, Shields, AT, Piner, LW, Bales, CW, Houmard, JA, and Kraus, WE. Effects of aerobic and/or resistance training on body mass and fat mass in overweight or obese adults. *J. Appl. Physiol.* (1985) 113: 1831-1837, 2012.
 79. Wilson, JM, Marin, PJ, Rhea, MR, Wilson, SM, Loenneke, JP, and Anderson, JC. Concurrent training: a meta-analysis examining interference of aerobic and resistance exercises. *J. Strength Cond Res.* 26: 2293-2307, 2012.
- ## Chapter 7
1. Abe, T, DeHoyos, DV, Pollock, ML, and Garzarella, L. Time course for strength and muscle thickness changes following upper and lower body resistance training in men and women. *Eur. J. Appl. Physiol.* 81: 174-180, 2000.
 2. Abernethy, PJ, Jurimae, J, Logan, PA, Taylor, AW, and Thayer, RE. Acute and chronic response of skeletal muscle to resistance exercise. *Sports Med.* 17: 22-38, 1994.
 3. Ahtiainen, JP, Pakarinen, A, Alen, M, Kraemer, WJ, and Häkkinen, K. Short vs. long rest period between the sets in hypertrophic resistance training: influence on muscle strength, size, and hormonal adaptations in trained men. *J. Strength Cond Res* 19: 572-582, 2005.
 4. Alway, SE, Grumbt, WH, Gonyea, WJ, and Stray-Gundersen, J. Contrasts in muscle and myofibers of elite male and female bodybuilders. *J. Appl. Physiol.* (1985) 67: 24-31, 1989.
 5. Alway, SE, Grumbt, WH, Stray-Gundersen, J, and Gonyea, WJ. Effects of resistance training on elbow flexors of highly competitive bodybuilders. *J. Appl. Physiol.* (1985) 72: 1512-1521, 1992.
 6. Bamman, MM, Hill, VJ, Adams, GR, Haddad, F, Wetzstein, CJ, Gower, BA, Ahmed, A, and Hunter, GR. Gender differences in resistance-training-induced myofiber hypertrophy among older adults. *J. Gerontol. A Biol. Sci. Med. Sci.* 58: 108-116, 2003.
 7. Bamman, MM, Petrella, JK, Kim, JS, Mayhew, DL, and Cross, JM. Cluster analysis tests the importance of myogenic gene expression during myofiber hypertrophy in humans. *J. Appl. Physiol.* 102: 2232-2239, 2007.
 8. Bellamy, LM, Joannis, S, Grubb, A, Mitchell, CJ, McKay, BR, Phillips, SM, Baker, S, and Parise, G. The Acute Satellite Cell Response and Skeletal Muscle Hypertrophy following Resistance Training. *PLoS One* 9: e109739, 2014.
 9. Bentwich, I, Avniel, A, Karov, Y, Aharonov, R, Gilad, S, Barad, O, Barzilai, A, Einat, P, Einav, U, Meiri, E, Sharon, E, Spector, Y, and Bentwich, Z. Identification of hundreds of conserved and nonconserved human microRNAs. *Nat. Genet.* 37: 766-770, 2005.
 10. Bickel, CS, Cross, JM, and Bamman, MM. Exercise dosing to retain resistance training adaptations in young and older adults. *Med. Sci. Sports Exerc.* 43: 1177-1187, 2011.
 11. Breen, L, and Phillips, SM. Skeletal muscle protein metabolism in the elderly: Interventions to counteract the 'anabolic resistance' of ageing. *Nutr. Metab. (Lond)* 8: 68-7075-8-68, 2011.
 12. Buford, TW, Anton, SD, Judge, AR, Marzetti, E, Wohlgemuth, SE, Carter, CS, Leeuwenburgh, C, Pahor, M, and Manini, TM. Models of accelerated sarcopenia: critical pieces for solving the puzzle of age-related muscle atrophy. *Ageing Res. Rev.* 9: 369-383, 2010.
 13. Burton, LA, and Sumukadas, D. Optimal management of sarcopenia. *Clin. Interv. Aging* 5: 217-228, 2010.
 14. Burton, LC, Shapiro, S, and German, PS. Determinants of physical activity initiation and maintenance among community-dwelling older persons. *Prev. Med.* 29: 422-430, 1999.
 15. Charette, SL, McEvoy, L, Pyka, G, Snow-Harter, C, Guido, D, Wiswell, RA, and Marcus, R. Muscle hypertrophy response to resistance training in older women. *J. Appl. Physiol.* (1985) 70: 1912-1916, 1991.
 16. Churchward-Venne, TA, Tieland, M, Verdijk, LB, Leenders, M, Dirks, ML, de Groot, LC, and van Loon, LJ. There Are No Nonresponders to Resistance-Type Exercise Training in Older Men and Women. *J. Am. Med. Dir. Assoc.* 16: 400-411, 2015.
 17. Coffey, VG, Zhong, Z, Shield, A, Canny, BJ, Chibalin, AV, Zierath, JR, and Hawley, JA. Early signaling responses to divergent exercise stimuli in skeletal muscle from well-trained humans. *FASEB J.* 20: 190-192, 2006.
 18. Damas, F, Phillips, SM, Libardi, CA, Vechin, FC, Lixandrao, ME, Jannig, PR, Costa, LA, Bacurau, AV, Snijders, T, Parise, G, Tricoli, V, Roschel, H, and Ugrasowitsch, C. Resistance training-induced changes in integrated myofibrillar protein synthesis are related to hypertrophy only after attenuation of muscle damage. *J. Physiol.* 594: 5209-5222, 2016.
 19. Davidsen, PK, Gallagher, IJ, Hartman, JW, Tarnopolsky, MA, Dela, F, Helge, JW, Timmons, JA, and Phillips, SM. High responders to resistance exercise training demonstrate differential regulation of skeletal muscle microRNA expression. *J. Appl. Physiol.* 110: 309-317, 2011.
 20. Devaney, JM, Tosi, LL, Fritz, DT, Gordish-Dressman, HA, Jiang, S, Orkunoglu-Suer, FE, Gordon, AH, Harmon, BT, Thompson, PD, Clarkson, PM, Angelopoulos, TJ, Gordon, PM, Moyna, NM, Pescatello, LS, Visich, PS, Zoeller, RE, Brandoli, C, Hoffman, EP, and Rogers, MB. Differences in fat and muscle mass associated with a functional human polymorphism in a post-transcriptional BMP2 gene regulatory element. *J. Cell. Biochem.* 107: 1073-1082, 2009.
 21. Dieli-Conwright, CM, Spector, TM, Rice, JC, Sattler, FR, and Schroeder, ET. Influence of hormone replacement therapy on eccentric exercise induced myogenic gene expression in postmenopausal women. *J. Appl. Physiol.* (1985) 107: 1381-1388, 2009.
 22. Dumont, NA, Wang, YX, and Rudnicki, MA. Intrinsic and extrinsic mechanisms regulating satellite cell function. *Development* 142: 1572-1581, 2015.
 23. Erskine, RM, Williams, AG, Jones, DA, Stewart, CE, and Degens, H. The individual and combined influence of ACE and ACTN3 genotypes on muscle phenotypes before and after strength training. *Scand. J. Med. Sci. Sports* 24: 642-648, 2014.
 24. Evans, W. Functional and metabolic consequences of sarcopenia. *J. Nutr.* 127: 998S-1003S, 1997.
 25. Fell, J, and Williams, D. The effect of aging on skeletal-muscle recovery from exercise: possible implications for aging athletes. *J. Aging Phys. Act.* 16: 97-115, 2008.
 26. Fisher, J, Steele, J, Bruce-Low, S, and Smith, D. Evidence-based resistance training recommendations. *Med Sportiva* 15: 147-162, 2011.
 27. Fry, CS, Drummond, MJ, Glynn, EL, Dickinson, JM, Gundermann, DM, Timmerman, KL, Walker, DK, Dhanani, S, Volpi, E, and Rasmussen, BB. Aging impairs contraction-induced human skeletal muscle mTORC1 signaling and protein synthesis. *Skelet Muscle* 1: 11-5040-1-11, 2011.

28. Galpin, AJ, Fry, AC, Nicoll, JX, Moore, CA, Schilling, BK, and Thomason, DB. Resting extracellular signal-regulated protein kinase 1/2 expression following a continuum of chronic resistance exercise training paradigms. *Res. Sports Med.* 24: 298-303, 2016.
29. Garg, K, and Boppart, MD. Influence of exercise and aging on extracellular matrix composition in the skeletal muscle stem cell niche. *J. Appl. Physiol.* (1985) 121: 1053-1058, 2016.
30. Gordon, EH, Peel, NM, Samanta, M, Theou, O, Howlett, SE, and Hubbard, RE. Sex differences in frailty: A systematic review and meta-analysis. *Exp. Gerontol.* 89: 30-40, 2017.
31. Grgić, J, and Schoenfeld, BJ. A case for considering age and sex when prescribing rest intervals in resistance training. *Kinesiology* 51: 78-82, 2019.
32. Häkkinen, K, Newton, RU, Gordon, SE, McCormick, M, Volek, JS, Nindl, BC, Gotshalk, LA, Campbell, WW, Evans, WJ, Häkkinen, A, Humphries, BJ, and Kraemer, WJ. Changes in muscle morphology, electromyographic activity, and force production characteristics during progressive strength training in young and older men. *J. Gerontol. A Biol. Sci. Med. Sci.* 53: B415-23, 1998.
33. Hand, BD, Kostek, MC, Ferrell, RE, Delmonico, MJ, Douglass, LW, Roth, SM, Hagberg, JM, and Hurley, BF. Influence of promoter region variants of insulin-like growth factor pathway genes on the strength-training response of muscle phenotypes in older adults. *J. Appl. Physiol.* 103: 1678-1687, 2007.
34. Hansen, M, and Kjaer, M. Influence of sex and estrogen on musculotendinous protein turnover at rest and after exercise. *Exerc. Sport Sci. Rev.* 42: 183-192, 2014.
35. Haun, CT, Vann, CG, Mobley, CB, Osburn, SC, Mumford, PW, Roberson, PA, Romero, MA, Fox, CD, Parry, HA, Kavazis, AN, Moon, JR, Young, KC, and Roberts, MD. Pre-training Skeletal Muscle Fiber Size and Predominant Fiber Type Best Predict Hypertrophic Responses to 6 Weeks of Resistance Training in Previously Trained Young Men. *Front. Physiol.* 10: 297, 2019.
36. Hubal, MJ, Gordish-Dressman, H, Thompson, PD, Price, TB, Hoffman, EP, Angelopoulos, TJ, Gordon, PM, Moyna, NM, Pescatello, LS, Visich, PS, Zoeller, RF, Seip, RL, and Clarkson, PM. Variability in muscle size and strength gain after unilateral resistance training. *Med. Sci. Sports Exerc.* 37: 964-972, 2005.
37. Ivey, FM, Roth, SM, Ferrell, RE, Tracy, BL, Lemmer, JT, Hurlbut, DE, Martel, GF, Siegel, EL, Fozard, JL, Jeffrey Metter, E, Fleg, JL, and Hurley, BF. Effects of age, gender, and myostatin genotype on the hypertrophic response to heavy resistance strength training. *J. Gerontol. A Biol. Sci. Med. Sci.* 55: M641-8, 2000.
38. Kadi, F, Charifi, N, Denis, C, and Lexell, J. Satellite cells and myonuclei in young and elderly women and men. *Muscle Nerve* 29: 120-127, 2004.
39. Karlsen, A, Bechshoft, RL, Malmgaard-Clausen, NM, Andersen, JL, Schjerling, P, Kjaer, M, and Mackey, AL. Lack of muscle fibre hypertrophy, myonuclear addition, and satellite cell pool expansion with resistance training in 83-94-year-old men and women. *Acta Physiol. (Oxf)* 227: e13271, 2019.
40. Kilikevicius, A, Bungler, L, and Lionikas, A. Baseline Muscle Mass Is a Poor Predictor of Functional Overload-Induced Gain in the Mouse Model. *Front. Physiol.* 7: 534, 2016.
41. Konopka, AR, and Harber, MP. Skeletal Muscle Hypertrophy after Aerobic Exercise Training. *Exerc. Sport Sci. Rev.* , 2014.
42. Kosek, DJ, Kim, JS, Petrella, JK, Cross, JM, and Bamman, MM. Efficacy of 3 days/wk resistance training on myofiber hypertrophy and myogenic mechanisms in young vs. older adults. *J. Appl. Physiol.* 101: 531-544, 2006.
43. Kumar, V, Selby, A, Rankin, D, Patel, R, Atherton, P, Hildebrandt, W, Williams, J, Smith, K, Seynnes, O, Hiscock, N, and Rennie, MJ. Age-related differences in the dose-response relationship of muscle protein synthesis to resistance exercise in young and old men. *J. Physiol.* 587: 211-217, 2009.
44. Lexell, J, Henriksson-Larsen, K, Winblad, B, and Sjostrom, M. Distribution of different fiber types in human skeletal muscles: effects of aging studied in whole muscle cross sections. *Muscle Nerve* 6: 588-595, 1983.
45. Lexell, J, Downham, D, and Sjostrom, M. Distribution of different fibre types in human skeletal muscles. Fibre type arrangement in m. vastus lateralis from three groups of healthy men between 15 and 83 years. *J. Neurol. Sci.* 72: 211-222, 1986.
46. Lixandrao, ME, Damas, F, Chacon-Mikahil, MP, Cavaglieri, CR, Ugrinowitsch, C, Bottaro, M, Vechin, FC, Conceicao, MS, Berton, R, and Libardi, CA. Time Course of Resistance Training-Induced Muscle Hypertrophy in the Elderly. *J. Strength Cond Res.* 30: 159-163, 2016.
47. Lysenko, EA, Popov, DV, Vepkhvadze, TF, Sharova, AP, and Vinogradova, OL. Moderate-Intensity Strength Exercise to Exhaustion Results in More Pronounced Signaling Changes in Skeletal Muscles of Strength-Trained Compared With Untrained Individuals. *J. Strength Cond Res.* , 2018.
48. MacDougall, JD, Sale, DG, Alway, SE, and Sutton, JR. Muscle fiber number in biceps brachii in bodybuilders and control subjects. *J. Appl. Physiol.* 57: 1399-1403, 1984.
49. MacDougall, JD, Gibala, MJ, Tarnopolsky, MA, MacDonald, JR, Interisano, SA, and Yarasheski, KE. The time course for elevated muscle protein synthesis following heavy resistance exercise. *Can. J. Appl. Physiol.* 20: 480-486, 1995.
50. Mero, AA, Hulmi, JJ, Salmijarvi, H, Katajavaruri, M, Haverinen, M, Holviala, J, Ridanpaa, T, Häkkinen, K, Kovanen, V, Ahtiainen, JP, and Selanne, H. Resistance training induced increase in muscle fiber size in young and older men. *Eur. J. Appl. Physiol.* 113: 641-650, 2013.
51. Miller, BF, Olesen, JL, Hansen, M, Dossing, S, Cramer, RM, Welling, RJ, Langberg, H, Flyvbjerg, A, Kjaer, M, Babraj, JA, Smith, K, and Rennie, MJ. Coordinated collagen and muscle protein synthesis in human patella tendon and quadriceps muscle after exercise. *J. Physiol.* 567: 1021-1033, 2005.
52. Moritani, T, and deVries, HA. Potential for gross muscle hypertrophy in older men. *J. Gerontol.* 35: 672-682, 1980.
53. Nader, GA, von Walden, F, Liu, C, Lindvall, J, Gutmann, L, Pistilli, EE, and Gordon, PM. Resistance exercise training modulates acute gene expression during human skeletal muscle hypertrophy. *J. Appl. Physiol.* (1985) 116: 693-702, 2014.
54. Negaresh, R, Ranjbar, R, Baker, JS, Habibi, A, Mokhtarzade, M, Gharibvand, MM, and Fokin, A. Skeletal Muscle Hypertrophy, Insulin-like Growth Factor 1, Myostatin and Follistatin in Healthy and Sarcopenic Elderly Men: The Effect of Whole-body Resistance Training. *Int. J. Prev. Med.* 10: 29, 2019.
55. Ntanas-Stathopoulos, J, Tzanninis, JG, Philippou, A, and Koutsilieris, M. Epigenetic regulation on gene expression induced by physical exercise. *J. Musculoskelet Neuronal Interact.* 1: 133-146, 2013.
56. Ogasawara, R, Kobayashi, K, Tsutaki, A, Lee, K, Abe, T, Fujita, S, Nakazato, K, and Ishii, N. mTOR signaling response to resistance exercise is altered by chronic resistance training and detraining in skeletal muscle. *J. Appl. Physiol.* , 2013.
57. Ogawa, K, Sanada, K, Machida, S, Okutsu, M, and Suzuki, K. Resistance exercise training-induced muscle hypertrophy was

- associated with reduction of inflammatory markers in elderly women. *Mediators Inflamm.* 2010: 171023, 2010.
58. Perez-Lopez, A, McKendry, J, Martin-Rincon, M, Morales-Alamo, D, Perez-Kohler, B, Valades, D, Bujan, J, Calbet, JAL, and Breen, L. Skeletal muscle IL-15/IL-15Ralpha and myofibrillar protein synthesis after resistance exercise. *Scand. J. Med. Sci. Sports* 28: 116-125, 2018.
 59. Pescatello, LS, Devaney, JM, Hubal, MJ, Thompson, PD, and Hoffman, EP. Highlights from the functional single nucleotide polymorphisms associated with human muscle size and strength or FAMuSS study. *Biomed. Res. Int.* 2013: 643575, 2013.
 60. Peterson, MD, Rhea, MR, and Alvar, BA. Applications of the dose-response for muscular strength development: a review of meta-analytic efficacy and reliability for designing training prescription. *J. Strength Cond Res.* 19: 950-958, 2005.
 61. Peterson, MD, Sen, A, and Gordon, PM. Influence of resistance exercise on lean body mass in aging adults: a meta-analysis. *Med. Sci. Sports Exerc.* 43: 249-258, 2011.
 62. Petrella, JK, Kim, J, Mayhew, DL, Cross, JM, and Bamman, MM. Potent myofiber hypertrophy during resistance training in humans is associated with satellite cell-mediated myonuclear addition: a cluster analysis. *J. Appl. Physiol.* 104: 1736-1742, 2008.
 63. Phillips, SM, Tipton, KD, Aarsland, A, Wolf, SE, and Wolfe, RR. Mixed muscle protein synthesis and breakdown after resistance exercise in humans. *Am. J. Physiol.* 273: E99-107, 1997.
 64. Pistilli, EE, Devaney, JM, Gordish-Dressman, H, Bradbury, MK, Seip, RL, Thompson, PD, Angelopoulos, TJ, Clarkson, PM, Moyna, NM, Pescatello, LS, Visich, PS, Zoeller, RE, Gordon, PM, and Hoffman, EP. Interleukin-15 and interleukin-15R alpha SNPs and associations with muscle, bone, and predictors of the metabolic syndrome. *Cytokine* 43: 45-53, 2008.
 65. Pistilli, EE, and Quinn, LS. From anabolic to oxidative: reconsidering the roles of IL-15 and IL-15Ralpha in skeletal muscle. *Exerc. Sport Sci. Rev.* 41: 100-106, 2013.
 66. Pollanen, E, Ronkainen, PH, Suominen, H, Takala, T, Koskinen, S, Puolakka, J, Sipila, S, and Kovanen, V. Muscular transcriptome in postmenopausal women with or without hormone replacement. *Rejuvenation Res.* 10: 485-500, 2007.
 67. Renault, V, Thornell, LE, Eriksson, PO, Butler-Browne, G, and Mouly, V. Regenerative potential of human skeletal muscle during aging. *Aging Cell.* 1: 132-139, 2002.
 68. Riechman, SE, Balasekaran, G, Roth, SM, and Ferrell, RE. Association of interleukin-15 protein and interleukin-15 receptor genetic variation with resistance exercise training responses. *J. Appl. Physiol.* 97: 2214-2219, 2004.
 69. Rossetti, ML, Steiner, JL, and Gordon, BS. Androgen-mediated regulation of skeletal muscle protein balance. *Mol. Cell. Endocrinol.* 447: 35-44, 2017.
 70. Roth, SM, Martel, GF, Ivey, FM, Lemmer, JT, Metter, EJ, Hurley, BF, and Rogers, MA. Skeletal muscle satellite cell populations in healthy young and older men and women. *Anat. Rec.* 260: 351-358, 2000.
 71. Roth, SM, Ivey, FM, Martel, GF, Lemmer, JT, Hurlbut, DE, Siegel, EL, Metter, EJ, Fleg, JL, Fozard, JL, Kostek, MC, Wernick, DM, and Hurley, BF. Muscle size responses to strength training in young and older men and women. *J. Am. Geriatr. Soc.* 49: 1428-1433, 2001.
 72. Sale, DG, MacDougall, JD, Alway, SE, and Sutton, JR. Voluntary strength and muscle characteristics in untrained men and women and male bodybuilders. *J. Appl. Physiol.* (1985) 62: 1786-1793, 1987.
 73. Schoenfeld, BJ, Ratamess, NA, Peterson, MD, Contreras, B, Tiryaki-Sonmez, G, and Alvar, BA. Effects of different volume-equated resistance training loading strategies on muscular adaptations in well-trained men. *J. Strength Cond Res.* 28: 2909-2918, 2014.
 74. Schoenfeld, BJ, Peterson, MD, Ogborn, D, Contreras, B, and Sonmez, GT. Effects of Low- Versus High-Load Resistance Training on Muscle Strength and Hypertrophy in Well-Trained Men. *J. Strength Cond Res.* 29: 2954-2963, 2015.
 75. Seaborne, RA, Strauss, J, Cocks, M, Shepherd, S, O'Brien, TD, van Someren, KA, Bell, PG, Murgatroyd, C, Morton, JP, Stewart, CE, and Sharples, AP. Human skeletal muscle possesses an epigenetic memory of hypertrophy. *Scientific Reports* 8: 1898, 2018.
 76. Seaborne, RA, Hughes, DC, Turner, DC, Owens, DJ, Baehr, LM, Gorski, P, Semenova, EA, Borisov, OV, Larin, AK, Popov, DV, Generozov, EV, Sutherland, H, Ahmetov, II, Jarvis, JC, Bodine, SC, and Sharples, AP. UBR5 is a novel E3 ubiquitin ligase involved in skeletal muscle hypertrophy and recovery from atrophy. *J. Physiol.* 597: 3727-3749, 2019.
 77. Sharples, AP, Stewart, CE, and Seaborne, RA. Does skeletal muscle have an 'epi'-memory? The role of epigenetics in nutritional programming, metabolic disease, aging and exercise. *Aging Cell.* 15: 603-616, 2016.
 78. Simoneau, JA, and Bouchard, C. Genetic determinism of fiber type proportion in human skeletal muscle. *FASEB J.* 9: 1091-1095, 1995.
 79. Singh, MA, Ding, W, Manfredi, TJ, Solares, GS, O'Neill, EE, Clements, KM, Ryan, ND, Kehayias, JJ, Fielding, RA, and Evans, WJ. Insulin-like growth factor I in skeletal muscle after weight-lifting exercise in frail elders. *Am. J. Physiol.* 277: E135-43, 1999.
 80. Smith, GI, Atherton, P, Villareal, DT, Frimel, TN, Rankin, D, Rennie, MJ, and Mittendorfer, B. Differences in muscle protein synthesis and anabolic signaling in the postabsorptive state and in response to food in 65-80 year old men and women. *PLoS One* 3: e1875, 2008.
 81. Smith, GI, Villareal, DT, Sinacore, DR, Shah, K, and Mittendorfer, B. Muscle protein synthesis response to exercise training in obese, older men and women. *Med. Sci. Sports Exerc.* 44: 1259-1266, 2012.
 82. Smith, GI, and Mittendorfer, B. Sexual dimorphism in skeletal muscle protein turnover. *J. Appl. Physiol.* (1985) 120: 674-682, 2016.
 83. Stewart, CE, and Rittweger, J. Adaptive processes in skeletal muscle: molecular regulators and genetic influences. *J. Musculoskelet. Neuronal Interact.* 6: 73-86, 2006.
 84. Stewart, VH, Saunders, DH, and Greig, CA. Responsiveness of muscle size and strength to physical training in very elderly people: a systematic review. *Scand. J. Med. Sci. Sports* 24: e1-10, 2014.
 85. Stragier, S, Baudry, S, Poortmans, J, Duchateau, J, and Carpentier, A. Leucine-enriched protein supplementation does not influence neuromuscular adaptations in response to a 6-month strength training programme in older adults. *Exp. Gerontol.* 82: 58-66, 2016.
 86. Tang, JE, Perco, JG, Moore, DR, Wilkinson, SB, and Phillips, SM. Resistance training alters the response of fed state mixed muscle protein synthesis in young men. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 294: R172-8, 2008.

87. Urban, RJ, Bodenbun, YH, Gilkison, C, Foxworth, J, Coggan, AR, Wolfe, RR, and Ferrando, A. Testosterone administration to elderly men increases skeletal muscle strength and protein synthesis. *Am. J. Physiol.* 269: E820-6, 1995.
88. Van Etten, LM, Verstappen, FT, and Westerterp, KR. Effect of body build on weight-training-induced adaptations in body composition and muscular strength. *Med. Sci. Sports Exerc.* 26: 515-521, 1994.
89. Verdijk, LB, Koopman, R, Schaart, G, Meijer, K, Savelberg, HH, and van Loon, LJ. Satellite cell content is specifically reduced in type II skeletal muscle fibers in the elderly. *Am. J. Physiol. Endocrinol. Metab.* 292: E151-7, 2007.
90. Watanabe, Y, Tanimoto, M, Ohgane, A, Sanada, K, Miyachi, M, and Ishii, N. Increased muscle size and strength from slow-movement, low-intensity resistance exercise and tonic force generation. *J. Aging Phys. Act.* 21: 71-84, 2013.
91. Waters, DL, Baumgartner, RN, Garry, PJ, and Vellas, B. Advantages of dietary, exercise-related, and therapeutic interventions to prevent and treat sarcopenia in adult patients: an update. *Clin. Interv. Aging* 5: 259-270, 2010.
92. Welle, S, Thornton, C, and Statt, M. Myofibrillar protein synthesis in young and old human subjects after three months of resistance training. *Am. J. Physiol.* 268: E422-7, 1995.
93. Welle, S, Totterman, S, and Thornton, C. Effect of age on muscle hypertrophy induced by resistance training. *J. Gerontol. A Biol. Sci. Med. Sci.* 51: M270-5, 1996.
94. Wilkinson, SB, Phillips, SM, Atherton, PJ, Patel, R, Yarasheski, KE, Tarnopolsky, MA, and Rennie, MJ. Differential effects of resistance and endurance exercise in the fed state on signalling molecule phosphorylation and protein synthesis in human muscle. *J. Physiol.* 586: 3701-3717, 2008.
95. Xu, M, Chen, X, Chen, D, Yu, B, Li, M, He, J, and Huang, Z. Regulation of skeletal myogenesis by microRNAs. *J. Cell. Physiol.* , 2019.
96. Yarasheski, KE. Managing sarcopenia with progressive resistance exercise training. *J. Nutr. Health Aging* 6: 349-356, 2002.
97. Zacker, RJ. Health-related implications and management of sarcopenia. *JAAAPA* 19: 24-29, 2006.
98. Zhao, W, Pan, J, Zhao, Z, Wu, Y, Bauman, WA, and Cardozo, CP. Testosterone protects against dexamethasone-induced muscle atrophy, protein degradation and MAFbx upregulation. *J. Steroid Biochem. Mol. Biol.* 110: 125-129, 2008.
5. Andersen, LL, Magnusson, SP, Nielsen, M, Haleem, J, Poulsen, K, and Aagaard, P. Neuromuscular activation in conventional therapeutic exercises and heavy resistance exercises: implications for rehabilitation. *Phys. Ther.* 86: 683-697, 2006.
6. Andersen, V, Fimland, MS, Wiik, E, Skoglund, A, and Sæterbakken, AH. Effects of grip width on muscle strength and activation in the lat pull-down. *J. Strength Cond Res.* 28: 1135-1142, 2014.
7. Antonio, J. Nonuniform response of skeletal muscle to heavy resistance training: can bodybuilders induce regional muscle hypertrophy. *Journal of Strength and Conditioning Research* 14: 102-113, 2000.
8. Baechle, TR, Earle, RW. Essentials of strength training and conditioning. In: Anonymous Champaign, IL: Human Kinetics, 2008.
9. Baker, D, Wilson, G, and Carolyn, R. Periodization: the effect on strength of manipulating volume and intensity. *J Strength Cond Res* 8: 235-242, 1994.
10. Balsalobre, C, Santos-Concejero, J, Baz, E, and Schoenfeld, BJ. The effects of exercise variation in muscle thickness, maximal strength and motivation in resistance trained men. *Plos One*, 2019.
11. Barbalho, M, Coswig, VS, Raiol, R, Steele, J, Fisher, J, Paoli, A, and Gentil, P. Effects of Adding Single Joint Exercises to a Resistance Training Programme in Trained Women. *Sports (Basel)* 6: 10.3390/sports6040160, 2018.
12. Barbalho, M, Coswig, VS, Raiol, R, Steele, J, Fisher, JP, Paoli, A, Bianco, A, and Gentil, P. Does the addition of single joint exercises to a resistance training program improve changes in performance and anthropometric measures in untrained men? *Eur. J. Transl. Myol* 28: 7827, 2018.
13. Barbalho, M, Gentil, P, Raiol, R, Fisher, J, Steele, J, and Coswig, V. Influence of Adding Single-Joint Exercise to a Multijoint Resistance Training Program in Untrained Young Women. *J. Strength Cond Res.*, 2018.
14. Barbalho, M, Coswig, V, Raiol, R, Fisher, J, Steele, J, Bianco, A, and Gentil, P. Single joint exercises do not provide benefits in performance and anthropometric changes in recreational bodybuilders. *Eur. J. Sport. Sci.* : 1-8, 2019.
15. Barnett, C, Kippers, V, and Turner, P. Effects of variations of the bench press exercise on the EMG activity of five shoulder muscles. *J Strength Cond Res* 9: 222-227, 1995.
16. Bjornsen, T, Wernbom, M, Lovstad, A, Paulsen, G, D'Souza, RF, Cameron-Smith, D, Flesche, A, Hisdal, J, Berntsen, S, and Raastad, T. Delayed myonuclear addition, myofiber hypertrophy, and increases in strength with high-frequency low-load blood flow restricted training to volitional failure. *J. Appl. Physiol.* (1985) 126: 578-592, 2019.
17. Bloomquist, K, Langberg, H, Karlsen, S, Madsgaard, S, Boesen, M, and Raastad, T. Effect of range of motion in heavy load squatting on muscle and tendon adaptations. *Eur. J. Appl. Physiol.* 113: 2133-2142, 2013.
18. Bompa, T, and Haff, GG. Theory and Methodology of Training. Champaign, IL; Human Kinetics, 2009.
19. Botton, CE, Wilhelm, EN, Ughini, CC, Pinto, RS, and Lima, CS. Electromyographical analysis of the deltoid between different strength training exercises. *Medicina Sportiva* 17: 67-71, 2013.
20. Bressel, E, Willardson, JM, Thompson, B, and Fontana, FE. Effect of instruction, surface stability, and load intensity on trunk muscle activity. *J. Electromyogr. Kinesiol.* 19: e500-4, 2009.

Chapter 8

1. Ahmadzad, S, Ghorbani, S, Ghasemikaram, M, and Bahmanzadeh, M. Effects of short-term nonperiodized, linear periodized and daily undulating periodized resistance training on plasma adiponectin, leptin and insulin resistance. *Clin. Biochem.* 47: 417-422, 2014.
2. Aisbett, B, Condo, D, Zacharewicz, E, and Lamon, S. The Impact of Shiftwork on Skeletal Muscle Health. *Nutrients* 9: 10.3390/nu9030248, 2017.
3. Alves Souza, RW, Aguiar, AF, Vechetti-Junior, JJ, Piedade, WP, Rocha Campos, GE, and Dal-Pai-Silva, M. Resistance training with excessive training load and insufficient recovery alters skeletal muscle mass-related protein expression. *J. Strength Cond Res.* 28: 2338-2345, 2014.
4. Amstrup, AK, Sikjaer, T, Pedersen, SB, Heickendorff, L, Mosekilde, L, and Rejnmark, L. Reduced fat mass and increased lean mass in response to 1 year of melatonin treatment in postmenopausal women: A randomized placebo-controlled trial. *Clin. Endocrinol. (Oxf)* 84: 342-347, 2016.

21. Buchmann, N, Spira, D, Norman, K, Demuth, I, Eckardt, R, and Steinhagen-Thiessen, E. Sleep, Muscle Mass and Muscle Function in Older People. *Dtsch. Arztebl Int.* 113: 253-260, 2016.
22. Buford, TW, Rossi, SJ, Smith, DB, and Warren, AJ. A comparison of periodization models during nine weeks with equated volume and intensity for strength. *J. Strength Cond Res.* 21: 1245-1250, 2007.
23. Burley, SD, Whittingham-Dowd, J, Allen, J, Grosset, JF, and Onambele-Pearson, GL. The Differential Hormonal Milieu of Morning versus Evening May Have an Impact on Muscle Hypertrophic Potential. *PLoS One* 11: e0161500, 2016.
24. Burnham, TR, Ruud, JD, and McGowan, R. Bench press training program with attached chains for female volleyball and basketball athletes. *Percept. Mot. Skills* 110: 61-68, 2010.
25. Cadegiani, FA, and Kater, CE. Body composition, metabolism, sleep, psychological and eating patterns of overtraining syndrome: Results of the EROS study (EROS-PROFILE). *J. Sports Sci.* 36: 1902-1910, 2018.
26. Cambridge, ED, Sidorkewicz, N, Ikeda, DM, and McGill, SM. Progressive hip rehabilitation: the effects of resistance band placement on gluteal activation during two common exercises. *Clin. Biomech. (Bristol, Avon)* 27: 719-724, 2012.
27. Campos, YD, and Silva, SF. Comparison of electromyographic activity during the bench press and barbell pullover exercises. *Motriz: Revista de Educação Física* 20: 200-205, 2014.
28. Cibulka, M, Wenthe, A, Boyle, Z, Callier, D, Schwerdt, A, Jarman, D, and Strube, MJ. Variation in Medial and Lateral Gastrocnemius Muscle Activity with Foot Position. *Int. J. Sports Phys. Ther.* 12: 233-241, 2017.
29. Clark, C. Assessment of 4 weeks of P90x (r) training on muscular strength and endurance, anaerobic power, and body composition. Middle Tennessee State University, JEWL Scholar@MTSU Repository, 2014.
30. Clark, KM, Holt, LE, and Sinyard, J. Electromyographic comparison of the upper and lower rectus abdominis during abdominal exercises. *J. Strength Cond Res.* 17: 475-483, 2003.
31. Conlon, JA, Newton, RU, Tufano, JJ, Banyard, HG, Hopper, AJ, Ridge, AJ, and Haff, GG. Periodization Strategies in Older Adults: Impact on Physical Function and Health. *Med. Sci. Sports Exerc.* 48: 2426-2436, 2016.
32. Contreras, B, and Schoenfeld, B. To crunch or not to crunch: An evidence-based examination of spinal flexion exercises, their potential risks, and their applicability to program design. *Strength & Conditioning Journal* 33: 8-18, 2011.
33. Contreras, B, Cronin, J, Schoenfeld, BJ, Nates, R, and Sonmez, GT. Are all hip extension exercises created equal? *Strength Cond J* 35: 17-22, 2013.
34. Contreras, B, Vigotsky, AD, Schoenfeld, BJ, Beardsley, C, and Cronin, J. A Comparison of Gluteus Maximus, Biceps Femoris, and Vastus Lateralis EMG Activity in the Back Squat and Barbell Hip Thrust Exercises. *J. Appl. Biomech.*, 2015.
35. Critchley, D. Instructing pelvic floor contraction facilitates transversus abdominis thickness increase during low-abdominal hollowing. *Physiother. Res. Int.* 7: 65-75, 2002.
36. Cunanan, AJ, DeWeese, BH, Wagle, JP, Carroll, KM, Sausaman, R, Hornsby, WG, 3rd, Haff, GG, Triplett, NT, Pierce, KC, and Stone, MH. The General Adaptation Syndrome: A Foundation for the Concept of Periodization. *Sports Med.* 48: 787-797, 2018.
37. de Franca, HS, Branco, PA, Guedes Junior, DP, Gentil, P, Steele, J, and Teixeira, CV. The effects of adding single-joint exercises to a multi-joint exercise resistance training program on upper body muscle strength and size in trained men. *Appl. Physiol. Nutr. Metab.* 40: 822-826, 2015.
38. de Lima, C, Boullousa, DA, Frollini, AB, Donatto, FF, Leite, RD, Gonelli, PR, Montebello, MI, Prestes, J, and Cesar, MC. Linear and daily undulating resistance training periodizations have differential beneficial effects in young sedentary women. *Int. J. Sports Med.* 33: 723-727, 2012.
39. De Souza, EO, Tricoli, V, Rauch, J, Alvarez, MR, Laurentino, G, Aihara, AY, Cardoso, FN, Roschel, H, and Ugrinowitsch, C. Different Patterns in Muscular Strength and Hypertrophy Adaptations in Untrained Individuals Undergoing Nonperiodized and Periodized Strength Regimens. *J. Strength Cond Res.* 32: 1238-1244, 2018.
40. Dos Santos, L, Ribeiro, AS, Cavalcante, EF, Nabuco, HC, Antunes, M, Schoenfeld, BJ, and Cyrino, ES. Effects of Modified Pyramid System on Muscular Strength and Hypertrophy in Older Women. *Int. J. Sports Med.* 39: 613-618, 2018.
41. Duncan, M. Muscle activity of the upper and lower rectus abdominis during exercises performed on and off a Swiss ball. *J. Bodyw Mov. Ther.* 13: 364-367, 2009.
42. Ebben, WP, Feldmann, CR, Dayne, A, Mitsche, D, Alexander, P, and Knetzger, KJ. Muscle activation during lower body resistance training. *Int. J. Sports Med.* 30: 1-8, 2009.
43. Ema, R, Wakahara, T, Miyamoto, N, Kanehisa, H, and Kawakami, Y. Inhomogeneous architectural changes of the quadriceps femoris induced by resistance training. *Eur. J. Appl. Physiol.* 113: 2691-2703, 2013.
44. Ema, R, Sakaguchi, M, Akagi, R, and Kawakami, Y. Unique activation of the quadriceps femoris during single- and multi-joint exercises. *Eur. J. Appl. Physiol.* 116: 1031-1041, 2016.
45. Escamilla, RE, Fleisig, GS, Zheng, N, Barrentine, SW, Wilk, KE, and Andrews, JR. Biomechanics of the knee during closed kinetic chain and open kinetic chain exercises. *Med. Sci. Sports Exerc.* 30: 556-569, 1998.
46. Escamilla, RE, Babb, E, DeWitt, R, Jew, P, Kelleher, P, Burnham, T, Busch, J, D'Anna, K, Mowbray, R, and Imamura, RT. Electromyographic analysis of traditional and nontraditional abdominal exercises: implications for rehabilitation and training. *Phys. Ther.* 86: 656-671, 2006.
47. Fink, J, Kikuchi, N, Yoshida, S, Terada, K, and Nakazato, K. Impact of high versus low fixed loads and non-linear training loads on muscle hypertrophy, strength and force development. *Springerplus* 5: 698-016-2333-z. eCollection 2016, 2016.
48. Fonseca, RM, Roschel, H, Tricoli, V, de Souza, EO, Wilson, JM, Laurentino, GC, Aihara, AY, de Souza Leao, AR, and Ugrinowitsch, C. Changes in exercises are more effective than in loading schemes to improve muscle strength. *J. Strength Cond Res.*, 2014.
49. Fry, AC, and Kraemer, WJ. Resistance exercise overtraining and overreaching. Neuroendocrine responses. *Sports Med.* 23: 106-129, 1997.
50. Fullagar, HH, Skorski, S, Duffield, R, Hammes, D, Coutts, AJ, and Meyer, T. Sleep and athletic performance: the effects of sleep loss on exercise performance, and physiological and cognitive responses to exercise. *Sports Med.* 45: 161-186, 2015.
51. Garcia-Lopez, D, Hernandez-Sanchez, S, Martin, E, Marin, PJ, Zarzosa, F, and Herrero, AJ. Free-weight augmentation with elastic bands improves bench-press kinematics in professional rugby players. *J. Strength Cond Res.*, 2014.
52. Gentil, P, Soares, SR, Pereira, MC, Cunha, RR, Martorelli, SS, Martorelli, AS, and Bottaro, M. Effect of adding single-joint exercises to a multi-joint exercise resistance-training program on strength and hypertrophy in untrained subjects. *Appl. Physiol. Nutr. Metab.* 38: 341-344, 2013.

53. Gentil, P, Soares, S, and Bottaro, M. Single vs. Multi-Joint Resistance Exercises: Effects on Muscle Strength and Hypertrophy. *Asian J. Sports Med.* 6: e24057, 2015.
54. Gentil, P, Fisher, J, and Steele, J. A Review of the Acute Effects and Long-Term Adaptations of Single- and Multi-Joint Exercises during Resistance Training. *Sports Med.* 47: 843-855, 2017.
55. Glass, SC, and Armstrong, T. Electromyographical activity of the pectoralis muscle during incline and decline bench presses. *J Strength Cond Res* 11: 163-167, 1997.
56. Gray, H. *Gray's Anatomy: The Anatomical Basis of Medicine and Surgery*. London; Pearson Professional Ltd, 1995.
57. Grgic, J, Mikulic, P, Podnar, H, and Pedisic, Z. Effects of linear and daily undulating periodized resistance training programs on measures of muscle hypertrophy: a systematic review and meta-analysis. *PeerJ* 5: e3695, 2017.
58. Grgic, J, Lazinica, B, Garofolini, A, Schoenfeld, BJ, Saner, NJ, and Mikulic, P. The effects of time of day-specific resistance training on adaptations in skeletal muscle hypertrophy and muscle strength: A systematic review and meta-analysis. *Chronobiol. Int.* 36: 449-460, 2019.
59. Habermeyer, P, Kaiser, E, Knappe, M, Kreusser, T, and Wiedemann, E. Functional anatomy and biomechanics of the long biceps tendon. *Unfallchirurg* 90: 319-329, 1987.
60. Hackett, DA, Amirthalingam, T, Mitchell, L, Mavros, Y, Wilson, GC, and Halaki, M. Effects of a 12-Week Modified German Volume Training Program on Muscle Strength and Hypertrophy-A Pilot Study. *Sports (Basel)* 6: 10.3390/sports6010007, 2018.
61. Harries, SK, Lubans, DR, and Callister, R. Comparison of resistance training progression models on maximal strength in sub-elite adolescent rugby union players. *J. Sci. Med. Sport*, 2015.
62. Hebert-Losier, K, Schneiders, AG, Garcia, JA, Sullivan, SJ, and Simoneau, GG. Influence of knee flexion angle and age on triceps surae muscle activity during heel raises. *J. Strength Cond Res.* 26: 3124-3133, 2012.
63. Helms, E, Fitschen, PJ, Aragon, A, Cronin, J, and Schoenfeld, BJ. Recommendations for natural bodybuilding contest preparation: resistance and cardiovascular training. *J. Sports Med. Phys. Fitness*, 2014.
64. Hung, YJ, and Gross, MT. Effect of foot position on electromyographic activity of the vastus medialis oblique and vastus lateralis during lower-extremity weight-bearing activities. *J. Orthop. Sports Phys. Ther.* 29: 93-102; discussion 103-5, 1999.
65. Hunter, GR, Wetzstein, CJ, McLafferty, CL, Jr, Zuckerman, PA, Landers, KA, and Bamman, MM. High-resistance versus variable-resistance training in older adults. *Med. Sci. Sports Exerc.* 33: 1759-1764, 2001.
66. Israetel, M, and Hoffmann, J. *How Much should I Train? an Introduction to Training Volume Landmarks*. Philadelphia, PA; Renaissance Periodization., 2017.
67. Junior, V, Gentil, P, Oliveira, E, and Carmo, J. Comparison among the EMG activity of the pectoralis major, anterior deltoidis and triceps brachii during the bench press and peck deck exercises. *Rev Bras Med Esporte* 13: 43-46, 2007.
68. Karst, GM, and Willett, GM. Effects of specific exercise instructions on abdominal muscle activity during trunk curl exercises. *J. Orthop. Sports Phys. Ther.* 34: 4-12, 2004.
69. Kiely, J. Periodization Theory: Confronting an Inconvenient Truth. *Sports Med.* 48: 753-764, 2018.
70. Kim, K, Shin, D, Jung, GU, Lee, D, and Park, SM. Association between sleep duration, fat mass, lean mass and obesity in Korean adults: the fourth and fifth Korea National Health and Nutrition Examination Surveys. *J. Sleep Res.* 26: 453-460, 2017.
71. Kim, M, Sasai, H, Kojima, N, and Kim, H. Objectively measured night-to-night sleep variations are associated with body composition in very elderly women. *J. Sleep Res.* 24: 639-647, 2015.
72. Kok, LY, Hamer, PW, and Bishop, DJ. Enhancing muscular qualities in untrained women: linear versus undulating periodization. *Med. Sci. Sports Exerc.* 41: 1797-1807, 2009.
73. Kraemer, WJ, Häkkinen, K, Triplett-Mcbride, NT, Fry, AC, Koziris, LP, Ratamess, NA, Bauer, JE, Volek, JS, McConnell, T, Newton, RU, Gordon, SE, Cummings, D, Hauth, J, Pullo, F, Lynch, JM, Fleck, SJ, Mazzetti, SA, and Knuttgen, HG. Physiological changes with periodized resistance training in women tennis players. *Med. Sci. Sports Exerc.* 35: 157-168, 2003.
74. Kramer, JB, Stone, MH, O'Bryant, HS, Conley, MS, Johnson, RL, Nieman, DC, Honeycutt, DR, and Hoke, TP. Effects of single vs. multiple sets of weight training: impact of volume, intensity, and variation. *Journal of strength and Conditioning Research* 11: 143-147, 1997.
75. Kubo, K, Ikebukuro, T, and Yata, H. Effects of squat training with different depths on lower limb muscle volumes. *Eur. J. Appl. Physiol.*, 2019.
76. Lauver, JD, Cayot, TE, and Scheuermann, BW. Influence of bench angle on upper extremity muscular activation during bench press exercise. *Eur. J. Sport. Sci.* 16: 309-316, 2016.
77. Le Bozec, S, Maton, B, and Cnockaert, JC. The synergy of elbow extensor muscles during dynamic work in man. I. Elbow extension. *Eur. J. Appl. Physiol. Occup. Physiol.* 44: 255-269, 1980.
78. Lehman, GJ, and McGill, SM. Quantification of the differences in electromyographic activity magnitude between the upper and lower portions of the rectus abdominis muscle during selected trunk exercises. *Phys. Ther.* 81: 1096-1101, 2001.
79. Lehman, GJ, Buchan, DD, Lundy, A, Myers, N, and Nalborczyk, A. Variations in muscle activation levels during traditional latissimus dorsi weight training exercises: An experimental study. *Dyn. Med.* 3: 4, 2004.
80. Levy, AS, Kelly, BT, Lintner, SA, Osbahr, DC, and Speer, KP. Function of the long head of the biceps at the shoulder: electromyographic analysis. *J. Shoulder Elbow Surg.* 10: 250-255, 2001.
81. Lewis, CL, and Sahrmann, SA. Muscle activation and movement patterns during prone hip extension exercise in women. *J. Athl Train.* 44: 238-248, 2009.
82. Lusk, SJ, Hale, BD, and Russell, DM. Grip width and forearm orientation effects on muscle activity during the lat pull-down. *J. Strength Cond Res.* 24: 1895-1900, 2010.
83. Lynn, SK, and Costigan, PA. Changes in the medial-lateral hamstring activation ratio with foot rotation during lower limb exercise. *J. Electromyogr. Kinesiol.* 19: e197-205, 2009.
84. MacIntosh, BR. Recent developments in understanding the length dependence of contractile response of skeletal muscle. *Eur. J. Appl. Physiol.* 117: 1059-1071, 2017.
85. Mannarino, P, Matta, T, Lima, J, Simao, R, and Freitas de Salles, B. Single-Joint Exercise Results in Higher Hypertrophy of Elbow Flexors Than Multijoint Exercise. *J. Strength Cond Res.*, 2019.
86. Marchant, DC, Greig, M, and Scott, C. Attentional focusing instructions influence force production and muscular activity during isokinetic elbow flexions. *J. Strength Cond Res.* 23: 2358-2366, 2009.

87. Marchetti, PH, and Uchida, MC. Effects of the pullover exercise on the pectoralis major and latissimus dorsi muscles as evaluated by EMG. *J. Appl. Biomech.* 27: 380-384, 2011.
88. Marcori, AJ, Moura, TBMA, and Okazaki, VHA. Gastrocnemius muscle activation during plantar flexion with different feet positioning in physically active young men. *Isokinetics and Exercise Science* 25: 121-125, 2017.
89. Marx, JO, Ratamess, NA, Nindl, BC, Gotshalk, LA, Volek, JS, Dohi, K, Bush, JA, Gomez, AL, Mazzetti, SA, Fleck, SJ, Häkkinen, K, Newton, RU, and Kraemer, WJ. Low-volume circuit versus high-volume periodized resistance training in women. *Med. Sci. Sports Exerc.* 33: 635-643, 2001.
90. McAllister, MJ, Schilling, BK, Hammond, KG, Weiss, LW, and Farney, TM. Effect of grip width on electromyographic activity during the upright row. *J. Strength Cond Res.* 27: 181-187, 2013.
91. McCaw, ST, and Melrose, DR. Stance width and bar load effects on leg muscle activity during the parallel squat. *Med. Sci. Sports Exerc.* 31: 428-436, 1999.
92. McGill, S. Core training: Evidence translating to better performance and injury prevention. *Strength Cond J* 32: 33-46, 2010.
93. Monico-Neto, M, Antunes, HK, Lee, KS, Phillips, SM, Giampa, SQ, de Sá Souza, H, Dattilo, M, Medeiros, A, de Moraes, WM, Tufik, S, and de Mello, MT. Resistance training minimizes catabolic effects induced by sleep deprivation in rats. *Appl. Physiol. Nutr. Metab.* 40: 1143-1150, 2015.
94. Monteiro, AG, Aoki, MS, Evangelista, AL, Alveno, DA, Monteiro, GA, Picarro, Ida, C, and Ugrinowitsch, C. Nonlinear periodization maximizes strength gains in split resistance training routines. *J. Strength Cond Res.* 23: 1321-1326, 2009.
95. Ninos, JC, Irrgang, JJ, Burdett, R, and Weiss, JR. Electromyographic analysis of the squat performed in self-selected lower extremity neutral rotation and 30 degrees of lower extremity turn-out from the self-selected neutral position. *J. Orthop. Sports Phys. Ther.* 25: 307-315, 1997.
96. O'Bryant, HS, Byrd, R, and Stone, MH. Cycle ergometer performance and maximum leg and hip strength adaptations to two different methods of weight training. *J Appl Sport Sci Res* 2: 27-30, 1988.
97. Ogasawara, R, Yasuda, T, Sakamaki, M, Ozaki, H, and Abe, T. Effects of periodic and continued resistance training on muscle CSA and strength in previously untrained men. *Clin. Physiol. Funct. Imaging* 31: 399-404, 2011.
98. Ogasawara, R, Kobayashi, K, Tsutaki, A, Lee, K, Abe, T, Fujita, S, Nakazato, K, and Ishii, N. mTOR signaling response to resistance exercise is altered by chronic resistance training and detraining in skeletal muscle. *J. Appl. Physiol.* , 2013.
99. Ogasawara, R, Yasuda, T, Ishii, N, and Abe, T. Comparison of muscle hypertrophy following 6-month of continuous and periodic strength training. *Eur. J. Appl. Physiol.* 113: 975-985, 2013.
100. Paoli, A, Marcolin, G, and Petrone, N. The effect of stance width on the electromyographical activity of eight superficial thigh muscles during back squat with different bar loads. *J. Strength Cond Res.* 23: 246-250, 2009.
101. Patel, SR. Reduced sleep as an obesity risk factor. *Obes. Rev.* 10 Suppl 2: 61-68, 2009.
102. Pelzer, T, Ullrich, B, and Pfeiffer, M. Periodization effects during short-term resistance training with equated exercise variables in females. *Eur. J. Appl. Physiol.* 117: 441-454, 2017.
103. Poliquin, C. Five Steps to Increasing the Effectiveness of Your Strength Training Program. . *Natl Strength Cond Assoc. J.* 10: 34-39, 1988.
104. Prestes, J, De Lima, C, Frollini, AB, Donatto, FF, and Conte, M. Comparison of linear and reverse linear periodization effects on maximal strength and body composition. *J. Strength Cond Res.* 23: 266-274, 2009.
105. Prestes, J, Frollini, AB, de Lima, C, Donatto, FF, Foschini, D, de Cassia Marqueti, R, Figueira, A, Jr, and Fleck, SJ. Comparison between linear and daily undulating periodized resistance training to increase strength. *J. Strength Cond Res.* 23: 2437-2442, 2009.
106. Rauch, JT, Ugrinowitsch, C, Barakat, CI, Alvarez, MR, Brummert, DL, Aube, DW, Barsuhn, AS, Hayes, D, Tricoli, V, and De Souza, EO. Auto-regulated exercise selection training regimen produces small increases in lean body mass and maximal strength adaptations in strength-trained individuals. *J. Strength Cond Res.* , 2017.
107. Riemann, BL, Limbaugh, GK, Eitner, JD, and LeFavi, RG. Medial and lateral gastrocnemius activation differences during heel-raise exercise with three different foot positions. *J. Strength Cond Res.* 25: 634-639, 2011.
108. Rocha Júnior, VA, Gentil, P, Oliveira, E, and do Carmo, J. Comparison among the EMG activity of the pectoralis major, anterior deltoidis and triceps brachii during the bench press and peck deck exercises. *Rev Bras Med Esporte* 13: 43e-46e, 2007.
109. Samuels, C. Sleep, recovery, and performance: the new frontier in high-performance athletics. *Neurol. Clin.* 26: 169-80; ix-x, 2008.
110. Sanchis-Moysi, J, Idoate, F, Dorado, C, Alayon, S, and Calbet, JA. Large asymmetric hypertrophy of rectus abdominis muscle in professional tennis players. *PLoS One* 5: e15858, 2010.
111. Sarti, MA, Monfort, M, Fuster, MA, and Villaplana, LA. Muscle activity in upper and lower rectus abdominus during abdominal exercises. *Arch. Phys. Med. Rehabil.* 77: 1293-1297, 1996.
112. Schiotz, MK, Potteiger, JA, Huntsinger, PG, and Denmark, LCDC. The short-term effects of periodized and constant-intensity training on body composition, strength, and performance. *Journal of Strength & Conditioning Research* 12: 173-178, 1998.
113. Schoenfeld, B, Kolber, MJ, and Haimes, JE. The Upright Row: Implications for Preventing Subacromial Impingement. . *Strength and Conditioning Journal* 33: 25-28, 2011.
114. Schoenfeld, BJ, Contreras, B, Ogborn, D, Galpin, A, Krieger, J, and Sonmez, GT. Effects of varied versus constant loading zones on muscular adaptations in well-trained men. *Int J Sports Med.* DOI: 10.1055/s-0035-1569369, 2016.
115. Schoenfeld, B, Sonmez, RG, Kolber, MJ, Contreras, B, Harris, R, and Ozen, S. Effect of hand position on EMG activity of the posterior shoulder musculature during a horizontal abduction exercise. *J. Strength Cond Res.* 27: 2644-2649, 2013.
116. Schoenfeld, BJ. The mechanisms of muscle hypertrophy and their application to resistance training. *J. Strength Cond Res.* 24: 2857-2872, 2010.
117. Schoenfeld, BJ, Contreras, B, Tiryaki-Sonmez, G, Willardson, JM, and Fontana, F. An electromyographic comparison of a modified version of the plank with a long lever and posterior tilt versus the traditional plank exercise. *Sports Biomech.* 13: 296-306, 2014.
118. Schoenfeld, BJ, Contreras, B, Tiryaki-Sonmez, G, Wilson, JM, Kolber, MJ, and Peterson, MD. Regional differences in muscle activation during hamstring exercise. *J. Strength Cond Res.* 29: 159-164, 2015.
119. Schoenfeld, BJ, Ogborn, D, and Krieger, JW. Dose-response relationship between weekly resistance training volume and

- increases in muscle mass: A systematic review and meta-analysis. *J. Sports Sci.* 35: 1073-1082, 2017.
120. Schoenfeld, BJ, Vigotsky, A, Contreras, B, Golden, S, Alto, A, Larson, R, Winkelman, N, and Paoli, A. Differential effects of attentional focus strategies during long-term resistance training. *Eur. J. Sport. Sci.* 18: 705-712, 2018.
 121. Schoenfeld, BJ, Grgic, J, Haun, C, Itagaki, T, and Helms, ER. Calculating Set-Volume for the Limb Muscles with the Performance of Multi-Joint Exercises: Implications for Resistance Training Prescription. *Sports (Basel)* 7: 10.3390/sports7070177, 2019.
 122. Sedliak, M, Finni, T, Cheng, S, Kraemer, WJ, and Häkkinen, K. Effect of time-of-day-specific strength training on serum hormone concentrations and isometric strength in men. *Chronobiol. Int.* 24: 1159-1177, 2007.
 123. Selkowitz, DM, Beneck, GJ, and Powers, CM. Comparison of Electromyographic Activity of the Superior and Inferior Portions of the Gluteus Maximus Muscle During Common Therapeutic Exercises. *J. Orthop. Sports Phys. Ther.* 46: 794-799, 2016.
 124. SELYE, H. Stress and the general adaptation syndrome. *Br. Med. J.* 1: 1383-1392, 1950.
 125. Signorile, JE, Lew, KM, Stoutenberg, M, Pluchino, A, Lewis, JE, and Gao, J. Range of motion and leg rotation affect electromyography activation levels of the superficial quadriceps muscles during leg extension. *J. Strength Cond Res.* 28: 2536-2545, 2014.
 126. Simao, R, Spinetti, J, de Salles, BF, Matta, T, Fernandes, L, Fleck, SJ, Rhea, MR, and Strom-Olsen, HE. Comparison between nonlinear and linear periodized resistance training: hypertrophic and strength effects. *J. Strength Cond Res.* 26: 1389-1395, 2012.
 127. Snyder, BJ, and Leech, JR. Voluntary increase in latissimus dorsi muscle activity during the lat pull-down following expert instruction. *J. Strength Cond Res.* 23: 2204-2209, 2009.
 128. Snyder, BJ, and Fry, WR. Effect of verbal instruction on muscle activity during the bench press exercise. *J. Strength Cond Res.* 26: 2394-2400, 2012.
 129. Souza, EO, Ugrinowitsch, C, Tricoli, V, Roschel, H, Lowery, RP, Aihara, AY, Leao, ARS, and Wilson, JM. Early Adaptations to Six Weeks of Non-Periodized and Periodized Strength Training Regimens in Recreational Males. *J Sports Sci Med.* 13: 604-609, 2014.
 130. Stasinaki, AN, Zaras, N, Methenitis, S, Tsitkanou, S, Krase, A, Kavvoura, A, and Terzis, G. Triceps brachii muscle strength and architectural adaptations with resistance training exercises at short or long fascicle length. *Journal of Functional Morphology and Kinesiology.* 3: 28, 2018.
 131. Stone, MH, O'Bryant, HS, Schilling, BK, and R.L. Johnson, RL. Periodization: Effects Of Manipulating Volume And Intensity. Part 1. *Strength Cond J* 21: 56-62, 1999.
 132. Stone, MH, Potteiger, JA, Pierce, KC, Proulx, CM, O'Bryant, HS, Johnson, RL, and Stone, ME. Comparison of the effects of three different weight-training programs on the one repetition maximum squat. *J Strength Cond Res* 14: 332-337, 2000.
 133. Stone, MH, O'Bryant, H, and Garhammer, J. A hypothetical model for strength training. *J. Sports Med. Phys. Fitness* 21: 342-351, 1981.
 134. Suhara, H, Ae, K, Nariai, M, Shiraki, H, and Miyakawa, S. A Three-Dimensional Biomechanical Analysis of KOSHIWARI Exercise. *Journal of Sports Science* 6: 149-158, 2018.
 135. Tan, X, Titova, OE, Lindberg, E, Elmstahl, S, Lind, L, Schioth, HB, and Benedict, C. Association Between Self-Reported Sleep Duration and Body Composition in Middle-Aged and Older Adults. *J. Clin. Sleep Med.* 15: 431-435, 2019.
 136. Trebs, AA, Brandenburg, JP, and Pitney, WA. An electromyography analysis of 3 muscles surrounding the shoulder joint during the performance of a chest press exercise at several angles. *J. Strength Cond Res.* 24: 1925-1930, 2010.
 137. Vance, J, Wulf, G, Tollner, T, McNevin, N, and Mercer, J. EMG activity as a function of the performer's focus of attention. *J. Mot. Behav.* 36: 450-459, 2004.
 138. Wakahara, T, Miyamoto, N, Sugisaki, N, Murata, K, Kanehisa, H, Kawakami, Y, Fukunaga, T, and Yanai, T. Association between regional differences in muscle activation in one session of resistance exercise and in muscle hypertrophy after resistance training. *Eur. J. Appl. Physiol.* 112: 1569-1576, 2012.
 139. Wakahara, T, Fukutani, A, Kawakami, Y, and Yanai, T. Nonuniform muscle hypertrophy: its relation to muscle activation in training session. *Med. Sci. Sports Exerc.* 45: 2158-2165, 2013.
 140. Wakahara, T, Ema, R, Miyamoto, N, and Kawakami, Y. Inter- and intramuscular differences in training-induced hypertrophy of the quadriceps femoris: association with muscle activation during the first training session. *Clin. Physiol. Funct. Imaging* 37: 405-412, 2017.
 141. Watanabe, K, Otsuki, S, Hisa, T, and Nagaoka, M. Functional difference between the proximal and distal compartments of the semitendinosus muscle. *J. Phys. Ther. Sci.* 28: 1511-1517, 2016.
 142. Weiss, LW, Coney, HD, and Clark, FC. Gross measures of exercise-induced muscular hypertrophy. *J. Orthop. Sports Phys. Ther.* 30: 143-148, 2000.
 143. Wickham, JB, and Brown, JM. Muscles within muscles: the neuromotor control of intra-muscular segments. *Eur. J. Appl. Physiol. Occup. Physiol.* 78: 219-225, 1998.
 144. Wickiewicz, TL, Roy, RR, Powell, PL, and Edgerton, VR. Muscle architecture of the human lower limb. *Clin. Orthop. Relat. Res.* (179): 275-283, 1983.
 145. Wilk, KE, Escamilla, RF, Fleisig, GS, Barrentine, SW, Andrews, JR, and Boyd, ML. A comparison of tibiofemoral joint forces and electromyographic activity during open and closed kinetic chain exercises. *Am. J. Sports Med.* 24: 518-527, 1996.
 146. Willett, GM, Hyde, JE, Uhrlaub, MB, Wendel, CL, and Karst, GM. Relative activity of abdominal muscles during commonly prescribed strengthening exercises. *J. Strength Cond Res.* 15: 480-485, 2001.
 147. Willoughby, DS. The effects of mesocycle-length weight training programs involving periodization and partially equated volumes on upper and lower body strength. *J Strength Cond Res* 7: 2-8, 1993.
 148. Woodley, SJ, and Mercer, SR. Hamstring muscles: architecture and innervation. *Cells Tissues Organs* 179: 125-141, 2005.
 149. Worrell, TW, Karst, G, Adamczyk, D, Moore, R, Stanley, C, Steimel, B, and Steimel, S. Influence of joint position on electromyographic and torque generation during maximal voluntary isometric contractions of the hamstrings and gluteus maximus muscles. *J. Orthop. Sports Phys. Ther.* 31: 730-740, 2001.
 150. Wright, GA, DeLong, T, and Gehlsen, G. Electromyographic activity of the hamstrings during performance of the leg curl, stiff-leg deadlift and back squat movements. *Journal of Strength and Conditioning Research. J Strength Cond Res* 13: 168-174, 1999.
 151. Wulf, G. Attentional focus and motor learning: a review of 15 years. *International Review of Sport and Exercise Psychology* 6: 77-104, 2013.

152. Yamashita, N. EMG activities in mono- and bi-articular thigh muscles in combined hip and knee extension. *Eur. J. Appl. Physiol. Occup. Physiol.* 58: 274-277, 1988.
 153. Yavuz, HU, Erdag, D, Amca, AM, and Aritan, S. Kinematic and EMG activities during front and back squat variations in maximum loads. *J. Sports Sci.* 33: 1058-1066, 2015.
 154. Youdas, JW, Amundson, CL, Cicero, KS, Hahn, JJ, Harezlak, DT, and Hollman, JH. Surface electromyographic activation patterns and elbow joint motion during a pull-up, chin-up, or perfect-pullup rotational exercise. *J. Strength Cond Res.* 24: 3404-3414, 2010.
 155. Zatsiorsky, VM, and Kraemer, WJ. Science and Practice of Strength Training. Champaign, IL; Human Kinetics, 2006.
 156. Zebis, MK, Skotte, J, Andersen, CH, Mortensen, P, Petersen, HH, Viskaer, TC, Jensen, TL, Bencke, J, and Andersen, LL. Kettlebell swing targets semitendinosus and supine leg curl targets biceps femoris: an EMG study with rehabilitation implications. *Br. J. Sports Med.* 47: 1192-1198, 2013.
- ### Chapter 9
1. Adams, G, and Bamman, MM. Characterization and regulation of mechanical loading-induced compensatory muscle hypertrophy. *Comprehensive Physiology* 2829, 2012.
 2. Adechian, S, Balage, M, Remond, D, Migne, C, Quignard-Bou-lange, A, Marset-Baglieri, A, Rousset, S, Boirie, Y, Gaudichon, C, Dardevet, D, and Mosoni, L. Protein feeding pattern, casein feeding, or milk-soluble protein feeding did not change the evolution of body composition during a short-term weight loss program. *Am. J. Physiol. Endocrinol. Metab.* 303: E973-82, 2012.
 3. Alexander, JW. Immunonutrition: the role of omega-3 fatty acids. *Nutrition* 14: 627-633, 1998.
 4. Anderson, KE, Rosner, W, Khan, MS, New, MI, Pang, SY, Wissel, PS, and Kappas, A. Diet-hormone interactions: protein/carbohydrate ratio alters reciprocally the plasma levels of testosterone and cortisol and their respective binding globulins in man. *Life Sci.* 40: 1761-1768, 1987.
 5. Antonio, J, Ellerbroek, A, Silver, T, Orris, S, Scheiner, M, Gonzalez, A, and Peacock, CA. A high protein diet (3.4 g/kg/d) combined with a heavy resistance training program improves body composition in healthy trained men and women—a follow-up investigation. *J. Int. Soc. Sports Nutr.* 12: 39-015-0100-0. eCollection 2015, 2015.
 6. Aragon, AA, and Schoenfeld, BJ. Nutrient timing revisited: is there a post-exercise anabolic window? *J. Int. Soc. Sports Nutr.* 10: 5-2783-10-5, 2013.
 7. Areta, JL, Burke, LM, Ross, ML, Camera, DM, West, DW, Broad, EM, Jeacocke, NA, Moore, DR, Stellingwerff, T, Phillips, SM, Hawley, JA, and Coffey, VG. Timing and distribution of protein ingestion during prolonged recovery from resistance exercise alters myofibrillar protein synthesis. *J. Physiol.* 591: 2319-2331, 2013.
 8. Areta, JL, Burke, LM, Camera, DM, West, DW, Crawshaw, S, Moore, DR, Stellingwerff, T, Phillips, SM, Hawley, JA, and Coffey, VG. Reduced resting skeletal muscle protein synthesis is rescued by resistance exercise and protein ingestion following short-term energy deficit. *Am. J. Physiol. Endocrinol. Metab.* 306: E989-97, 2014.
 9. Arnal, MA, Mosoni, L, Boirie, Y, Houlier, ML, Morin, L, Verdier, E, Ritz, P, Antoine, JM, Prugnaud, J, Beaufre, B, and Mirand, PP. Protein pulse feeding improves protein retention in elderly women. *Am. J. Clin. Nutr.* 69: 1202-1208, 1999.
 10. Arnal, MA, Mosoni, L, Boirie, Y, Houlier, ML, Morin, L, Verdier, E, Ritz, P, Antoine, JM, Prugnaud, J, Beaufre, B, and Mirand, PP. Protein feeding pattern does not affect protein retention in young women. *J. Nutr.* 130: 1700-1704, 2000.
 11. Atherton, PJ, Etheridge, T, Watt, PW, Wilkinson, D, Selby, A, Rankin, D, Smith, K, and Rennie, MJ. Muscle full effect after oral protein: time-dependent concordance and discordance between human muscle protein synthesis and mTORC1 signaling. *Am. J. Clin. Nutr.* 92: 1080-1088, 2010.
 12. Atherton, PJ, and Smith, K. Muscle protein synthesis in response to nutrition and exercise. *J. Physiol.* 590: 1049-1057, 2012.
 13. Atherton, PJ. Is there an optimal time for warfighters to supplement with protein? *J. Nutr.* 143: 1848S-1851S, 2013.
 14. Bandegan, A, Courtney-Martin, G, Rafii, M, Pencharz, PB, and Lemon, PW. Indicator Amino Acid-Derived Estimate of Dietary Protein Requirement for Male Bodybuilders on a Nontraining Day Is Several-Fold Greater than the Current Recommended Dietary Allowance. *J. Nutr.* 147: 850-857, 2017.
 15. Bergeron, K, Julien, P, Davis, TA, Myre, A, and Thivierge, MC. Long-chain n-3 fatty acids enhance neonatal insulin-regulated protein metabolism in piglets by differentially altering muscle lipid composition. *J. Lipid Res.* 48: 2396-2410, 2007.
 16. Bilsborough, S, and Mann, N. A review of issues of dietary protein intake in humans. *Int. J. Sport Nutr. Exerc. Metab.* 16: 129-152, 2006.
 17. Biolo, G, Williams, BD, Fleming, RY, and Wolfe, RR. Insulin action on muscle protein kinetics and amino acid transport during recovery after resistance exercise. *Diabetes* 48: 949-957, 1999.
 18. Borkman, M, Storlien, LH, Pan, DA, Jenkins, AB, Chisholm, DJ, and Campbell, LV. The relation between insulin sensitivity and the fatty-acid composition of skeletal-muscle phospholipids. *N. Engl. J. Med.* 328: 238-244, 1993.
 19. Borsheim, E, Tipton, KD, Wolf, SE, and Wolfe, RR. Essential amino acids and muscle protein recovery from resistance exercise. *Am. J. Physiol. Endocrinol. Metab.* 283: E648-57, 2002.
 20. Breen, L, and Churchward-Venne, TA. Leucine: a nutrient 'trigger' for muscle anabolism, but what more? *J. Physiol.* 590: 2065-2066, 2012.
 21. Camera, DM, West, DW, Burd, NA, Phillips, SM, Garnham, AP, Hawley, JA, and Coffey, VG. Low muscle glycogen concentration does not suppress the anabolic response to resistance exercise. *J. Appl. Physiol.* 113: 206-214, 2012.
 22. Campbell, B, Kreider, RB, Ziegenfuss, T, La Bounty, P, Roberts, M, Burke, D, Landis, J, Lopez, H, and Antonio, J. International Society of Sports Nutrition position stand: protein and exercise. *J. Int. Soc. Sports Nutr.* 4: 8, 2007.
 23. Campbell, BI, Aguilar, D, Conlin, L, Vargas, A, Schoenfeld, BJ, Corson, A, Gai, C, Best, S, Galvan, E, and Couvillion, K. Effects of High Versus Low Protein Intake on Body Composition and Maximal Strength in Aspiring Female Physique Athletes Engaging in an 8-Week Resistance Training Program. *Int. J. Sport Nutr. Exerc. Metab.* 28: 580-585, 2018.
 24. Candow, DG, and Chilibeck, PD. Timing of creatine or protein supplementation and resistance training in the elderly. *Appl. Physiol. Nutr. Metab.* 33: 184-190, 2008.
 25. Capaldo, B, Gastaldelli, A, Antonello, S, Auletta, M, Pardo, F, Ciociaro, D, Guida, R, Ferrannini, E, and Sacca, L. Splanchnic and leg substrate exchange after ingestion of a natural mixed meal in humans. *Diabetes* 48: 958-966, 1999.
 26. Churchley, EG, Coffey, VG, Pedersen, DJ, Shield, A, Carey, KA, Cameron-Smith, D, and Hawley, JA. Influence of preexercise muscle glycogen content on transcriptional activity of metabolic and myogenic genes in well-trained humans. *J. Appl. Physiol.* 102: 1604-1611, 2007.

27. Churchward-Venne, TA, Murphy, CH, Longland, TM, and Phillips, SM. Role of protein and amino acids in promoting lean mass accretion with resistance exercise and attenuating lean mass loss during energy deficit in humans. *Amino Acids* 45: 231-240, 2013.
28. Coffey, VG, Shield, A, Canny, BJ, Carey, KA, Cameron-Smith, D, and Hawley, JA. Interaction of contractile activity and training history on mRNA abundance in skeletal muscle from trained athletes. *Am. J. Physiol. Endocrinol. Metab.* 290: E849-55, 2006.
29. Creer, A, Gallagher, P, Slivka, D, Jemiolo, B, Fink, W, and Trappe, S. Influence of muscle glycogen availability on ERK1/2 and Akt signaling after resistance exercise in human skeletal muscle. *J. Appl. Physiol.* 99: 950-956, 2005.
30. Denne, SC, Liechty, EA, Liu, YM, Brechtel, G, and Baron, AD. Proteolysis in skeletal muscle and whole body in response to euglycemic hyperinsulinemia in normal adults. *Am. J. Physiol.* 261: E809-14, 1991.
31. Dennis, PB, Jaeschke, A, Saitoh, M, Fowler, B, Kozma, SC, and Thomas, G. Mammalian TOR: a homeostatic ATP sensor. *Science* 294: 1102-1105, 2001.
32. Derave, W, Lund, S, Holman, GD, Wojtaszewski, J, Pedersen, O, and Richter, EA. Contraction-stimulated muscle glucose transport and GLUT-4 surface content are dependent on glycogen content. *Am. J. Physiol.* 277: E1103-10, 1999.
33. Deutz, NE, and Wolfe, RR. Is there a maximal anabolic response to protein intake with a meal? *Clin. Nutr.* 32: 309-313, 2013.
34. Dideriksen, K, Reitelseder, S, and Holm, L. Influence of amino acids, dietary protein, and physical activity on muscle mass development in humans. *Nutrients* 5: 852-876, 2013.
35. Dorgan, JF, Judd, JT, Longcope, C, Brown, C, Schatzkin, A, Clevidence, BA, Campbell, WS, Nair, PP, Franz, C, Kahle, L, and Taylor, PR. Effects of dietary fat and fiber on plasma and urine androgens and estrogens in men: a controlled feeding study. *Am. J. Clin. Nutr.* 64: 850-855, 1996.
36. Drummond, MJ, Dreyer, HC, Fry, CS, Glynn, EL, and Rasmussen, BB. Nutritional and contractile regulation of human skeletal muscle protein synthesis and mTORC1 signaling. *J. Appl. Physiol.* (1985) 106: 1374-1384, 2009.
37. Edmonds, MS, and Baker, DH. Amino acid excesses for young pigs: effects of excess methionine, tryptophan, threonine or leucine. *J. Anim. Sci.* 64: 1664-1671, 1987.
38. Essen-Gustavsson, B, and Tesch, PA. Glycogen and triglyceride utilization in relation to muscle metabolic characteristics in men performing heavy-resistance exercise. *Eur. J. Appl. Physiol. Occup. Physiol.* 61: 5-10, 1990.
39. Fluck, M. Regulation of protein synthesis in skeletal muscle. *Dtsch Z Sportmed* 63: 75-80, 2012.
40. Fluckey, JD, Vary, TC, Jefferson, LS, and Farrell, PA. Augmented insulin action on rates of protein synthesis after resistance exercise in rats. *Am. J. Physiol.* 270: E313-9, 1996.
41. Fox, AK, Kaufman, AE, and Horowitz, JF. Adding fat calories to meals after exercise does not alter glucose tolerance. *J. Appl. Physiol.* 97: 11-16, 2004.
42. Garthe, I, Raastad, T, Refsnes, PE, and Sundgot-Borgen, J. Effect of nutritional intervention on body composition and performance in elite athletes. *Eur. J. Sport. Sci.* 13: 295-303, 2013.
43. Gelfand, RA, and Barrett, EJ. Effect of physiologic hyperinsulinemia on skeletal muscle protein synthesis and breakdown in man. *J. Clin. Invest.* 80: 1-6, 1987.
44. Gingras, AA, White, PJ, Chouinard, PY, Julien, P, Davis, TA, Dombrowski, L, Couture, Y, Dubreuil, P, Myre, A, Bergeron, K, Marette, A, and Thivierge, MC. Long-chain omega-3 fatty acids regulate bovine whole-body protein metabolism by promoting muscle insulin signalling to the Akt-mTOR-S6K1 pathway and insulin sensitivity. *J. Physiol.* 579: 269-284, 2007.
45. Glynn, EL, Fry, CS, Timmerman, KL, Drummond, MJ, Volpi, E, and Rasmussen, BB. Addition of carbohydrate or alanine to an essential amino Acid mixture does not enhance human skeletal muscle protein anabolism. *J. Nutr.* 143: 307-314, 2013.
46. Goodman, CA, Mayhew, DL, and Hornberger, TA. Recent progress toward understanding the molecular mechanisms that regulate skeletal muscle mass. *Cell. Signal.* 23: 1896-1906, 2011.
47. Greene, DA, Varley, BJ, Hartwig, TB, Chapman, P, and Rigney, M. A Low-Carbohydrate Ketogenic Diet Reduces Body Mass Without Compromising Performance in Powerlifting and Olympic Weightlifting Athletes. *J. Strength Cond Res.* 32: 3373-3382, 2018.
48. Greenhaff, PL, Karagounis, LG, Peirce, N, Simpson, EJ, Hazell, M, Layfield, R, Wackerhage, H, Smith, K, Atherton, P, Selby, A, and Rennie, MJ. Disassociation between the effects of amino acids and insulin on signaling, ubiquitin ligases, and protein turnover in human muscle. *Am. J. Physiol. Endocrinol. Metab.* 295: E595-604, 2008.
49. Gropper, SS, Smith, JL, and Groff, JL. Advanced Nutrition and Human Metabolism. Belmont, CA; Wadsworth Cengage Learning, 2009.
50. Hamalainen, EK, Adlercreutz, H, Puska, P, and Pietinen, P. Decrease of serum total and free testosterone during a low-fat high-fibre diet. *J. Steroid Biochem.* 18: 369-370, 1983.
51. Hamer, HM, Wall, BT, Kiskini, A, de Lange, A, Groen, BB, Bakker, JA, Gijsen, AP, Verdijk, LB, and van Loon, LJ. Carbohydrate co-ingestion with protein does not further augment post-prandial muscle protein accretion in older men. *Nutr. Metab. (Lond)* 10: 15-7075-10-15, 2013.
52. Hawley, JA, Burke, LM, Phillips, SM, and Spriet, LL. Nutritional modulation of training-induced skeletal muscle adaptations. *J. Appl. Physiol.* (1985) 110: 834-845, 2011.
53. Helms, ER, Aragon, AA, and Fitschen, PJ. Evidence-based recommendations for natural bodybuilding contest preparation: nutrition and supplementation. *J. Int. Soc. Sports Nutr.* 11: 20-2783-11-20. eCollection 2014, 2014.
54. Helms, ER, Zinn, C, Rowlands, DS, and Brown, SR. A systematic review of dietary protein during caloric restriction in resistance trained lean athletes: a case for higher intakes. *Int. J. Sport Nutr. Exerc. Metab.* 24: 127-138, 2014.
55. Hepburn, D, and Maughan, RJ. Glycogen availability as a limiting factor in the performance of isometric exercise. *J. Physiol.* 342: 52P-53P, 1982.
56. Heslin, MJ, Newman, E, Wolf, RF, Pisters, PW, and Brennan, MF. Effect of hyperinsulinemia on whole body and skeletal muscle leucine carbon kinetics in humans. *Am. J. Physiol.* 262: E911-8, 1992.
57. Holwerda, AM, Paulussen, KJM, Overkamp, M, Goessens, JPB, Kramer, IE, Wodzig, WKWH, Verdijk, LB, and van Loon, LJC. Dose-Dependent Increases in Whole-Body Net Protein Balance and Dietary Protein-Derived Amino Acid Incorporation into Myofibrillar Protein During Recovery from Resistance Exercise in Older Men. *J. Nutr.* 149: 221-230, 2019.
58. Ivy, J, and Ferguson-Stegall, L. Nutrient Timing: The Means to Improved Exercise Performance, Recovery, and Training. *American Journal of Lifestyle Medicine* : 1-14, 2013.
59. Ivy, JL. Glycogen resynthesis after exercise: effect of carbohydrate intake. *Int. J. Sports Med.* 19 Suppl 2: S142-5, 1998.
60. Jabekk, PT, Moe, IA, Meen, HD, Tomten, SE, and Hostmark, AT. Resistance training in overweight women on a ketogenic diet conserved lean body mass while reducing body fat. *Nutr. Metab. (Lond)* 7: 17-7075-7-17, 2010.

61. Joy, JM, Vogel, RM, Shane Broughton, K, Kudla, U, Kerr, NY, Davison, JM, Wildman, REC, and DiMarco, NM. Daytime and nighttime casein supplements similarly increase muscle size and strength in response to resistance training earlier in the day: a preliminary investigation. *J. Int. Soc. Sports Nutr.* 15: 24-018-0228-9, 2018.
62. Katsanos, CS, Kobayashi, H, Sheffield-Moore, M, Aarsland, A, and Wolfe, RR. A high proportion of leucine is required for optimal stimulation of the rate of muscle protein synthesis by essential amino acids in the elderly. *Am. J. Physiol. Endocrinol. Metab.* 291: E381-7, 2006.
63. Kawanaka, K, Nolte, LA, Han, DH, Hansen, PA, and Holloszy, JO. Mechanisms underlying impaired GLUT-4 translocation in glycogen-supercompensated muscles of exercised rats. *Am. J. Physiol. Endocrinol. Metab.* 279: E1311-8, 2000.
64. Kephart, WC, Pledge, CD, Roberson, PA, Mumford, PW, Romero, MA, Mobley, CB, Martin, JS, Young, KC, Lowery, RP, Wilson, JM, Huggins, KW, and Roberts, MD. The Three-Month Effects of a Ketogenic Diet on Body Composition, Blood Parameters, and Performance Metrics in CrossFit Trainees: A Pilot Study. *Sports (Basel)* 6: 10.3390/sports6010001, 2018.
65. Kerksick, C, Harvey, T, Stout, J, Campbell, B, Wilborn, C, Kreider, R, Kalman, D, Ziegenfuss, T, Lopez, H, Landis, J, Ivy, JL, and Antonio, J. International Society of Sports Nutrition position stand: nutrient timing. *J. Int. Soc. Sports Nutr.* 5: 17, 2008.
66. Kettelhut, IC, Wing, SS, and Goldberg, AL. Endocrine regulation of protein breakdown in skeletal muscle. *Diabetes Metab. Rev.* 4: 751-772, 1988.
67. Kim, DH, Kim, JY, Yu, BP, and Chung, HY. The activation of NF-kappaB through Akt-induced FOXO1 phosphorylation during aging and its modulation by calorie restriction. *Biogerontology* 9: 33-47, 2008.
68. Kimball, SR. Integration of signals generated by nutrients, hormones, and exercise in skeletal muscle. *Am. J. Clin. Nutr.* 99: 237S-242S, 2014.
69. Koopman, R, Manders, RJ, Jonkers, RA, Hul, GB, Kuipers, H, and van Loon, LJ. Intramyocellular lipid and glycogen content are reduced following resistance exercise in untrained healthy males. *Eur. J. Appl. Physiol.* 96: 525-534, 2006.
70. Koopman, R, Beelen, M, Stellingwerff, T, Pennings, B, Saris, WH, Kies, AK, Kuipers, H, and van Loon, LJ. Coingestion of carbohydrate with protein does not further augment postexercise muscle protein synthesis. *Am. J. Physiol. Endocrinol. Metab.* 293: E833-42, 2007.
71. Kumar, V, Atherton, P, Smith, K, and Rennie, MJ. Human muscle protein synthesis and breakdown during and after exercise. *J. Appl. Physiol.* 106: 2026-2039, 2009.
72. Lambert, CP, and Flynn, MG. Fatigue during high-intensity intermittent exercise: application to bodybuilding. *Sports Med.* 32: 511-522, 2002.
73. Lane, AR, Duke, JW, and Hackney, AC. Influence of dietary carbohydrate intake on the free testosterone: cortisol ratio responses to short-term intensive exercise training. *Eur. J. Appl. Physiol.* 108: 1125-1131, 2010.
74. Langfort, J, Zarzecny, R, Pilis, W, Nazar, K, and Kaciuba-Usitko, H. The effect of a low-carbohydrate diet on performance, hormonal and metabolic responses to a 30-s bout of supramaximal exercise. *Eur. J. Appl. Physiol. Occup. Physiol.* 76: 128-133, 1997.
75. Layman, DK. Protein quantity and quality at levels above the RDA improves adult weight loss. *J. Am. Coll. Nutr.* 23: 631S-636S, 2004.
76. Lemon, PW, Tarnopolsky, MA, MacDougall, JD, and Atkinson, SA. Protein requirements and muscle mass/strength changes during intensive training in novice bodybuilders. *J. Appl. Physiol.* 73: 767-775, 1992.
77. Levenhagen, DK, Gresham, JD, Carlson, MC, Maron, DJ, Borel, MJ, and Flakoll, PJ. Postexercise nutrient intake timing in humans is critical to recovery of leg glucose and protein homeostasis. *Am. J. Physiol. Endocrinol. Metab.* 280: E982-93, 2001.
78. Leveritt, M, and Abernethy, PJ. Effects of carbohydrate restriction on strength performance. 13: 52-57, 1999.
79. Lima-Silva, AE, Pires, FO, Bertuzzi, R, Silva-Cavalcante, MD, Oliveira, RS, Kiss, MA, and Bishop, D. Effects of a low- or a high-carbohydrate diet on performance, energy system contribution, and metabolic responses during supramaximal exercise. *Appl. Physiol. Nutr. Metab.* 38: 928-934, 2013.
80. Longland, TM, Oikawa, SY, Mitchell, CJ, Devries, MC, and Phillips, SM. Higher compared with lower dietary protein during an energy deficit combined with intense exercise promotes greater lean mass gain and fat mass loss: a randomized trial. *Am. J. Clin. Nutr.* 103: 738-746, 2016.
81. Lynch, CJ, Hutson, SM, Patson, BJ, Vaval, A, and Vary, TC. Tissue-specific effects of chronic dietary leucine and norleucine supplementation on protein synthesis in rats. *Am. J. Physiol. Endocrinol. Metab.* 283: E824-35, 2002.
82. MacDougall, JD, Ray, S, Sale, DG, McCartney, N, Lee, P, and Garner, S. Muscle substrate utilization and lactate production. *Can. J. Appl. Physiol.* 24: 209-215, 1999.
83. MacKenzie-Shalders, KL, King, NA, Byrne, NM, and Slater, GJ. Increasing Protein Distribution Has No Effect on Changes in Lean Mass During a Rugby Preseason. *Int. J. Sport Nutr. Exerc. Metab.* 26: 1-7, 2016.
84. Macnaughton, LS, Wardle, SL, Witard, OC, McGlory, C, Hamilton, DL, Jeromson, S, Lawrence, CE, Wallis, GA, and Tipton, KD. The response of muscle protein synthesis following whole-body resistance exercise is greater following 40 g than 20 g of ingested whey protein. *Physiol. Rep.* 4: 10.14814/phy2.12893, 2016.
85. Mamerow, MM, Mettler, JA, English, KL, Casperson, SL, Arentson-Lantz, E, Sheffield-Moore, M, Layman, DK, and Paddon-Jones, D. Dietary Protein Distribution Positively Influences 24-h Muscle Protein Synthesis in Healthy Adults. *J. Nutr.*, 2014.
86. Margolis, LM, Rivas, DA, Berrone, M, Ezzyat, Y, Young, AJ, McClung, JP, Fielding, RA, and Pasiakos, SM. Prolonged Calorie Restriction Downregulates Skeletal Muscle mTORC1 Signaling Independent of Dietary Protein Intake and Associated microRNA Expression. *Front. Physiol.* 7: 445, 2016.
87. McBride, A, Ghilagaber, S, Nikolaev, A, and Hardie, DG. The glycogen-binding domain on the AMPK beta subunit allows the kinase to act as a glycogen sensor. *Cell. Metab.* 9: 23-34, 2009.
88. McIver, CM, Wycherley, TP, and Clifton, PM. mTOR signaling and ubiquitin-proteasome gene expression in the preservation of fat free mass following high protein, calorie restricted weight loss. *Nutr. Metab. (Lond)* 9: 83-7075-9-83, 2012.
89. Meirelles, CM, and Gomes, PS. Effects of Short-Term Carbohydrate Restrictive and Conventional Hypoenergetic Diets and Resistance Training on Strength Gains and Muscle Thickness. *J. Sports Sci. Med.* 15: 578-584, 2016.
90. Mitchell, JB, DiLauro, PC, Pizza, FX, and Cavender, DL. The effect of preexercise carbohydrate status on resistance exercise performance. *Int. J. Sport Nutr.* 7: 185-196, 1997.
91. Moore, DR, Del Bel, NC, Nizi, KI, Hartman, JW, Tang, JE, Armstrong, D, and Phillips, SM. Resistance training reduces fasted- and fed-state leucine turnover and increases dietary nitrogen retention in previously untrained young men. *J. Nutr.* 137: 985-991, 2007.

92. Moore, DR, Robinson, MJ, Fry, JL, Tang, JE, Glover, EI, Wilkinson, SB, Prior, T, Tarnopolsky, MA, and Phillips, SM. Ingested protein dose response of muscle and albumin protein synthesis after resistance exercise in young men. *Am. J. Clin. Nutr.* 89: 161-168, 2009.
93. Moro, T, Tinsley, G, Bianco, A, Marcolin, G, Pacelli, QF, Battaglia, G, Palma, A, Gentil, P, Neri, M, and Paoli, A. Effects of eight weeks of time-restricted feeding (16/8) on basal metabolism, maximal strength, body composition, inflammation, and cardiovascular risk factors in resistance-trained males. *J. Transl. Med.* 14: 290, 2016.
94. Morton, RW, McGlory, C, and Phillips, SM. Nutritional interventions to augment resistance training-induced skeletal muscle hypertrophy. *Front. Physiol.* 6: 245, 2015.
95. Morton, RW, Murphy, KT, McKellar, SR, Schoenfeld, BJ, Henselmans, M, Helms, E, Aragon, AA, Devries, MC, Banfield, L, Krieger, JW, and Phillips, SM. A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults. *Br. J. Sports Med.*, 2017.
96. Murphy, MG. Dietary fatty acids and membrane protein function. *J. Nutr. Biochem.* 1: 68-79, 1990.
97. Noreen, EE, Sass, MJ, Crowe, ML, Pabon, VA, Brandauer, J, and Averill, LK. Effects of supplemental fish oil on resting metabolic rate, body composition, and salivary cortisol in healthy adults. *J. Int. Soc. Sports Nutr.* 7: 31-2783-7-31, 2010.
98. O'Gorman, DJ, Del Aguila, LE, Williamson, DL, Krishnan, RK, and Kirwan, JP. Insulin and exercise differentially regulate PI3-kinase and glycogen synthase in human skeletal muscle. *J. Appl. Physiol.* 89: 1412-1419, 2000.
99. Okamura, K, Doi, T, Hamada, K, Sakurai, M, Matsumoto, K, Imaizumi, K, Yoshioka, Y, Shimizu, S, and Suzuki, M. Effect of amino acid and glucose administration during postexercise recovery on protein kinetics in dogs. *Am. J. Physiol.* 272: E1023-30, 1997.
100. Ortenblad, N, Westerblad, H, and Nielsen, J. Muscle glycogen stores and fatigue. *J. Physiol.* 591: 4405-4413, 2013.
101. Paoli, A, Grimaldi, K, D'Agostino, D, Cenci, L, Moro, T, Bianco, A, and Palma, A. Ketogenic diet does not affect strength performance in elite artistic gymnasts. *J. Int. Soc. Sports Nutr.* 9: 34-2783-9-34, 2012.
102. Parkin, JA, Carey, MF, Martin, IK, Stojanovska, L, and Febbraio, MA. Muscle glycogen storage following prolonged exercise: effect of timing of ingestion of high glycemic index food. *Med. Sci. Sports Exerc.* 29: 220-224, 1997.
103. Pasiakos, SM, Vislocky, LM, Carbone, JW, Altieri, N, Konopelski, K, Freake, HC, Anderson, JM, Ferrando, AA, Wolfe, RR, and Rodriguez, NR. Acute energy deprivation affects skeletal muscle protein synthesis and associated intracellular signaling proteins in physically active adults. *J. Nutr.* 140: 745-751, 2010.
104. Pasiakos, SM. Exercise and amino acid anabolic cell signaling and the regulation of skeletal muscle mass. *Nutrients* 4: 740-758, 2012.
105. Pasiakos, SM, and Carbone, JW. Assessment of skeletal muscle proteolysis and the regulatory response to nutrition and exercise. *IUBMB Life* 66: 478-484, 2014.
106. Phillips, SM. The science of muscle hypertrophy: making dietary protein count. *Proc. Nutr. Soc.* 70: 100-103, 2011.
107. Phillips, SM, and Van Loon, LJ. Dietary protein for athletes: from requirements to optimum adaptation. *J. Sports Sci.* 29 Suppl 1: S29-38, 2011.
108. Phillips, SM. Current Concepts and Unresolved Questions in Dietary Protein Requirements and Supplements in Adults. *Front. Nutr.* 4: 13, 2017.
109. Pitkanen, HT, Nykanen, T, Knuutinen, J, Lahti, K, Keinanen, O, Alen, M, Komi, PV, and Mero, AA. Free amino acid pool and muscle protein balance after resistance exercise. *Med. Sci. Sports Exerc.* 35: 784-792, 2003.
110. Power, O, Hallihan, A, and Jakeman, P. Human insulinotropic response to oral ingestion of native and hydrolysed whey protein. *Amino Acids* 37: 333-339, 2009.
111. Rasmussen, BB, Tipton, KD, Miller, SL, Wolf, SE, and Wolfe, RR. An oral essential amino acid-carbohydrate supplement enhances muscle protein anabolism after resistance exercise. *J. Appl. Physiol.* 88: 386-392, 2000.
112. Reidy, PT, Walker, DK, Dickinson, JM, Gundermann, DM, Drummond, MJ, Timmerman, KL, Fry, CS, Borack, MS, Cope, MB, Mukherjee, R, Jennings, K, Volpi, E, and Rasmussen, BB. Protein blend ingestion following resistance exercise promotes human muscle protein synthesis. *J. Nutr.* 143: 410-416, 2013.
113. Rennie, MJ, Bohe, J, Smith, K, Wackerhage, H, and Greenhaff, P. Branched-chain amino acids as fuels and anabolic signals in human muscle. *J. Nutr.* 136: 264S-8S, 2006.
114. Ribeiro, AS, Nunes, JP, Schoenfeld, BJ, Aguiar, AF, and Cyrino, ES. Effects of different dietary energy intake following resistance training on muscle mass and body fat in bodybuilders: a pilot study. *J Hum Kinet.* DOI: 10.2478/hukin-2019-0038, 2019.
115. Ribeiro, AS, Nunes, JP, and Schoenfeld, BJ. Should Competitive Bodybuilders Ingest More Protein than Current Evidence-Based Recommendations? *Sports Med.* 49: 1481-1485, 2019.
116. Richter, EA, Derave, W, and Wojtaszewski, JF. Glucose, exercise and insulin: emerging concepts. *J. Physiol.* 535: 313-322, 2001.
117. Rosqvist, E, Igman, D, Kullberg, J, Cedernaes, J, Johansson, HE, Larsson, A, Johansson, L, Ahlstrom, H, Arner, P, Dahlman, I, and Riserus, U. Overfeeding polyunsaturated and saturated fat causes distinct effects on liver and visceral fat accumulation in humans. *Diabetes* 63: 2356-2368, 2014.
118. Rossato, LT, Schoenfeld, BJ, and de Oliveira, EP. Is there sufficient evidence to supplement omega-3 fatty acids to increase muscle mass and strength in young and older adults? *Clin. Nutr.*, 2019.
119. Rozenek, R, Ward, P, Long, S, and Garhammer, J. Effects of high-calorie supplements on body composition and muscular strength following resistance training. *J. Sports Med. Phys. Fitness* 42: 340-347, 2002.
120. Ryan, AM, Reynolds, JV, Healy, L, Byrne, M, Moore, J, Brannelly, N, McHugh, A, McCormack, D, and Flood, P. Enteral nutrition enriched with eicosapentaenoic acid (EPA) preserves lean body mass following esophageal cancer surgery: results of a double-blinded randomized controlled trial. *Ann. Surg.* 249: 355-363, 2009.
121. Sallinen, J, Pakarinen, A, Ahtiainen, J, Kraemer, WJ, Volek, JS, and Häkkinen, K. Relationship between diet and serum anabolic hormone responses to heavy-resistance exercise in men. *Int. J. Sports Med.* 25: 627-633, 2004.
122. Schaafsma, G. The protein digestibility-corrected amino acid score. *J. Nutr.* 130: 1865S-7S, 2000.
123. Schoeller, DA. How accurate is self-reported dietary energy intake? *Nutr. Rev.* 48: 373-379, 1990.
124. Schoenfeld, BJ, Aragon, AA, and Krieger, JW. The effect of protein timing on muscle strength and hypertrophy: a meta-analysis. *J. Int. Soc. Sports Nutr.* 10: 53-2783-10-53, 2013.
125. Schoenfeld, BJ, Aragon, A, Wilborn, C, Urbina, SL, Hayward, SE, and Krieger, J. Pre- versus post-exercise protein intake has similar effects on muscular adaptations. *PeerJ* 5: e2825, 2017.
126. Schoenfeld, BJ, and Aragon, AA. How much protein can the body use in a single meal for muscle-building? Implications

- for daily protein distribution. *J. Int. Soc. Sports Nutr.* 15: 10-018-0215-1. eCollection 2018, 2018.
127. Simmons, E, Fluckey, JD, and Riechman, SE. Cumulative Muscle Protein Synthesis and Protein Intake Requirements. *Annu. Rev. Nutr.* 36: 17-43, 2016.
 128. Slater, G, and Phillips, SM. Nutrition guidelines for strength sports: sprinting, weightlifting, throwing events, and bodybuilding. *J. Sports Sci.* 29 Suppl 1: S67-77, 2011.
 129. Smith, GI, Atherton, P, Reeds, DN, Mohammed, BS, Rankin, D, Rennie, MJ, and Mittendorfer, B. Dietary omega-3 fatty acid supplementation increases the rate of muscle protein synthesis in older adults: a randomized controlled trial. *Am. J. Clin. Nutr.* 93: 402-412, 2011.
 130. Snijders, T, Trommelen, J, Kouw, IWK, Holwerda, AM, Verdijk, LB, and van Loon, LJC. The Impact of Pre-sleep Protein Ingestion on the Skeletal Muscle Adaptive Response to Exercise in Humans: An Update. *Front. Nutr.* 6: 17, 2019.
 131. Sonmez, GT, Schoenfeld, BJ, and Vatansever-Ozen, S. Omega-3 fatty acids and exercise: a review of their combined effect on body composition and physical performance. *Biomedical Human Kinetics* 3: 23-29, 2011.
 132. Staples, AW, Burd, NA, West, DW, Currie, KD, Atherton, PJ, Moore, DR, Rennie, MJ, Macdonald, MJ, Baker, SK, and Phillips, SM. Carbohydrate does not augment exercise-induced protein accretion versus protein alone. *Med. Sci. Sports Exerc.* 43: 1154-1161, 2011.
 133. Stokes, T, Hector, AJ, Morton, RW, McGlory, C, and Phillips, SM. Recent Perspectives Regarding the Role of Dietary Protein for the Promotion of Muscle Hypertrophy with Resistance Exercise Training. *Nutrients* 10: 10.3390/nu10020180, 2018.
 134. Tang, JE, Moore, DR, Kujbida, GW, Tarnopolsky, MA, and Phillips, SM. Ingestion of whey hydrolysate, casein, or soy protein isolate: effects on mixed muscle protein synthesis at rest and following resistance exercise in young men. *J. Appl. Physiol.* (1985) 107: 987-992, 2009.
 135. Tarnopolsky, MA, Atkinson, SA, MacDougall, JD, Chesley, A, Phillips, S, and Swartz, HP. Evaluation of protein requirements for trained strength athletes. *J. Appl. Physiol.* (1985) 73: 1986-1995, 1992.
 136. Timmons, JA. Variability in training-induced skeletal muscle adaptation. *J. Appl. Physiol.* 110: 846-853, 2011.
 137. Tinsley, GM, Forsse, JS, Butler, NK, Paoli, A, Bane, AA, La Bounty, PM, Morgan, GB, and Grandjean, PW. Time-restricted feeding in young men performing resistance training: A randomized controlled trial. *Eur. J. Sport. Sci.* 17: 200-207, 2017.
 138. Tinsley, GM, Moore, ML, Graybeal, AJ, Paoli, A, Kim, Y, Gonzales, JU, Harry, JR, VanDusseldorp, TA, Kennedy, DN, and Cruz, MR. Time-restricted feeding plus resistance training in active females: a randomized trial. *Am. J. Clin. Nutr.* , 2019.
 139. Tipton, KD, Ferrando, AA, Phillips, SM, Doyle, D, Jr, and Wolfe, RR. Postexercise net protein synthesis in human muscle from orally administered amino acids. *Am. J. Physiol.* 276: E628-34, 1999.
 140. Tipton, KD, Gurkin, BE, Matin, S, and Wolfe, RR. Nonessential amino acids are not necessary to stimulate net muscle protein synthesis in healthy volunteers. *J. Nutr. Biochem.* 10: 89-95, 1999.
 141. Tipton, KD, Elliott, TA, Cree, MG, Wolf, SE, Sanford, AP, and Wolfe, RR. Ingestion of casein and whey proteins result in muscle anabolism after resistance exercise. *Med. Sci. Sports Exerc.* 36: 2073-2081, 2004.
 142. Tipton, KD, Elliott, TA, Ferrando, AA, Aarsland, AA, and Wolfe, RR. Stimulation of muscle anabolism by resistance exercise and ingestion of leucine plus protein. *Appl. Physiol. Nutr. Metab.* 34: 151-161, 2009.
 143. van Wessel, T, de Haan, A, van der Laarse, WJ, and Jaspers, RT. The muscle fiber type-fiber size paradox: hypertrophy or oxidative metabolism? *Eur. J. Appl. Physiol.* 110: 665-694, 2010.
 144. Vargas, S, Romance, R, Petro, JL, Bonilla, DA, Galancho, I, Espinar, S, Kreider, RB, and Benitez-Porres, J. Efficacy of ketogenic diet on body composition during resistance training in trained men: a randomized controlled trial. *J. Int. Soc. Sports Nutr.* 15: 31-018-0236-9, 2018.
 145. Volek, JS, Kraemer, WJ, Bush, JA, Incledon, T, and Boetes, M. Testosterone and cortisol in relationship to dietary nutrients and resistance exercise. *J. Appl. Physiol.* (1985) 82: 49-54, 1997.
 146. Volek, JS, Gomez, AL, Love, DM, Avery, NG, Sharman, MJ, and Kraemer, WJ. Effects of a high-fat diet on postabsorptive and postprandial testosterone responses to a fat-rich meal. *Metabolism* 50: 1351-1355, 2001.
 147. Wax, B, Kavazis, AN, and Brown, SP. Effects of supplemental carbohydrate ingestion during superimposed electromyostimulation exercise in elite weightlifters. *J. Strength Cond Res.* 27: 3084-3090, 2013.
 148. Whitehouse, AS, and Tisdale, MJ. Downregulation of ubiquitin-dependent proteolysis by eicosapentaenoic acid in acute starvation. *Biochem. Biophys. Res. Commun.* 285: 598-602, 2001.
 149. Wilkinson, SB, Tarnopolsky, MA, Macdonald, MJ, Macdonald, JR, Armstrong, D, and Phillips, SM. Consumption of fluid skim milk promotes greater muscle protein accretion after resistance exercise than does consumption of an isonitrogenous and isoenergetic soy-protein beverage. *Am. J. Clin. Nutr.* 85: 1031-1040, 2007.
 150. Witard, OC, Jackman, SR, Breen, L, Smith, K, Selby, A, and Tipton, KD. Myofibrillar muscle protein synthesis rates subsequent to a meal in response to increasing doses of whey protein at rest and after resistance exercise. *Am. J. Clin. Nutr.* 99: 86-95, 2014.
 151. Wojtaszewski, JF, MacDonald, C, Nielsen, JN, Hellsten, Y, Hardie, DG, Kemp, BE, Kiens, B, and Richter, EA. Regulation of 5'AMP-activated protein kinase activity and substrate utilization in exercising human skeletal muscle. *Am. J. Physiol. Endocrinol. Metab.* 284: E813-22, 2003.
 152. Wooding, DJ, Packer, JE, Kato, H, West, DWD, Courtney-Martin, G, Pencharz, PB, and Moore, DR. Increased Protein Requirements in Female Athletes after Variable-Intensity Exercise. *Med. Sci. Sports Exerc.* 49: 2297-2304, 2017.
 153. Wu, G. Amino acids: metabolism, functions, and nutrition. *Amino Acids* 37: 1-17, 2009.
 154. Yamada, S, Buffinger, N, DiMario, J, and Strohman, RC. Fibroblast growth factor is stored in fiber extracellular matrix and plays a role in regulating muscle hypertrophy. *Med. Sci. Sports Exerc.* 21: 173-180, 1989.
 155. Yang, Y, Breen, L, Burd, NA, Hector, AJ, Churchward-Venne, TA, Josse, AR, Tarnopolsky, MA, and Phillips, SM. Resistance exercise enhances myofibrillar protein synthesis with graded intakes of whey protein in older men. *Br. J. Nutr.* : 1-9, 2012.
 156. Young, K, and Davies, CT. Effect of diet on human muscle weakness following prolonged exercise. *Eur. J. Appl. Physiol. Occup. Physiol.* 53: 81-85, 1984.