

## REFERENCES

- Abildgaard, J. *et al.* (2013), 'Menopause is associated with decreased whole body fat oxidation during exercise', *American Journal of Physiology. Endocrinology and Metabolism*, 304(11), pp. E1227-1236. doi: 10.1152/ajpendo.00492.2012.
- Aceto, C. (1997), *Everything You Need to Know about Fat Loss*, (Adamsville TN: Fundco).
- Acheson, K. J. *et al.* (1988), 'Glycogen storage capacity and de novo lipogenesis during massive carbohydrate overfeeding in man', *The American Journal of Clinical Nutrition*, 48(2), pp. 240-247. doi: 10.1093/ajcn/48.2.240.
- Achten J. *et al.* (2004), 'Higher dietary carbohydrate content during intensified running training results in better maintenance of performance and mood state', *J. Appl. Physiol.*, vol. 96(4), pp. 1331-40.
- Ackerman, K. E. *et al.* (2019), 'Low energy availability surrogates correlate with health and performance consequences of Relative Energy Deficiency in Sport', *British Journal of Sports Medicine*, 53(10), pp. 628-633. doi: 10.1136/bjsports-2017-098958.
- Ackland T. *et al.* (2012), 'Current Assessment of Body Composition in Sport'. *Sports Med.*, vol. 42(3): 227-49.
- ACSM (1982), 'American College of Sports Medicine Position Statement on: The Use of Alcohol in Sports', *Medicine & Science in Sports & Exercise*, 14(6), p. ix.
- ACSM (1996), 'Position stand on exercise and fluid replacement', *Med. Sci. Sports and Ex.*, vol. 28, pp. i-vii.
- ACSM (2007), Armstrong, L. E., Casa, D. J. *et al.*, 'American College of Sports Medicine position stand on: Exertional heat illness during training and competition', *Med. Sci. Sports and Ex.*, vol. 39(3): 556-72.
- Adams, R. B., Egbo, K. N., and Demmig-Adams, B. (2014), 'High-dose vitamin C supplements diminish the benefits of exercise in athletic training and disease prevention', *Nutrition & Food Science*, vol. 44(2), pp. 95-101
- Ahlborg, B. *et al.* (1967), 'Human muscle glycogen content and capacity for prolonged exercise after different diets', *Forsvarsmedicin*, vol. 3, pp. 85-99.
- Ahnen, R. T., Jonnalagadda, S. S. and Slavin, J. L. (2019), 'Role of plant protein in nutrition, wellness, and health', *Nutrition Reviews*, 77(11), pp. 735-747. doi: 10.1093/nutrit/nuz028.
- Aird, T. P., Davies, R. W. and Carson, B. P. (2018), 'Effects of fasted vs fed-state exercise on performance and post-exercise metabolism: A systematic review and meta-analysis', *Scandinavian Journal of Medicine & Science in Sports*, 28(5), pp. 1476-1493. doi: 10.1111/sms.13054.
- Alexander, D. D. *et al.* (2016), 'Dairy consumption and CVD: a systematic review and meta-analysis', *The British Journal of Nutrition*, 115(4), pp. 737-750. doi: 10.1017/S0007114515005000.

- Alexander, D., Ball, M. J. and Mann, J. (1994), 'Nutrient intake and hematological status of vegetarians and age-sex matched omnivores', *Eur. J. Clin. Nutr.*, vol. 48, pp. 538–46.
- Alghannam, A. F. *et al.* (2016), 'Impact of muscle glycogen availability on the capacity for repeated exercise in man', *Med. Sci. Sports and Ex.*, vol. 48(1), pp. 123–31.
- Alghannam, A. F., Gonzalez, J. T. and Betts, J. A. (2018), 'Restoration of Muscle Glycogen and Functional Capacity: Role of Post-Exercise Carbohydrate and Protein Co-Ingestion', *Nutrients*, 10(2). doi: 10.3390/nu10020253.
- Alghannam, A. F., Gonzalez, J. T. and Betts, J. A. (2018), 'Restoration of Muscle Glycogen and Functional Capacity: Role of Post-Exercise Carbohydrate and Protein Co-Ingestion', *Nutrients*, 10(2), p. E253. doi: 10.3390/nu10020253.
- Allaway, H. C. M., Southmayd, E. A. and Souza, M. J. D. (2016), 'The physiology of functional hypothalamic amenorrhea associated with energy deficiency in exercising women and in women with anorexia nervosa', *Hormone Molecular Biology and Clinical Investigation*, 25(2), pp. 91–119. doi: 10.1515/hmbci-2015-0053.
- Allen, J. M. *et al.* (2018), 'Exercise Alters Gut Microbiota Composition and Function in Lean and Obese Humans', *Medicine and Science in Sports and Exercise*, 50(4), pp. 747–757. doi: 10.1249/MSS.0000000000001495.
- Alvares, T. S. *et al.* (2012), 'Acute L-Arginine supplementation does not increase nitric oxide production in healthy subjects', *Nutrition & Metabolism*, 9(1), p. 54. doi: 10.1186/1743-7075-9-54.
- Anderson, L. *et al.* (2017), 'Energy Intake and Expenditure of Professional Soccer Players of the English Premier League: Evidence of Carbohydrate Periodization', *International Journal of Sport Nutrition and Exercise Metabolism*, 27(3), pp. 228–238. doi: 10.1123/ijsnem.2016-0259.
- Anthony, J. C. *et al.* (2001), 'Signaling pathways involved in translational control of protein synthesis in skeletal muscle by leucine', *J. Nutr.*, vol. 131(3), pp. 856S–860S.
- Antoni, R. *et al.* (2018), 'A pilot feasibility study exploring the effects of a moderate time-restricted feeding intervention on energy intake, adiposity and metabolic physiology in free-living human subjects', *Journal of Nutritional Science*, 7. doi: 10.1017/jns.2018.13.
- Antonio, J. and Street, C. (1999), 'Glutamine: a potentially useful supplement for athletes', *Can. J. Appl. Physiol.*, vol. 24(1): S69–77.
- Appleby, P. N. *et al.* (1999), 'The Oxford Vegetarian Study: an overview', *Am. J. Clin. Nutr.*, vol. 70 (3 Suppl), pp. 525S–31S.
- Aragon, A. A. *et al.* (2017), 'International society of sports nutrition position stand: diets and body composition', *Journal of the International Society of Sports Nutrition*, 14(1), p. 16. doi: 10.1186/s12970-017-0174-y.
- Areces, F. *et al.* (2014), 'A 7-day oral supplementation with branched-chain amino acids was ineffective to prevent muscle damage during a marathon', *Amino Acids*, 46(5), pp. 1169–1176. doi: 10.1007/s00726-014-1677-3.
- Areta J. L. *et al.* (2013), 'Timing and distribution of protein ingestion during prolonged recovery from resistance exercise alters myofibrillar protein synthesis', *J. Physiol.*, vol. 591(9), pp. 2319–31.
- Armento, A. *et al.* (2021), 'Presence and Perceptions of Menstrual Dysfunction and Associated Quality of Life Measures Among High School Female Athletes', *Journal of Athletic Training*. doi: 10.4085/624-20.
- Armstrong, L. E. (2002), 'Caffeine, body fluid-electrolyte balance and exercise performance', *Int. J. Sport Nutr.*, vol. 12, pp. 189–206.
- Armstrong, L. E. *et al.* (1985), 'Influence of diuretic induced dehydration on competitive

- running performance', *Med. Sci. Sports Ex.*, vol. 17, pp. 456–61.
- Armstrong, L. E. *et al.* (1998), 'Urinary indices during dehydration, exercise and rehydration', *Int. J. Sport Nutr.*, vol. 8, pp. 345–55.
- Armstrong, L. E. *et al.* (2005), 'Fluid, electrolyte and renal indices of hydration during 11 days of controlled caffeine consumption', *Int. J. Sport Nutr. Exerc. Metab.*, vol. 15, pp. 252–65.
- Ashtary-Larky, D. *et al.* (2021), 'Ketogenic diets, physical activity, and body composition: A review', *The British Journal of Nutrition*, pp. 1–68. doi: 10.1017/S0007114521002609.
- Ashwell, M. *et al.* (2011), 'Waist-to-height ratio is a better screening tool than waist circumference and BMI for adult cardiometabolic risk factors: systematic review and meta-analysis'. *Obes. Rev.* vol 13(3), pp. 275–86.
- Asnicar, F. *et al.* (2021), 'Microbiome connections with host metabolism and habitual diet from 1,098 deeply phenotyped individuals', *Nature Medicine*, 27(2), pp. 321–332. doi: 10.1038/s41591-020-01183-8.
- Astrup, A. *et al.* (2019), 'WHO draft guidelines on dietary saturated and trans fatty acids: time for a new approach?', *BMJ*, 366, p. l4137. doi: 10.1136/bmj.l4137.
- Astrup, A., *et al.* (2010), 'The role of reducing intakes of saturated fat in the prevention of cardiovascular disease: where does the evidence stand in 2010?', *Am. J. Clin. Nutr.*, vol. 93, pp. 684–8.
- Aykan, N. F. (2015), 'Red Meat and Colorectal Cancer', *Oncology Reviews*, 9(1), p. 288. doi: 10.4081/oncol.2015.288.
- Bacon, L. *et al.* (2005), 'Size acceptance and intuitive eating improve health for obese, female chronic dieters', *J. Am. Diet. Assoc.*, vol. 105 (6), pp. 929–36.
- Barnes, M. J. (2014), 'Alcohol: impact on sports performance and recovery in male athletes', *Sports Medicine (Auckland, N.Z.)*, 44(7), pp. 909–919. doi: 10.1007/s40279-014-0192-8.
- Barr, S. I. and Costill, D. L. (1989), 'Water. Can the endurance athlete get too much of a good thing?', *J. Am. Diet. Assoc.*, vol. 89, pp. 1629–32.
- Barr, S. I. and Rideout, C. A. (2004), 'Nutritional considerations for vegetarian athletes', *Nutrition*, vol. 20 (7–8), pp. 696–703.
- Barrett, E. C., McBurney, M. I. and Ciappio, E. D. (2014), 'ω-3 fatty acid supplementation as a potential therapeutic aid for the recovery from mild traumatic brain injury/concussion', *Advances in Nutrition (Bethesda, Md.)*, 5(3), pp. 268–277. doi: 10.3945/an.113.005280.
- Bartlett, J. D. *et al.* (2013), 'Reduced carbohydrate availability enhances exercise-induced p53 signaling in human skeletal muscle: implications for mitochondrial biogenesis', *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*, 304(6), pp. R450–458. doi: 10.1152/ajpregu.00498.2012.
- Bartlett, J. D. *et al.* (2015), 'Carbohydrate availability and exercise training adaptation: too much of a good thing?', *Eur. J. Sport Sci.*, vol. 15(1), pp. 3–12.
- Barton, W. *et al.* (2018), 'The microbiome of professional athletes differs from that of more sedentary subjects in composition and particularly at the functional metabolic level', *Gut*, 67(4), pp. 625–633. doi: 10.1136/gutjnl-2016-313627.
- Bates *et al.* (2014), National Diet and Nutrition Survey. Headline results from years 1, 2, 3, 4 of the Rolling Programme (2208/9–2011/12) (Public Health England, London).
- Bauer, J. *et al.* (2013), 'Evidence-based recommendations for optimal dietary protein intake in older people: a position paper from the PROT-AGE Study Group', *Journal of the American Medical Directors Association*, 14(8), pp. 542–559. doi: 10.1016/j.jamda.2013.05.021.
- Bazzare, T. L. *et al.* (1986), 'Incidence of poor nutritional status among triathletes, endurance athletes and controls', *Med. Sci. Sports Ex.*, vol. 18, p. 590.
- BDA, *Vegetarian, vegan and plant-based diet.* Available at:

<https://www.bda.uk.com/resource/vegetarian-vegan-plant-based-diet.html> (Accessed: 6 May 2021).

- Beals, K. A. and Hill, A. K. (2006), 'The prevalence of disordered eating, menstrual dysfunction and low bone mineral density among US collegiate athletes', *Int. J. Sports Nutr. Exerc. Metab.*, vol. 16, pp. 1–23.
- Beelen, M. *et al.* (2008), 'Protein coingestion stimulates muscle protein synthesis during resistance type exercise', *Am. J. Physiol. Endocrinol Metab.*, vol. 295, pp. E70–7.
- Beelen, M. *et al.* (2010), 'Nutritional strategies to promote postexercise recovery', *Int. J. Sport Nutr. Exerc. Metab.*, vol. 20, pp. S15–32.
- Beis, L. Y. *et al.* (2011), 'Food and macronutrient intake of elite Ethiopian distance runners', *J. Int. Soc. Sports Nutr.*, vol. 8 pp. 7–11.
- Bell, K. E. *et al.* (2016), 'Muscle Disuse as a Pivotal Problem in Sarcopenia-related Muscle Loss and Dysfunction', *The Journal of Frailty & Aging*, 5(1), pp. 33–41. doi: 10.14283/jfa.2016.78.
- Bell, P. G. *et al.* (2015), 'Recovery facilitation with Montmorency cherries following high-intensity, metabolically challenging exercise', *Appl. Physiol. Nutr. Metab.*, vol. 40, pp. 414–23.
- Below, P. R. *et al.* (1995), 'Fluid and carbohydrate ingestion independently improve performance during one hour of intense exercise', *Med. Sci. Sports Exerc.*, vol. 27, pp. 200–10.
- Belza, A., Ritz, C., Sørensen, M. Q. *et al.* (2013), 'Contribution of gastroenteropancreatic appetite hormones to protein-induced satiety', *Am. J. Clin. Nutr.*, vol. 97(5), pp. 980–9.
- Bennell K. L. *et al.* (1995), 'Risk factors for stress fractures in female track and field athletes: a retrospective analysis'. *Clin. J. Sports Med.*, vol. 5, pp. 229–35.
- Beradi, J. M. *et al.* (2008), 'Recovery from a cycling time trial is enhanced with carbohydrate-protein supplementation vs. isoenergetic carbohydrate supplementation', *J. Int. Soc. Sports Nutr.* 2008, vol. 5, p. 24.
- Berg, A. and Keul, J. (1988), 'Biomechanical changes during exercise in children', *Young athletes: Biological, psychological and educational perspectives*, ed. R. M. Malina (Champaign, IL, Human Kinetics), pp. 61–77.
- Bergstrom, J. *et al.* (1967), 'Diet, muscle glycogen and physical performance', *Acta. Physiol. Scand.*, vol. 71, pp. 140–50.
- Berkulo, M. A. R. *et al.* (2016), 'Ad-libitum drinking and performance during a 40-km cycling time trial in the heat', *European Journal of Sport Science*, 16(2), pp. 213–220. doi: 10.1080/17461391.2015.1009495.
- Bermon, S. *et al.* (2017), 'Consensus Statement Immunonutrition and Exercise', *Exercise Immunology Review*, 23, pp. 8–50.
- Berrazaga, Insaf, *et al.* (2019), "The Role of the Anabolic Properties of Plant- versus Animal-Based Protein Sources in Supporting Muscle Mass Maintenance: A Critical Review.", *Nutrients*, vol. 11, no. 8, Aug. 2019. PubMed Central, doi:10.3390/nu11081825.
- Berry, D. J. *et al.* (2011), 'Vitamin D status has a linear association with seasonal infections and lung function in British adults', *British Journal of Nutrition*, 106(9), pp. 1433–1440. doi: 10.1017/S0007114511001991.
- Bescos, R. *et al.* (2012), 'The effect of nitric oxide related supplements on human performance', *Sports Med.*, vol. 42(2), pp. 99–117.
- Betts, J. A. and Williams, C. (2010), 'Short-term recovery from prolonged exercise: exploring the potential for protein ingestion to accentuate the benefits of carbohydrate supplements', *Sports Medicine (Auckland, N.Z.)*, 40(11), pp. 941–959. doi: 10.2165/11536900-000000000-00000.
- Betts, J. A. *et al.* (2016), 'Is breakfast the most important meal of the day?', *The Proceedings of the*

- Nutrition Society*, 75(4), pp. 464–474. doi: 10.1017/S0029665116000318.
- Betts, J. *et al.* (2007), ‘The influence of carbohydrate and protein ingestion during recovery from prolonged exercise on subsequent endurance performance’, *J. Sports Sci.*, vol. 25(13), pp. 1449–60.
- Betts, J. *et al.* (2014), ‘The causal role of breakfast in energy balance and health: a randomized controlled trial in lean adults’, *Am. J. Clin. Nutr.*, vol. 100(2), pp. 539–47.
- Biolo, G. *et al.* (2008), ‘Positive energy balance is associated with accelerated muscle atrophy and increased erythrocyte glutathione turnover during 5 wk of bed rest’, *The American Journal of Clinical Nutrition*, 88(4), pp. 950–958. doi: 10.1093/ajcn/88.4.950.
- Biolo, G., K. D. Tipton, S. Klein, and R. R. Wolfe (1997), ‘An abundant supply of amino acids enhances the metabolic effect of exercise on muscle protein’, *Am. J. Physiol.* 273: E122-E129.
- Bishop, N. C. *et al.* (2001), ‘Pre-exercise carbohydrate status and immune responses to prolonged cycling: I. Effect on neutrophil degranulation’, *International Journal of Sport Nutrition and Exercise Metabolism*, 11(4), pp. 490–502. doi: 10.1123/ijsnem.11.4.490.
- Bishop, N. C. *et al.* (2002), ‘Influence of carbohydrate supplementation on plasma cytokine and neutrophil degranulation responses to high intensity intermittent exercise’, *Int. J. Sport Nutr.*, vol. 12, pp. 145–56.
- Bloomer, R. J. *et al.* (2000), ‘Effects of meal form and composition on plasma testosterone, cortisol and insulin following resistance exercise’, *Int. J. Sport Nutr.*, vol. 10, pp. 415–24.
- Bodine, S. C. *et al.* (2001), ‘Akt/mTOR pathway is a crucial regulator of skeletal muscle hypertrophy and can prevent muscle atrophy in vivo’, *Nature Cell Biology*, 3(11), pp. 1014–1019. doi: 10.1038/ncb1101-1014.
- Boirie, Y. *et al.* (1997), ‘Slow and fast dietary proteins differently modulate postprandial protein accretion’, *Proc. Nat. Acad. Sci. USA.*, vol. 94(26), pp. 14930–5.
- Bonn-Miller, M. O. *et al.* (2017), ‘Labeling Accuracy of Cannabidiol Extracts Sold Online’, *JAMA*, 318(17), p. 1708. doi: 10.1001/jama.2017.11909.
- Booth, F. and Zwetsloot, K. (2010), ‘Basic concepts about genes, inactivity and aging’, *Scand. J. Med. Sci. Sports*, vol. 20, pp. 1–4.
- Borsheim E *et al.* (2004), ‘Effect of an amino acid, protein, and carbohydrate mixture on net muscle protein balance after resistance training’, *Int. J. Sport. Nutr. Exerc. Metab.*, vol. 14, pp. 255–71.
- Bosch, A. N. *et al.* (1994), ‘Influence of carbohydrate ingestion on fuel substrate turnover and oxidation during prolonged exercise’, *J. Appl. Physiol.*, vol. 76, pp. 2364–72.
- Bounous, G. and Gold, P. (1991), ‘The biological activity of un-denatured whey proteins: role of glutathione’, *Clin. Invest. Med.*, vol. 4, pp. 296–309.
- Boutcher, Y. N. *et al.* (2019), ‘The Effect of Sprint Interval Training on Body Composition of Postmenopausal Women’, *Medicine & Science in Sports & Exercise*, 51(7), pp. 1413–1419. doi: 10.1249/MSS.0000000000001919.
- Bouvard, V. *et al.* (2015), ‘Carcinogenicity of consumption of red and processed meat’, *The Lancet Oncology*, Oct. 26 2015.
- Bowtell, J. and Kelly, V. (2019), ‘Fruit-Derived Polyphenol Supplementation for Athlete Recovery and Performance’, *Sports Medicine*, 49(1), pp. 3–23. doi: 10.1007/s40279-018-0998-x.
- Bowtell, J. L., Sumners, D. P., Dyer, A. *et al.* (2011), ‘Montmorency cherry juice reduces muscle damage caused by intensive strength exercise’, *Medicine and Science in Sports Exercise*, vol. 43(8), pp. 1544–51.
- Brand-Miller, J. *et al.* (2003), ‘Low GI diet in the management of diabetes’. *Diabetes Care*, vol. 26, pp. 2261–7.

- Brand-Miller, J., Foster-Powell, K. and McMillan Price J. (2005), *The Low GI Diet*. (Hodder Mobius).
- Breen, L. and Phillips, S. M. (2013), 'Interactions between exercise and nutrition to prevent muscle waste during ageing', *British Journal of Clinical Pharmacology*, 75(3), pp. 708–715. doi: 10.1111/j.1365-2125.2012.04456.x.
- Brilla, L. R. and Conte, V. (2000), 'Effects of a novel zinc-magnesium formulation on hormones and strength', *J. Exerc. Physiol. Online*, vol. 3(4), pp. 1–15.
- Brinkworth, G. D. and Buckley, J. D. (2003), 'Concentrated bovine colostrum protein supplementation reduces the incidence of self-reported symptoms of upper respiratory tract infection in adult males', *Eur. J. Nutr.*, vol. 42, pp. 228–32.
- Brinkworth, G. D. *et al.* (2004), 'Effect of bovine colostrum supplementation on the composition of resistance trained and untrained limbs in healthy young men', *Eur. J. Appl. Physiol.*, vol. 91, pp. 53–60.
- Broeder, C. E. *et al.* (2000), 'The Andro Project', *Arch. Intern. Med.*, vol. 160(20), pp. 3093–104.
- Brouns, F. *et al.* (1998), 'The effect of different rehydration drinks on post-exercise electrolyte excretion in trained athletes', *Int. J. Sports Med.*, vol. 19, pp. 56–60.
- Brown, E. C. *et al.* (2004), 'Soya versus whey protein bars: Effects of exercise training impact on lean body mass and antioxidant status', *J. Nutr.*, vol. 3, pp. 22–7.
- Brown, G. A. *et al.* (2000), 'Effects of anabolic precursors on serum testosterone concentrations and adaptations to resistance training in young men', *Int. J. Sport Nutr.*, vol. 10, pp. 340–59.
- Brown, M. A., Stevenson, E. J. and Howatson, G. (2019), 'Montmorency tart cherry (*Prunus cerasus* L.) supplementation accelerates recovery from exercise-induced muscle damage in females', *European Journal of Sport Science*, 19(1), pp. 95–102. doi: 10.1080/17461391.2018.1502360.
- Brown, R. C. and Cox, C. M. (1998), 'Effects of high fat versus high carbohydrate diets on plasma lipids and lipoproteins in endurance athletes', *Medicine & Science in Sports & Exercise*, 30(12), pp. 1677–1683.
- Brownlie, T., *et al.* (2004), 'Tissue iron deficiency without anemia impairs adaptation in endurance capacity after aerobic training in previously untrained women', *Am. J. Clin. Nutr.*, vol. 79(3), pp. 437–43.
- Bryer S. C. and Goldfarb, A. H. (2006), 'Effect of high dose vitamin C supplementation on muscle soreness, damage, function and oxidative stress to eccentric exercise', *Int. J. Sport Nutr. Exerc. Metab.*, vol. 16, pp. 270–80.
- Buckley, J. D. *et al.* (2003), 'Effect of bovine colostrum on anaerobic exercise performance and plasma insulin-like growth factor', *J. Sports Sci.*, vol. 21, pp. 577–88.
- Buford, T. W., *et al.* (2007), 'International Society of Sports Nutrition position stand: creatine supplementation and exercise', *J. Int. Soc. Sports Nutr.*, vol. 4, p. 6.
- Burd, N. A. *et al.* (2009), 'Exercise training and protein metabolism: influences of contraction, protein intake, and sex-based differences', *J. Appl. Physiol.*, vol. 106(5), pp. 1692–701.
- Burd, N. A. *et al.* (2011), 'Enhanced amino acid sensitivity of myofibrillar protein synthesis persists for up to 24 h after resistance exercise in young men', *J. Nutr.*, vol. 141(4), pp. 568–73.
- Burd, N. A. *et al.* (2013), 'Anabolic resistance of muscle protein synthesis with aging', *Exerc. Sport Sci. Rev.*, vol. 41, pp. 169–73.
- Burd, N. A. *et al.* (2015), 'Differences in postprandial protein handling after beef compared with milk ingestion during postexercise recovery: a randomized controlled trial', *The American Journal of Clinical Nutrition*, 102(4), pp. 828–836. doi: 10.3945/ajcn.114.103184.

- Burd, N. A. *et al.* (2019), 'Food-First Approach to Enhance the Regulation of Post-exercise Skeletal Muscle Protein Synthesis and Remodeling', *Sports Medicine (Auckland, N.z.)*, 49(Suppl 1), pp. 59–68. doi: 10.1007/s40279-018-1009-y.
- Burke L. M. *et al.* (2011), 'Carbohydrates for training and competition', *J. Sports Sci.*, vol. 29, Suppl. 1, pp. S17–27.
- Burke, D. G. *et al.* (1993), 'Muscle glycogen storage after prolonged exercise: effect of glycaemic index of carbohydrate feedings', *J. Appl. Physiol.*, vol. 75, pp. 1019–23.
- Burke, D. G. *et al.* (2000), 'The effect of continuous low dose creatine supplementation on force, power and total work', *Int. J. Sport Nutr.*, vol. 10, pp. 235–44.
- Burke, L. M. (2010), 'Fueling strategies to optimize performance: training high or training low?', *Scand. J. Med. Sci. Sports*, vol. 20, Suppl. 2, pp. 48–58.
- Burke, L. M. (2021), 'Ketogenic low-CHO, high-fat diet: the future of elite endurance sport?', *The Journal of Physiology*, 599(3), pp. 819–843. doi: 10.1113/JP278928.
- Burke, L. M. and Kiens, B. (2006), "'Fat adaptation" for athletic performance: the nail in the coffin?', *Journal of Applied Physiology (Bethesda, Md.: 1985)*, 100(1), pp. 7–8. doi: 10.1152/jappphysiol.01238.2005.
- Burke, L. M. and Kiens, B. (2006), "'Fat adaptation" for athletic performance: the nail in the coffin?', *Journal of Applied Physiology (Bethesda, Md.: 1985)*, 100(1), pp. 7–8. doi: 10.1152/jappphysiol.01238.2005.
- Burke, L. M. *et al.* (2003), 'Effect of alcohol intake on muscle glycogen storage after prolonged exercise', *Journal of Applied Physiology (Bethesda, Md.: 1985)*, 95(3), pp. 983–990. doi: 10.1152/jappphysiol.00115.2003.
- Burke, L. M. *et al.* (2004), 'Carbohydrates and fat for training and recovery', *J. Sports Sci.*, vol. 22(1), pp. 15–30.
- Burke, L. M. *et al.* (2012), 'Effect of intake of different dietary protein sources on plasma amino acid profiles at rest and after exercise', *Int. J. Sport Nutr. and Exerc. Metab.*, vol. 22, pp. 452–62.
- Burke, L. M. *et al.* (2017), 'Low carbohydrate, high fat diet impairs exercise economy and negates the performance benefit from intensified training in elite race walkers', *The Journal of Physiology*, 595(9), pp. 2785–2807. doi: 10.1113/JP273230.
- Burke, L. M. *et al.* (2018), 'Relative Energy Deficiency in Sport in Male Athletes: A Commentary on Its Presentation Among Selected Groups of Male Athletes', *International Journal of Sport Nutrition and Exercise Metabolism*, 28(4), pp. 364–374. doi: 10.1123/ijsnem.2018-0182.
- Burke, L. M. *et al.* (2019), 'International Association of Athletics Federations Consensus Statement 2019: Nutrition for Athletics', *International Journal of Sport Nutrition and Exercise Metabolism*, 29(2), pp. 73–84. doi: 10.1123/ijsnem.2019-0065.
- Burke, L. M. *et al.* (2020), 'Crisis of confidence averted: Impairment of exercise economy and performance in elite race walkers by ketogenic low carbohydrate, high fat (LCHF) diet is reproducible', *PloS One*, 15(6), p. e0234027. doi: 10.1371/journal.pone.0234027.
- Bussau, V. A. *et al.* (2002), 'Carbohydrate loading in human muscle: an improved 1-day protocol', *Eur. J. Appl. Physiol.*, vol. 87, pp. 290–5.
- Butterfield G. E. (1996), 'Ergogenic Aids: Evaluating sport nutrition products', *Int. Sport Nutr.*, vol. 6, pp. 191–7.
- Byrne, S. and McLean, N. (2002), 'Elite athletes: effects of the pressure to be thin', *Journal of Science and Medicine in Sport*, 5(2), pp. 80–94. doi: 10.1016/s1440-2440(02)80029-9.
- Cahill, C. F. (1976), 'Starvation in Man', *J. Clin. Endocrinol. Metab.*, vol. 5, pp. 397–415.
- Calder, P. C. (2013), 'n-3 fatty acids, inflammation and immunity: new mechanisms to explain old actions', *The Proceedings of the Nutrition Society*, 72(3), pp. 326–336. doi:

10.1017/S0029665113001031.

- Calder, P. C. (2020), 'Nutrition, immunity and COVID-19', *BMJ nutrition, prevention & health*, 3(1), pp. 74–92. doi: 10.1136/bmjnph-2020-000085.
- Cameron S. L. *et al.* (2010), 'Increased blood pH but not performance with sodium bicarbonate supplementation in elite rugby union players', *Int. J. Sport Nutr. Exerc. Metab.*, vol. 20(4), pp. 307–21.
- Campbell, B. *et al.* (2013), 'International Society of Sports Nutrition position stand: energy drinks', *Journal of the International Society of Sports Nutrition*, 10(1), p. 1. doi: 10.1186/1550-2783-10-1.
- Campbell, W. W. *et al.* (1995), 'Effects of resistance training and dietary protein intake in protein metabolism in older adults', *Am. J. Physiol.*, Vol 268, pp. 1143–53.
- Candow, D. G. *et al.* (2001), 'Effect of glutamine supplementation combined with resistance training in young adults', *Eur. J. Appl. Physiol.*, vol. 86(2), pp. 142–9.
- Candow, D. G. *et al.* (2006), 'Effect of whey and soya protein supplementation combined with resistance training in young adults', *Int. J. Sports Nutr. Exerc. Metab.*, vol. 16, pp. 233–44.
- Candow, D. G., Chilibeck, P. D. and Forbes, S. C. (2014), 'Creatine supplementation and aging musculoskeletal health', *Endocrine*, 45(3), pp. 354–coyle361. doi: 10.1007/s12020-013-0070-4.
- Cann, C. E. *et al.* (1984), 'Decreased spinal mineral content in amenorrhoeic women', *JAMA*, vol. 251, pp. 626–9.
- Cannell, J. *et al.* (2009), 'Athletic performance and vitamin D', *Med. Sci. Sports Exerc.*, vol. 41, pp. 1102–10.
- Carbon, R. (2002), 'The female athlete triad does not exist', *Sports Care News*, 26, pp. 3–5.
- Carr, A. J. (2011), 'Effects of acute alkalosis and acidosis on performance: a meta-analysis', *Sports Med.*, vol. 41(10), pp. 801–14.
- Carswell, A. T. *et al.* (2018), 'Influence of Vitamin D Supplementation by Sunlight or Oral D3 on Exercise Performance', *Medicine and Science in Sports and Exercise*, 50(12), pp. 2555–2564. doi: 10.1249/MSS.0000000000001721.
- Carter, J. M. *et al.* (2004), 'The effect of carbohydrate mouth rinse on 1-h cycle time-trial performance', *Med. Sci. Sports Exerc.*, vol. 36, pp. 2107–11.
- Castell, L. M. and Newsholme, E. A. (1997), 'The effects of oral glutamine supplementation on athletes after prolonged exhaustive exercise', *Nutrition*, vol. 13, pp. 738–42.
- Castell, L. M., Poortmans, J. R. and Newsholme, E. A. (1996), 'Does glutamine have a role in reducing infections in athletes?', *European Journal of Applied Physiology and Occupational Physiology*, 73(5), pp. 488–490. doi: 10.1007/BF00334429.
- Cermak N. M. *et al.* (2012), 'Nitrate supplementation's improvement of 10 km time-trial performance in trained cyclists'. *Int. J. Sport Nutr. Exerc. Metab.*, vol. 1, pp. 64–71.
- Chaix, A. *et al.* (2014), 'Time-restricted feeding is a preventive and therapeutic intervention against diverse nutritional challenges', *Cell Metabolism*, 20(6), pp. 991–1005. doi: 10.1016/j.cmet.2014.11.001.
- Chen, Z. *et al.* (2005), 'Postmenopausal hormone therapy and body composition--a substudy of the estrogen plus progestin trial of the Women's Health Initiative', *The American Journal of Clinical Nutrition*, 82(3), pp. 651–656. doi: 10.1093/ajcn.82.3.651.
- Cheung, S. S. *et al.* (2015), 'Separate and combined effects of dehydration and thirst sensation on exercise performance in the heat'. *Scand. J. Med. Sci. Sports*, vol. 25, Suppl. 1, pp. 104–11.
- Chevront, S. N., Carter, R., and Sawka, M. N. (2003), 'Fluid balance and endurance exercise performance', *Current Sports Medicine Reports*, vol. 2 pp. 202–8.
- Chevront, S. N., Montain, S. J. and Sawka, M. N. (2007), 'Fluid replacement and performance



- during the marathon', *Sports Medicine (Auckland, N.Z.)*, 37(4–5), pp. 353–357. doi: 10.2165/00007256-200737040-00020.
- Chowdhury, E. A. *et al.* (2015), 'Carbohydrate-rich breakfast attenuates glycaemic, insulinaemic and ghrelin response to *ad libitum* lunch relative to morning fasting in lean adults'. *Br. J. Nutr.*, vol. 114(1), pp. 98–107.
- Chowdhury, R. *et al.* (2014), 'Association of dietary, circulating, and supplement fatty acids with coronary risk: A systematic review and meta-analysis', *Ann. Intern. Med.*, vol. 160(6), pp. 398–406.
- Christensen, E. H. and Hansen, O. (1939), 'Arbeitsfähigkeit und Ernährung', *Skand. Arch. Physiol.*, vol. 81, pp. 160–71.
- Chryssanthopoulos, C. *et al.* (2002), 'The effect of a high carbohydrate meal on endurance running capacity', *Int. J. Sport Nutr.*, vol. 12, pp. 157–71.
- Churchward-Venne, T. A. *et al.* (2012), 'Supplementation of a suboptimal protein dose with leucine or essential amino acids: effects on myofibrillar protein synthesis at rest and following resistance exercise in men', *The Journal of Physiology*, 590(11), pp. 2751–2765. doi: 10.1113/jphysiol.2012.228833.
- Churchward-Venne, T. A. *et al.* (2020), 'Dose-response effects of dietary protein on muscle protein synthesis during recovery from endurance exercise in young men: a double-blind randomized trial', *The American Journal of Clinical Nutrition*, 112(2), pp. 303–317. doi: 10.1093/ajcn/nqaa073.
- Churuangsuk, C. *et al.* (2018), 'Low-carbohydrate diets for overweight and obesity: a systematic review of the systematic reviews', *Obesity Reviews: An Official Journal of the International Association for the Study of Obesity*, 19(12), pp. 1700–1718. doi: 10.1111/obr.12744.
- Ciuris, C. *et al.* (2019), 'A Comparison of Dietary Protein Digestibility, Based on DIAAS Scoring, in Vegetarian and Non-Vegetarian Athletes', *Nutrients*, 11(12). doi: 10.3390/nu11123016.
- Clark, J. F. (1997), 'Creatine and phosphocreatine: a review', *J. Athletic Training*, vol. 32(1), pp. 45–50.
- Clark, N. (1995), 'Nutrition quackery: when claims are too good to be true', *Phys. Sports Med.*, vol. 23, pp. 7–8.
- Clayton D. J. *et al.* (2015), 'Effect of breakfast omission on energy intake and evening exercise performance', *Med. Sci. Sports Exerc.*, vol. 47(12), pp. 2645–52.
- Clayton, D. J. and James, L. (2015), 'The effect of breakfast on appetite regulation, energy balance and exercise performance', *Proc. Nutr. Soc.*, vol. 14, pp. 1–9 (Epub ahead of print).
- Close, G. L., Cobley, R. J., Owens, D. J. *et al.* (2013a), 'Assessment of vitamin D concentration in non-supplemented professional athletes and healthy adults during the winter months in the UK: implications for skeletal muscle function', *J. Sports Sci.*, vol. 31(4), pp. 344–53.
- Close, G. L., *et al.* (2013b), 'The effects of vitamin D3 supplementation on serum total 25(OH) D concentration and physical performance: a randomised dose-response study', *Br. J. Sports Med.*, vol. 47, pp. 692–6.
- Close, G. L., Kasper, A. M. and Morton, J. P. (2019), 'From Paper to Podium: Quantifying the Translational Potential of Performance Nutrition Research', *Sports Medicine*, 49(1), pp. 25–37. doi: 10.1007/s40279-018-1005-2.
- Cobb, K. L. *et al.* (2003), 'Disordered eating, menstrual irregularity and bone mineral density in female runners', *Med. Sci. Sports Exerc.*, vol. 35, pp. 711–19.
- Cochran, A. J. R. *et al.* (2010), 'Carbohydrate feeding during recovery alters the skeletal muscle metabolic response to repeated sessions of high-intensity interval exercise in humans', *Journal of Applied Physiology (Bethesda, Md.: 1985)*, 108(3), pp. 628–636. doi:

10.1152/jappphysiol.00659.2009.

- Cockburn, E. *et al.* (2008), 'Acute milk-based protein-CHO supplementation attenuates exercise-induced muscle damage', *Appl. Physiol. Nutr. Metab.*, Aug; 33(4): 775–83.
- Cockburn, E., Robson-Ansley, P., Hayes, P. R., Stevenson, E. (2012), 'Effect of volume of milk consumed on the attenuation of exercise-induced muscle damage', *Eur. J. Appl. Physiol.*, Jan 7. (Epub ahead of print.)
- Coggan, A. R. and Coyle, E. F. (1987), 'Reversal of fatigue during prolonged exercise by carbohydrate infusion or ingestion'. *J. Appl. Physiol.*, vol. 63, pp. 2388–95.
- Coggan, A. R. and Coyle, E. F. (1991), 'Carbohydrate ingestion during prolonged exercise: effects on metabolism and performance'. In J. Holloszy (ed.), *Exercise and Sports Science Reviews*, vol. 19 (Williams and Wilkins), pp. 1–40.
- Cole, T. J., Bellizzi, M., Flegal, K. and Dietz, W. H. (2000), 'Establishing a standard definition for child overweight and obesity worldwide: international survey'. *British Medical Journal*, vol. 320, pp. 1240–3.
- Collins, J. *et al.* (2021), 'UEFA expert group statement on nutrition in elite football. Current evidence to inform practical recommendations and guide future research', *British Journal of Sports Medicine*, 55(8), p. 416. doi: 10.1136/bjsports-2019-101961.
- Cook, M. D. *et al.* (2015), 'New Zealand blackcurrant extract improves cycling performance and fat oxidation in cyclists'. *Eur. J. Appl. Physiol.*, vol. 115(11), pp. 2357–65.
- Cooper, R. *et al.* (2012), 'Creatine supplementation with specific view to exercise/sports performance: an update'. *J. Int. Soc. Sports Nutr.*, vol. 9, p. 33.
- Corder, K. E. *et al.* (2016), 'Effects of short-term docosahexaenoic acid supplementation on markers of inflammation after eccentric strength exercise in women'. *J. Sports Sci. Med.*, vol 15, pp. 176–83.
- Costa, R. J. S. *et al.* (2017) (a) 'Systematic review: exercise-induced gastrointestinal syndrome-implications for health and intestinal disease', *Alimentary Pharmacology & Therapeutics*, 46(3), pp. 246–265. doi: 10.1111/apt.14157.
- Costa, Ricardo J. S. *et al.* (2017) (b), 'Gut-training: the impact of two weeks repetitive gut-challenge during exercise on gastrointestinal status, glucose availability, fuel kinetics, and running performance', *Applied Physiology, Nutrition, and Metabolism = Physiologie Appliquee, Nutrition Et Metabolisme*, 42(5), pp. 547–557. doi: 10.1139/apnm-2016-0453.
- Costill, D. L. (1985), 'Carbohydrate nutrition before, during and after exercise'. *Fed. Proc.*, vol. 44, pp. 364–368.
- Costill, D. L. (1986), *Inside Running: Basics of Sports Physiology* (Benchmark Press), p. 189.
- Costill, D. L. (1988), 'Carbohydrates for exercise: dietary demands for optimal performance'. *Int. J. Sports Med.*, vol. 9, pp. 1–18.
- Costill, D. L. and Hargreaves, M. (1992), 'Carbohydrate nutrition and fatigue'. *Sports Med.*, vol. 13, pp. 86–92.
- Costill, D. L. *et al.* (1971), 'Muscle glycogen utilisation during prolonged exercise on successive days'. *J. Appl. Physiol.*, vol. 31, pp. 834–8.
- Cox, A. J., Pyne, D. B., Saunders, P. U., and Fricker, P. A. (2008), 'Oral administration of the probiotic *Lactobacillus fermentum* VRI-003 and mucosal immunity in endurance athletes'. *Br. J. Sports Med.* (Epub Feb. 13).
- Cox, G. *et al.* (2002), 'Acute creatine supplementation and performance during a field test simulating match play in elite female soccer players'. *Int. J. Sport Nutr.*, vol. 12, pp. 33–46.
- Cox, G. R. *et al.* (2010), 'Daily training with high carbohydrate availability increases exogenous carbohydrate oxidation during endurance cycling', *Journal of Applied Physiology*, 109(1), pp. 126–134. doi: 10.1152/jappphysiol.00950.2009.

- Cox, P. J. *et al.* (2016), 'Nutritional Ketosis Alters Fuel Preference and Thereby Endurance Performance in Athletes', *Cell Metabolism*, 24(2), pp. 256–268. doi: 10.1016/j.cmet.2016.07.010.
- Coyle, E. (2004), 'Fluid and fuel intake during exercise'. *J. Sports Sci.*, vol. 22, pp. 39–55.
- Coyle, E. F. (1988), 'Carbohydrates and athletic performance'. *Sports Sci. Exch. Sports Nutr.*, Gatorade Sports Science Institute, vol. 1.
- Coyle, E. F. (1991), 'Timing and method of increased carbohydrate intake to cope with heavy training, competition and recovery'. *J. Sports Sci.*, vol. 9 (suppl.), pp. 29–52.
- Coyle, E. F. (1995), 'Substrate utilization during exercise in active people'. *Am. J. Clin. Nutr.*, vol. 61 (suppl), pp. 968–79.
- Coyle, E. F. and Coggan, A. R. (1984), 'Effectiveness of carbohydrate feeding in delaying fatigue during prolonged exercise', *Sports Medicine (Auckland, N.Z.)*, 1(6), pp. 446–458. doi: 10.2165/00007256-198401060-00004.
- Coyle, E. F. *et al.* (1986), 'Muscle glycogen utilization during prolonged strenuous exercise when fed carbohydrate', *Journal of Applied Physiology (Bethesda, Md.: 1985)*, 61(1), pp. 165–172. doi: 10.1152/jappl.1986.61.1.165.
- Craddock, J. *et al.* (2015), 'Vegetarian and omnivorous nutrition – comparing physical performance'. *Int. J. Sport Nutr. Exerc. Metab.*, Nov 16 (Epub ahead of print).
- Craig, W. J., Mangels, A. R.; American Dietetic Association (2009), Position of the American Dietetic Association: vegetarian diets'. *J. Am. Diet. Assoc.*, vol. 109(7), pp. 1266–82.
- Craven, J. *et al.* (2021), 'The Effect of Consuming Carbohydrate With and Without Protein on the Rate of Muscle Glycogen Re-synthesis During Short-Term Post-exercise Recovery: a Systematic Review and Meta-analysis', *Sports Medicine – Open*, 7(1), p. 9. doi: 10.1186/s40798-020-00297-0.
- Cribb P. J. *et al.* (2006), 'The effect of whey isolate and resistance training on strength, body composition and plasma glutamine'. *Int. J. Sports Nutr. Exerc. Metab.*, vol. 16, pp. 494–509.
- Cronise, R. J., Sinclair, D. A. and Bremer, A. A. (2017), 'Oxidative Priority, Meal Frequency, and the Energy Economy of Food and Activity: Implications for Longevity, Obesity, and Cardiometabolic Disease', *Metabolic Syndrome and Related Disorders*, 15(1), pp. 6–17. doi: 10.1089/met.2016.0108.
- Crooks, C. V. *et al.* (2006), 'The effect of bovine colostrum supplementation on salivary IgA in distance runners'. *Int. J. Sport Nutr. Exerc. Metab.*, vol. 16, pp. 47–64.
- Crowe, F. L. *et al.* (2013), 'Risk of hospitalization or death from ischemic heart disease among British vegetarians and nonvegetarians: results from the EPIC-Oxford cohort study', *The American Journal of Clinical Nutrition*, 97(3), pp. 597–603. doi: 10.3945/ajcn.112.044073.
- Crowe, M. J., Weatherson, J. N., Bowden, B. F. (2006), 'Effects of dietary leucine supplementation on exercise performance'. *Eur. J. Appl. Physiol.*, vol. 97(6), p. 664.
- Cupisti, A. *et al.* (2002), 'Nutrition knowledge and dietary composition in Italian female athletes and non-athletes'. *Int. J. Sport Nutr.*, vol. 12, pp. 207–19.
- Currell, K. and Jeukendrup, A. E. (2008), 'Superior endurance performance with ingestion of multiple transportable carbohydrates'. *Med. Sci. Sports Exerc.*, vol. 40(2), pp. 275–81.
- Dangin, M. *et al.* (2001), 'The digestion rate of protein is an independent regulating factor of postprandial protein retention'. *Am. Physiol. Soc. Abstracts*, vol. 7: 022E.
- Danz, M. *et al.* (2016), 'Hyponatremia among triathletes in the Ironman European Championship'. *N. Engl. J. Med.*, vol. 374, pp. 997–9.
- Daries, H. N., Noakes, T. D. and Dennis, S. C. (2000), 'Effect of fluid intake volume on 2-h running performances in a 25 degrees C environment', *Medicine and Science in Sports and Exercise*, 32(10), pp. 1783–1789. doi: 10.1097/00005768-200010000-00019.

- Darling, A. L. *et al.* (2019), 'Dietary protein and bone health across the life-course: an updated systematic review and meta-analysis over 40 years', *Osteoporosis international: a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*, 30(4), pp. 741–761. doi: 10.1007/s00198-019-04933-8.
- Davey, G. K. *et al.* (2003), 'EPIC-Oxford: Lifestyle characteristics and nutrient intakes in a cohort of 33,883 meat-eaters and 31,546 non-meat-eaters in the UK'. *Public Health Nutrition*, vol. 6(3), pp. 259–68.
- David, L. A. *et al.* (2014), 'Diet rapidly and reproducibly alters the human gut microbiome', *Nature*, 505(7484), pp. 559–563. doi: 10.1038/nature12820.
- Davis, C. (1993), 'Body image, dieting behaviours and personality factors: a study of high-performance female athletes'. *Int. J. Sport Psych.*, vol. 23, pp. 179–92.
- Davis, J. M. *et al.* (1988), 'Carbohydrate-electrolyte drinks: effects on endurance cycling in the heat'. *Am. J. Clin. Nutr.*, vol. 48, pp. 1023–30.
- Davis, S. R. *et al.* (2012), 'Understanding weight gain at menopause', *Climacteric*, 15(5), pp. 419–429. doi: 10.3109/13697137.2012.707385.
- Davison, G. (2012), 'Bovine colostrum and immune function after exercise'. *Med. Sport Sci.* vol. 59, pp. 62–9.
- de Ataide e Silva, T. *et al.* (2013), 'Can carbohydrate mouth rinse improve performance during exercise? A systematic review'. *Nutrients*, Dec 19, 6(1), pp. 1–10. doi: 10.3390/nu6010001.
- de Oliveira Otto, M. C., Mozaffarian, D., Kromhout, D. *et al.* (2012), 'Dietary intake of saturated fat by food source and incident cardio vascular disease: the multi-ethnic study of atherosclerosis'. *Am. J. of Clin. Nutr.*, vol. 96(2), pp. 397–404.
- de Oliveira, E. P. and Burini, R. C. (2014), 'Carbohydrate-dependent, exercise-induced gastrointestinal distress', *Nutrients*, 6(10), pp. 4191–4199. doi: 10.3390/nu6104191.
- de Oliveira, E. P., Burini, R. C. and Jeukendrup, A. (2014), 'Gastrointestinal Complaints During Exercise: Prevalence, Etiology, and Nutritional Recommendations', *Sports Medicine (Auckland, N.z.)*, 44(Suppl 1), pp. 79–85. doi: 10.1007/s40279-014-0153-2.
- De Souza, M. J. *et al.* (2007), 'Drive for thinness score is a proxy indicator of energy deficiency in exercising women', *Appetite*, 48(3), pp. 359–367. doi: 10.1016/j.appet.2006.10.009.
- De Souza, R. J. *et al.* (2015), 'Intake of saturated and trans unsaturated fatty acids and risk of all cause mortality, cardiovascular disease, and type 2 diabetes: systematic review and meta-analysis of observational studies'. *BMJ*, vol. 351, h3978.
- De Souza, R. J. *et al.* (2015), 'Intake of saturated and trans unsaturated fatty acids and risk of all cause mortality, cardiovascular disease, and type 2 diabetes: systematic review and meta-analysis of observational studies'. *BMJ*, vol. 351, h3978.
- DellaValle, D. M. (2013), 'Iron supplementation for female athletes: effects on iron status and performance outcomes', *Current Sports Medicine Reports*, 12(4), pp. 234–239. doi: 10.1249/JSR.0b013e31829a6f6b.
- DeMarco, H. M. *et al.* (1999), *Med. Sci. Sports Ex.*, vol. 31(1), pp. 164–70.
- Department of Health & Social Care (2016), 'UK Chief Medical Officers' Low Risk Drinking Guidelines'.
- Department of Health & Social Care (2019), 'UK Chief Medical Officers' Physical Activity Guidelines'.
- Department of Health (1991), 'Dietary Reference Values for Food Energy and Nutrients for the United Kingdom'. London: HMSO.
- Department of Health (2004), 'At least five a week: Evidence on the impact of physical activity and its relationship to health. A report from the Chief Medical Officer'.

- Derave, W. *et al.* (2007), 'Beta-alanine supplementation augments muscle carnosine content and attenuates fatigue during repeated isokinetic contraction bouts in trained sprinters'. *J. Appl. Physiol.*, vol. 103, pp. 1736–43.
- Desbrow, B. *et al.* (2004), 'Carbohydrate-electrolyte feedings and 1 h time-trial cycling performance'. *Int. J. Sport Nutr. Exerc. Metab.*, vol. 14, pp. 541–9.
- Desbrow, B. *et al.* (2012), 'The effects of different doses of caffeine on endurance cycling time trial performance', *Journal of Sports Sciences*, 30(2), pp. 115–120. doi: 10.1080/02640414.2011.632431.
- Desbrow, B. *et al.* (2014a), 'Comparing the rehydration potential of different milk-based drinks to a carbohydrate-electrolyte beverage'. *Appl. Physiol. Nutr. Metab.*, vol. 39(12), pp. 1366–72.
- Desbrow, B. *et al.* (2014b), 'Sports Dietitians Australia position statement: sports nutrition for the adolescent athlete', *International Journal of Sport Nutrition and Exercise Metabolism*, 24(5), pp. 570–584. doi: 10.1123/ijsem.2014-0031.
- Deutz, N. E. P. *et al.* (2014), 'Protein intake and exercise for optimal muscle function with aging: Recommendations from the ESPEN Expert Group'. *Clin. Nutr.*, vol. 33(6), pp. 929–36.
- Devries, M. C. and Phillips, S. M. (2014), 'Creatine supplementation during resistance training in older adults-a meta-analysis', *Medicine and Science in Sports and Exercise*, 46(6), pp. 1194–1203. doi: 10.1249/MSS.0000000000000220.
- Devries, M. C. *et al.* (2018), 'Changes in Kidney Function Do Not Differ between Healthy Adults Consuming Higher- Compared with Lower- or Normal-Protein Diets: A Systematic Review and Meta-Analysis', *The Journal of Nutrition*, 148(11), pp. 1760–1775. doi: 10.1093/jn/nxy197.
- Dodd, H. *et al.* (2011), 'Calculating meal glycaemic index by using measured and published food values compared with directly measured meal glycaemic index'. *Am. J. Clin. Nutr.*, vol. 95, pp. 992–6.
- Dodd, S. L. *et al.* (1993), 'Caffeine and exercise performance'. *Sports Med.*, vol. 15, pp. 14–23.
- Donnelly, J. E. *et al.* (2009), 'American College of Sports Medicine Position Stand. Appropriate physical activity intervention strategies for weight loss and prevention of weight regain for adults', *Medicine and Science in Sports and Exercise*, 41(2), pp. 459–471. doi: 10.1249/MSS.0b013e3181949333.
- Downes, J. W. (2002), 'The master's athlete: Defying aging'. *Topics in Clinical Chiropractic*, vol. 9(2), pp. 53–9.
- Drew, M. *et al.* (2018), 'Prevalence of illness, poor mental health and sleep quality and low energy availability prior to the 2016 Summer Olympic Games', *British Journal of Sports Medicine*, 52(1), pp. 47–53. doi: 10.1136/bjsports-2017-098208.
- Drew, M. *et al.* (2018), 'Prevalence of illness, poor mental health and sleep quality and low energy availability prior to the 2016 Summer Olympic Games', *British Journal of Sports Medicine*, 52(1), pp. 47–53. doi: 10.1136/bjsports-2017-098208.
- Drew, M. K. *et al.* (2017), 'A multifactorial evaluation of illness risk factors in athletes preparing for the Summer Olympic Games', *Journal of Science and Medicine in Sport*, 20(8), pp. 745–750. doi: 10.1016/j.jsams.2017.02.010.
- Drinkwater, B. L. (1986), 'Bone mineral content after resumption of menses in amenorrheic athletes'. *JAMA*, vol. 256, pp. 380–2.
- Drinkwater, B. L. *et al.* (1984), 'Bone mineral content of amenorrheic and eumenorrheic athletes'. *New England J. Med.*, vol. 311, pp. 277–81.
- Drouin-Chartier, J.-P. *et al.* (2016), 'Comprehensive Review of the Impact of Dairy Foods and Dairy Fat on Cardiometabolic Risk', *Advances in Nutrition*, 7(6), pp. 1041–1051. doi:

10.3945/an.115.011619.

- Drummond, M. J. and Rasmussen, B. B. (2008), 'Leucine-enriched nutrients and the regulation of mammalian target of rapamycin signalling and human skeletal muscle protein synthesis'. *Curr. Opin. Clin. Nutr. Metab. Care*, vol. 11, pp. 222–6.
- Ducker, K. J., Dawson, B., Wallman, K. E. (2013), 'Effect of beta-alanine supplementation on 800-m running performance'. *Int. J. Sport Nutr. Exerc. Metab.*, vol. 23(6), pp. 554–61.
- Dueck, C. A. *et al.* (1996), 'Role of energy balance in athletic menstrual dysfunction'. *Int. J. Sport Nutr.*, vol. 6, pp. 165–90.
- Duiven, E. *et al.* (2021), 'Undeclared Doping Substances are Highly Prevalent in Commercial Sports Nutrition Supplements', *Journal of Sports Science and Medicine*, pp. 328–338. doi: 10.52082/jssm.2021.328.
- Dulloo, A. G. and Jacquet, J. (1998), 'Adaptive reduction in basal metabolic rate in response to food deprivation in humans: a role for feedback signals from fat stores', *The American Journal of Clinical Nutrition*, 68(3), pp. 599–606. doi: 10.1093/ajcn/68.3.599.
- Dulloo, A. G. *et al.* (1999), 'Efficacy of a green tea extract rich in catechin polyphenols and caffeine in increasing 24 hour energy expenditure and fat oxidation in humans'. *Am. J. Clin. Nutr.*, vol. 70, pp. 1040–5.
- Durnin, J. V. G. A. and Womersley, J. (1974), 'Body fat assessed from total body density and its estimation from skinfold thickness: measurements on 481 men and women ages from 16 to 72 Years'. *Brit. J. Nutr.*, vol. 32, p. 77.
- Easton, C. *et al.* (2007), 'Creatine and glycerol hyperhydration in trained subjects before exercise in the heat'. *Int. J. Sports Nutr. Exerc. Metab.*, vol. 17, pp. 70–91.
- Eckerson, J. M., Bull, A. J., Baechle, T. R., Fischer, C. A., O'Brien, D. C., Moore, G. A., Yee, J. C., Pulverenti, T. S. (2013), 'Acute ingestion of sugar-free Red Bull energy drink has no effect on upper body strength and muscular endurance in resistance trained men'. *J. Strength Cond. Res.*, vol. 27(8), pp. 2248–54.
- Edwards, J. R. *et al.* (1993), 'Energy balance in highly trained female endurance runners'. *Med. Sci. Sports Ex.*, vol. 25(12), pp. 1398–404.
- EFSA (2015), 'Scientific and technical assistance on food intended for sportspeople'. [http://www.efsa.europa.eu/sites/default/files/scientific\\_output/files/main\\_documents/871c](http://www.efsa.europa.eu/sites/default/files/scientific_output/files/main_documents/871c) accessed March 2016.
- EFSA Panel on Dietetic Products, Nutrition, and Allergies (2010), 'Scientific opinion on Dietary Reference Values for water'. *EFSA Journal*, vol. 8(3), p. 1459.
- Egli, L. *et al.* (2013), 'Exercise prevents fructose-induced hypertriglyceridemia in healthy young subjects'. *Diabetes*, vol. 62(7), pp. 2259–65.
- Eisinger, M. (1994), 'Nutrient intake of endurance runners with lacto-ovo vegetarian diet and regular Western diet'. *Z. Ernahrungswiss*, vol. 33, pp. 217–29.
- Elliot, T. A. *et al.* (2006), 'Milk ingestion stimulates net muscle protein synthesis following resistance exercise', *Med. Sci. Sports Exer.*, vol. 384, pp. 667–74.
- English, K. L. *et al.* (2016), 'Leucine partially protects muscle mass and function during bed rest in middle-aged adults', *The American Journal of Clinical Nutrition*, 103(2), pp. 465–473. doi: 10.3945/ajcn.115.112359.
- Erlenbusch, M. *et al.* (2005), 'Effect of high fat or high carbohydrate diets on endurance exercise: a meta-analysis'. *Int. J. Sport Nutr. Exerc. Metab.*, vol. 14, pp. 1–14.
- Estaki, M. *et al.* (2016), 'Cardiorespiratory fitness as a predictor of intestinal microbial diversity and distinct metagenomic functions', *Microbiome*, 4. doi: 10.1186/s40168-016-0189-7.
- Estruch, R. *et al.* (2013), 'Primary prevention of cardiovascular disease with a Mediterranean diet'. *N. Engl. J. Med.*, vol. 368, pp. 1279–90.

- European Food Safety Authority EFSA (2009), 'Scientific Opinion: Labelling reference intake values for n-3 and n-6 polyunsaturated fatty acids', *The EFSA Journal*, 2009; 1176, 1–11.
- Evans, G. H. *et al.* (2017), 'Optimizing the restoration and maintenance of fluid balance after exercise-induced dehydration', *Journal of Applied Physiology (Bethesda, Md.: 1985)*, 122(4), pp. 945–951. doi: 10.1152/jappphysiol.00745.2016.
- Evans, M. *et al.* (2019), 'No Benefit of Ingestion of a Ketone Monoester Supplement on 10-km Running Performance', *Medicine and Science in Sports and Exercise*, 51(12), pp. 2506–2515. doi: 10.1249/MSS.0000000000002065.
- Fagerberg, P. (2018), 'Negative Consequences of Low Energy Availability in Natural Male Bodybuilding: A Review', *International Journal of Sport Nutrition and Exercise Metabolism*, 28(4), pp. 385–402. doi: 10.1123/ijsnem.2016-0332.
- Fairchild, T. J. *et al.* (2002), 'Rapid carbohydrate loading after a short bout of near maximal-intensity exercise'. *Med. Sci. Sports Exer.*, pp. 980–6.
- Farajian, P. *et al.* (2004), 'Dietary intake and nutritional practices of elite Greek aquatic athletes'. *Int. J. Sport Nutr. Exerc. Metab.*, vol 14, pp. 574–85.
- Febbraio, M. A. and Stewart, K. L. (1996), 'CHO feeding before prolonged exercise: effect of glycaemic index on muscle glycogenolysis and exercise performance'. *J. Appl. Physiol.*, vol. 81, pp. 1115–20.
- Febbraio, M. A. *et al.* (2000), 'Effects of carbohydrate ingestion before and during exercise on glucose kinetics and performance'. *J. Appl. Physiol.*, vol. 89, pp. 2220–6.
- Feinman, R. D. *et al.* (2015), 'Dietary carbohydrate restriction as the first approach in diabetes management: critical review and evidence base', *Nutrition (Burbank, Los Angeles County, Calif.)*, 31(1), pp. 1–13. doi: 10.1016/j.nut.2014.06.011.
- Ference, B. A. *et al.* (2017), 'Low-density lipoproteins cause atherosclerotic cardiovascular disease. 1. Evidence from genetic, epidemiologic, and clinical studies. A consensus statement from the European Atherosclerosis Society Consensus Panel', *European Heart Journal*, 38(32), pp. 2459–2472. doi: 10.1093/eurheartj/ehx144.
- Ferguson-Stegall, L. *et al.* (2011), 'Aerobic exercise training adaptations are increased by post-exercise carbohydrate-protein supplementation'. *J. Nutr. Metab.*, Epub Jun 9.
- Fiala, K. A. *et al.* (2004), 'Rehydration with a caffeinated beverage during the non-exercise periods of 3 consecutive days of 2-a-day practices'. *Int. J. Sport Nutr. Exerc. Metab.*, vol 14. pp. 419–29.
- Fleck, S. J. and Reimers, K. J. (1994), 'The practice of making weight: does it affect performance?' *Strength and Cond.*, vol. 1, pp. 66–7.
- Flood, T. R. *et al.* (2020), 'Addition of pectin-alginate to a carbohydrate beverage does not maintain gastrointestinal barrier function during exercise in hot-humid conditions better than carbohydrate ingestion alone', *Applied Physiology, Nutrition, and Metabolism = Physiologie Appliquee, Nutrition Et Metabolisme*, 45(10), pp. 1145–1155. doi: 10.1139/apnm-2020-0118.
- Fogelholm, M. (1994), 'Effects of bodyweight reduction on sports performance'. *Sports Med.*, vol. 18(14), pp. 249–67.
- Fogelholm, M. (1995), 'Indicators of vitamin and mineral status in athletes' blood: a review'. *Int. J. Sports Nutr.*, vol. 5, pp. 267–84.
- Food Standards Agency (2003), *Safe upper levels for vitamins and minerals* (HMSO).
- Forbes, G. B. (2000), 'Body fat content influences the body composition response to nutrition and exercise', *Annals of the New York Academy of Sciences*, 904, pp. 359–365. doi: 10.1111/j.1749-6632.2000.tb06482.x.
- Forbes, S. C., Candow, D. G., Little, J. P., *et al.* (2007), 'Effect of Red Bull energy drink on

- repeated Wingate cycle performance and bench-press muscle endurance'. *Int. J. Sport Nutr. Exerc. Metab.*, vol. 17(5), pp. 433–44.
- Foster-Powell, K., Holt, S. and Brand-Miller, J. C. (2002), 'International table of glycaemic index and glycaemic load values: 2002'. *Am. J. Clin. Nutr.*, vol. 76, pp. 5–56.
- Frankenfield D. C. *et al.* (2005), 'Comparison of predictive equations for Resting Metabolic Rate in healthy nonobese and obese adults: a systematic review'. *J. Am. Diet. Assoc.*, vol. 105, pp. 775–89.
- Frentsos, J. A. and Baer, J. T. (1997), 'Increased energy and nutrient intake during training and competition improves elite triathletes' endurance performance'. *Int. J. Sport Nutr.*, vol. 7, pp. 61–71.
- Funnell, M. P. *et al.* (2019), 'Blinded and unblinded hypohydration similarly impair cycling time trial performance in the heat in trained cyclists', *Journal of Applied Physiology (Bethesda, Md.: 1985)*, 126(4), pp. 870–879. doi: 10.1152/jappphysiol.01026.2018.
- Gabel, K. *et al.* (2018), 'Effects of 8-hour time restricted feeding on body weight and metabolic disease risk factors in obese adults: A pilot study', *Nutrition and Healthy Aging*, 4(4), pp. 345–353. doi: 10.3233/NHA-170036.
- Gacini, A. A., Rahnema, N., Hamedinia, M. R. (2006), 'Effects of vitamin E supplementation on oxidative stress at rest and after exercise to exhaustion in athletic students'. *J. Sports Med. Phys. Fitness*, vol. 46(3), pp. 458–61.
- Gallagher, D. *et al.* (2000) 'Healthy percentage body fat ranges: an approach for developing guidelines based on body mass index', *The American Journal of Clinical Nutrition*, 72(3), pp. 694–701. doi:10.1093/ajcn/72.3.694.
- Gallagher, J. *et al.* (2018), 'Oral health and performance impacts in elite and professional athletes', *Community Dentistry and Oral Epidemiology*, 46(6), pp. 563–568. doi: <https://doi.org/10.1111/cdoe.12392>.
- Galloway, S. D. R. and Maughan, R. (2000), 'The effects of substrate and fluid provision on thermoregulatory and metabolic responses to prolonged exercise in a hot environment'. *J. Sports Sci.*, vol. 18(5), pp. 339–51.
- Ganio, M. S. *et al.* (2009), 'Effect of caffeine on sport-specific endurance performance: a systematic review', *Journal of Strength and Conditioning Research*, 23(1), pp. 315–324. doi: 10.1519/JSC.0b013e31818b979a.
- Gant, N., Ali, A. and Foskett, A. (2010), 'The influence of caffeine and carbohydrate coingestion on simulated soccer performance'. *Int. J. Sport Nutr. Exerc. Metab.*, vol. 20, pp. 191–7.
- Gardner, C. D. *et al.* (2018), 'Effect of Low-Fat vs Low-Carbohydrate Diet on 12-Month Weight Loss in Overweight Adults and the Association With Genotype Pattern or Insulin Secretion: The DIETFITS Randomized Clinical Trial', *JAMA*, 319(7), pp. 667–679. doi: 10.1001/jama.2018.0245.
- Gardner, C. D. *et al.* (2019), 'Maximizing the intersection of human health and the health of the environment with regard to the amount and type of protein produced and consumed in the United States', *Nutrition Reviews*, 77(4), pp. 197–215. doi: 10.1093/nutrit/nuy073.
- Garfinkel, P. E. and Garner, D. M. (1982), *Anorexia nervosa: a multidimensional perspective*. (Brunner/Mazel).
- Garthe, I. and Maughan, R. J. (2018), 'Athletes and Supplements: Prevalence and Perspectives', *International Journal of Sport Nutrition and Exercise Metabolism*, 28(2), pp. 126–138. doi: 10.1123/ijsnem.2017-0429.
- Garthe, I. *et al.* (2011), 'Effect of two different weight-loss rates on body composition and strength and power-related performance in elite athletes', *International Journal of Sport Nutrition and Exercise Metabolism*, 21(2), pp. 97–104. doi: 10.1123/ijsnem.21.2.97.



- Garvican-Lewis, L. A. *et al.* (2018), 'Intravenous Iron Does Not Augment the Hemoglobin Mass Response to Simulated Hypoxia', *Medicine and Science in Sports and Exercise*, 50(8), pp. 1669–1678. doi: 10.1249/MSS.0000000000001608.
- Gaskell, S. K. *et al.* (2020), 'Impact of 24-h high and low fermentable oligo-, di-, monosaccharide, and polyol diets on markers of exercise-induced gastrointestinal syndrome in response to exertional heat stress', *Applied Physiology, Nutrition, and Metabolism = Physiologie Appliquee, Nutrition Et Metabolisme*, 45(6), pp. 569–580. doi: 10.1139/apnm-2019-0187.
- GBD 2016 Alcohol Collaborators (20182016), 'Alcohol use and burden for 195 countries and territories, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016', *Lancet (London, England)*, 392(10152), pp. 1015–1035. doi: 10.1016/S0140-6736(18)31310-2.
- Gejl, K. D. and Nybo, L. (2021), 'Performance effects of periodized carbohydrate restriction in endurance trained athletes – a systematic review and meta-analysis', *Journal of the International Society of Sports Nutrition*, 18(1), p. 37. doi: 10.1186/s12970-021-00435-3.
- German, J. B. *et al.* (2009), 'A reappraisal of the impact of dairy foods and milk fat on cardiovascular disease risk'. *European Journal of Nutrition*, vol. 48(4), pp. 191–203.
- Geyer, H., Parr. M. K., Mareck, U., *et al.* (2004), 'Analysis of non-hormonal nutritional supplements for anabolic-androgenic steroids – results of an international study'. *Int. J. Sports Med.*, vol. 25(2), pp. 124–9.
- Gibala, M. J. (2000), 'Nutritional supplementation and resistance exercise: what is the evidence for enhanced skeletal muscle hypertrophy?' *Can. J. Appl. Physiol.*, vol. 25(6), pp. 524–35.
- Gilchrist, M., Winyard, P. G., and Benjamin, N. (2010), 'Dietary nitrate – good or bad?' *Nitric Oxide*, vol. 22, pp. 104–9.
- Gill, S. and Panda, S. (2015), 'A smartphone app reveals erratic diurnal eating patterns in humans that can be modulated for health benefits', *Cell metabolism*, 22(5), pp. 789–798. doi: 10.1016/j.cmet.2015.09.005.
- Gilson, S. F. *et al.* (2009), 'Effects of chocolate milk consumption on markers of muscle recovery during intensified soccer training'. *Medicine and Science in Sports and Exercise*, vol. 41, p. S577.
- Gisolphi, C. V. *et al.* (1992), 'Intestinal water absorption from select carbohydrate solutions in humans'. *J. Appl. Physiol.*, vol. 73, pp. 2142–50.
- Gisolphi, C. V. *et al.* (1995), 'Effect of sodium concentration in a carbohydrate-electrolyte solution on intestinal absorption', *Med. Sci. Sports Ex.*, vol. 27(10), pp. 1414–20.
- Gleeson, M. (2011), 'Nutrition and immunity', In *Diet, Immunity and Inflammation*, Calder, P. C. and Yaqoob, P (eds), (Woodhead Publishing).
- Gleeson, M. (2016), 'Immunological aspects of sport nutrition', *Immunology & Cell Biology*, 94(2), pp. 117–123. doi: <https://doi.org/10.1038/icb.2015.109>.
- Gleeson, M. *et al.* (2008), 'Exercise and immune function: is there any evidence for probiotic benefit for sportspeople?' *Complete Nutrition*, vol. 8, pp. 35–7.
- Gleeson, M., Nieman, D. C. and Pedersen, B. K. (2004), 'Exercise, nutrition and immune function', *Journal of Sports Sciences*, 22(1), pp. 115–125. doi: 10.1080/0264041031000140590.
- Goldstone, A. P. *et al.* (2009), 'Fasting biases brain reward systems towards high-calorie foods', *Eur. J. Neurosci.*, vol 30(8), pp. 1625–35.
- Gombart, A. F., Pierre, A. and Maggini, S. (2020), 'A Review of Micronutrients and the Immune System-Working in Harmony to Reduce the Risk of Infection', *Nutrients*, 12(1). doi: 10.3390/nu12010236.
- Gomez-Cabrera *et al.* (2008), 'Oral administration of vitamin C decreases muscle mitochondrial

- biogenesis and hampers training-induced adaptations in endurance performance', *Am. J. Clin. Nutr.* 87(1), pp. 142–9.
- Gontzea, I. *et al.* (1975), 'The influence of adaptation to physical effort on nitrogen balance in man', *Nutr. Rep. Int.*, vol. 22, pp. 213–16.
- Gonzalez, J. T. *et al.* (2017), 'Glucose Plus Fructose Ingestion for Post-Exercise Recovery—Greater than the Sum of Its Parts?', *Nutrients*, 9(4), p. 344. doi: 10.3390/nu9040344.
- Gonzalez-Alonzo, J. *et al.* (1992), 'Rehydration after exercise with common beverages and water', *Int. J. Sports Med.*, vol. 13, pp. 399–406.
- Gordon, C. M. *et al.* (2017), 'Functional Hypothalamic Amenorrhea: An Endocrine Society Clinical Practice Guideline', *The Journal of Clinical Endocrinology and Metabolism*, 102(5), pp. 1413–1439. doi: 10.1210/jc.2017-00131.
- Goulet, E. D. B. (2011), 'Effects of exercise-induced dehydration on time-trial exercise performance: a metaanalysis', *Brit. J. Sports Med.*, vol. 45, pp. 1149–56.
- Goulet, E. D. B. (2013), 'Effect of exercise-induced dehydration on endurance performance: evaluating the impact of exercise protocols on outcomes using a meta-analytic procedure'. *Br. J. Sports Med.*, vol. 47(11), pp. 679–86.
- Graham, T. E. and Spriet, L. L. (1991), 'Performance and metabolic responses to a high caffeine dose during prolonged exercise', *J. Appl. Physiol.*, vol. 71(6), pp. 2292–8.
- Grandjean, A. (2000), 'The effect of caffeinated, non-caffeinated, caloric and non-caloric beverages on hydration', *J. Am. Coll. Nutr.*, vol. 19, pp. 591–600.
- Gray-Donald, K. *et al.* (2014), 'Protein intake protects against weight loss in healthy community-dwelling older adults', *J. Nutr.* vol. 144, pp. 321–6.
- Green, A. L. *et al.* (1996), 'Carbohydrate augments creatine accumulation during creatine supplementation in humans', *Am. J. Physiol.*, vol. 271, E821–6.
- Greenhaff, P. L. (1997), 'Creatine supplementation and implications for exercise performance and guidelines for creatine supplementation', In A. Jeukendrup *et al.* (eds), *Advances in Training and Nutrition for Endurance Sports* (Maastricht: Novertis Nutrition Research Unit), pp. 8–11.
- Greenwood, M. *et al.* (2003), 'Creatine supplementation during college football training does not increase the incidence of cramping or injury', *Molecular and Cellular Biochemistry*, 244(1–2), pp. 83–88.
- Grgic, J. and Pickering, C. (2019), 'The effects of caffeine ingestion on isokinetic muscular strength: A meta-analysis', *Journal of Science and Medicine in Sport*, 22(3), pp. 353–360. doi: 10.1016/j.jsams.2018.08.016.
- Grgic, J. *et al.* (2018), 'Effects of caffeine intake on muscle strength and power: a systematic review and meta-analysis', *Journal of the International Society of Sports Nutrition*, 15, p. 11. doi: 10.1186/s12970-018-0216-0.
- Groen, B. *et al.* (2012), 'Intragastric protein administration stimulates overnight muscle protein synthesis in elderly men', *Am. J. Physiol. Endocrinol. Metab.*, vol. 302, pp. 52–60.
- GSSI (1995), 'Roundtable on methods of weight gain in athletes'. *Sports Science Exchange*, vol. 6(3), pp. 1–4.
- Gualano, B., *et al.* (2012), 'In sickness and in health: The widespread application of creatine supplementation', *Amino Acids*, vol. 43, pp. 519–29.
- Guest, N. *et al.* (2018), 'Caffeine, CYP1A2 Genotype, and Endurance Performance in Athletes', *Medicine and Science in Sports and Exercise*, 50(8), pp. 1570–1578. doi: 10.1249/MSS.0000000000001596.
- Guest, N. S. and Barr, S. (2005), 'Cognitive dietary restraint is associated with stress fractures in women runners', *Int. J. Sports Nutr. Exerc. Metab.*, vol. 15, pp. 147–59.

- Guest, N. S. *et al.* (2021), 'International society of sports nutrition position stand: caffeine and exercise performance', *Journal of the International Society of Sports Nutrition*, 18(1), p. 1. doi: 10.1186/s12970-020-00383-4.
- Hackney, A. C., Sinning, W. E. and Bruot, B. C. (1988), 'Reproductive hormonal profiles of endurance-trained and untrained males', *Medicine and Science in Sports and Exercise*, 20(1), pp. 60–65. doi: 10.1249/00005768-198802000-00009.
- Haghighatdoost, F. *et al.* (2017), 'Association of vegetarian diet with inflammatory biomarkers: a systematic review and meta-analysis of observational studies', *Public Health Nutrition*, 20(15), pp. 2713–2721. doi: 10.1017/S1368980017001768.
- Hall, K. D. and Guo, J. (2017), 'Obesity Energetics: Body Weight Regulation and the Effects of Diet Composition', *Gastroenterology*, 152(7), pp. 1718-1727.e3. doi: 10.1053/j.gastro.2017.01.052.
- Hall, Kevin D. *et al.* (2015), 'Calorie for calorie, dietary fat restriction results in more body fat loss than carbohydrate restriction in people with obesity', *Cell Metabolism*, vol. 22(3), pp. 427–36.
- Halliday, T. *et al.* (2011), 'Vitamin D status relative to diet, lifestyle, injury and illness in college athletes', *Med. Sci. Sports Exerc.*, vol. 43, pp. 335–43.
- Halliwell, B. and Gutteridge, J. M. C. (1985), *Free Radicals in Biology and Medicine* (Clarendon Press), pp. 162–4.
- Hamilton, B. (2011), 'Vitamin D and athletic performance: the potential role of muscle', *Asian J. Sports Med.*, vol. 2(4), pp. 211–19.
- Hammond, K. M. *et al.* (2019), 'Post-exercise carbohydrate and energy availability induce independent effects on skeletal muscle cell signalling and bone turnover: implications for training adaptation', *The Journal of Physiology*, 597(18), pp. 4779–4796. doi: <https://doi.org/10.1113/JP278209>.
- Hanne, N., Dlin, R. and Rotstein, A. (1986), 'Physical fitness, anthropometric and metabolic parameters in vegetarian athletes'. *J. Sports Med. Phys. Fitness.*, vol. 26, pp. 180–5.
- Hansen, A. K. *et al.* (2005), 'Skeletal muscle adaptation: training twice every second day vs. training once daily', *J. Appl. Physiol.*, vol 98(1), pp. 93–9.
- Hao, Q. *et al.* (2011), 'Probiotics for preventing acute upper respiratory tract infections', *The Cochrane Database of Systematic Reviews*, (9), p. CD006895. doi: 10.1002/14651858.CD006895.pub2.
- Hao, Q. *et al.* (2011), 'Probiotics for preventing acute upper respiratory tract infections', *The Cochrane Database of Systematic Reviews*, (9), p. CD006895. doi: 10.1002/14651858.CD006895.pub2.
- Hao, Q., Dong, B. R. and Wu, T. (2015), 'Probiotics for preventing acute upper respiratory tract infections', *The Cochrane Database of Systematic Reviews*, (2), p. CD006895. doi: 10.1002/14651858.CD006895.pub3.
- Hargreaves, M. and Snow, R. (2001), 'Amino acids and endurance exercise', *Int. J. Sport Nutr.*, vol. 11, pp. 133–145.
- Hargreaves, M. *et al.* (2004), 'Pre-exercise carbohydrate and fat ingestion: effects on metabolism and performance', *J. Sports Sci.*, vol. 22(1), pp. 31–38.
- Harris, R. C. (1998), 'Ergogenics 1', *Peak Performance*, vol. 112, pp. 2–6.
- Harris, R. C. *et al.* (2006), 'The absorption of orally supplied beta-alanine and its effect on muscle carnosine synthesis in human vastus lateralis', *Amino Acids*, vol. 30(3), pp. 279–89.
- Harrison, S. E. *et al.* (2021), 'Influence of Vitamin D Supplementation by Simulated Sunlight or Oral D3 on Respiratory Infection during Military Training', *Medicine and Science in Sports and Exercise*. doi: 10.1249/MSS.0000000000002604.

- Hartman, J. W. *et al.* (2007), 'Consumption of fat-free fluid milk after resistance exercise promotes greater lean mass accretion than does consumption of soysoya or carbohydrate in young, novice, male weightlifters', *Am. J. Clin. Nutr.*, vol. 86(2), pp. 373–81.
- Harvey, K. L., Holcomb, L. E. and Kolwicz, S. C. (2019), 'Ketogenic Diets and Exercise Performance', *Nutrients*, 11(10), p. 2296. doi: 10.3390/nu11102296.
- Hatori, M. *et al.* (2012), 'Time-Restricted Feeding without Reducing Caloric Intake Prevents Metabolic Diseases in Mice Fed a High-Fat Diet', *Cell Metabolism*, 15(6), pp. 848–860. doi: 10.1016/j.cmet.2012.04.019.
- Haub, M. D. (1998), 'Acute l-glutamine ingestion does not improve maximal effort exercise', *J. Sport Med. Phys. Fitness*, vol. 38, pp. 240–4.
- Haub, M. D. *et al.* (2002), 'Effect of protein source on resistive-training-induced changes in body composition and muscle size in older men', *Am. J. Clin. Nutr.*, vol. 76, pp. 511–17.
- Haussinger, D. *et al.* (1996), 'The role of cellular hydration in the regulation of cell functioning', *Biochem. J.*, vol. 31, pp. 697–710.
- Havemann, L. *et al.* (2006), 'Fat adaptation followed by carbohydrate loading compromises high-intensity sprint performance', *J. Appl. Physiol.*, vol. 100(1), pp. 194–202.
- Hawley, J. A. and Burke, L. M. (2010), 'Carbohydrate availability and training adaptation: effects on cell metabolism', *Exerc. Sports Sci. Rev.*, vol. 38, pp. 152–160.
- Hawley, J. A. and Leckey, J. (2015), 'Carbohydrate dependence during prolonged, intense endurance exercise', *Sports Med.*, vol. 45 (Suppl 1), pp. S5–S12.
- Hawley, J. A. and Lessard, S. J. (2008), 'Exercise training-induced improvements in insulin action', *Acta Physiol (Oxf)*, vol. 192(1), pp. 127–35.
- Hawley, J. *et al.* (1997), 'Carbohydrate loading and exercise performance', *Sports Med.*, vol. 24(1), pp. 1–10.
- Hawley, J. *et al.* (2011), 'Nutritional modulation of training-induced skeletal muscle adaptations', *J. Appl. Physiol.*, vol. 100, pp. 834–45.
- He, C.-S. *et al.* (2013), 'Influence of vitamin D status on respiratory infection incidence and immune function during 4 months of winter training in endurance sport athletes', *Exercise Immunology Review*, 19, pp. 86–101.
- Heaney, R. P. (2011), 'Assessing vitamin D status', *Curr. Opin. Clin. Nutr. Metab. Care*, vol. 14, pp. 440–4.
- Heaney, R. P. (2013), 'Health is better at serum 25(OH) D above 30ng/mL', *J. Steroid Biochem. Mol. Biol.*, vol. 136, pp. 224–8.
- Hector, A. J. and Phillips, S. M. (2018), 'Protein Recommendations for Weight Loss in Elite Athletes: A Focus on Body Composition and Performance', *International Journal of Sport Nutrition and Exercise Metabolism*, 28(2), pp. 170–177. doi: 10.1123/ijsnem.2017-0273.
- Heikura, I. A., Burke, L. M., *et al.* (2018) (a), 'Impact of Energy Availability, Health, and Sex on Hemoglobin-Mass Responses Following Live-High-Train-High Altitude Training in Elite Female and Male Distance Athletes', *International Journal of Sports Physiology and Performance*, 13(8), pp. 1090–1096. doi: 10.1123/ijsp.2017-0547.
- Heikura, I. A., Uusitalo, A. L. T., *et al.* (2018) (b), 'Low Energy Availability Is Difficult to Assess but Outcomes Have Large Impact on Bone Injury Rates in Elite Distance Athletes', *International Journal of Sport Nutrition and Exercise Metabolism*, 28(4), pp. 403–411. doi: 10.1123/ijsnem.2017-0313.
- Helge, J. W. *et al.* (2001), 'Fat utilisation during exercise: adaptation to a fat rich diet increases utilisation of plasma fatty acids and very low density lipoprotein-triacylglycerol in humans', *J. Physiol.*, vol. 537; 3, pp. 1009–20.
- Helms, E. R. *et al.* (2014) (a), 'A systematic review of dietary protein during caloric restriction in

- resistance trained lean athletes: a case for higher intakes', *International Journal of Sport Nutrition and Exercise Metabolism*, 24(2), pp. 127–138. doi: 10.1123/ijsnem.2013-0054.
- Helms, E. R. *et al.* (2014), 'A systematic review of dietary protein during caloric restriction in resistance trained lean athletes: a case for higher intakes'. *Int. J. Sport Nutr. Exerc. Metab.*, vol. 24(2), pp. 127–38.
- Helms, E. R., Aragon, A. A. and Fitschen, P. J. (2014) (b), 'Evidence-based recommendations for natural bodybuilding contest preparation: nutrition and supplementation', *Journal of the International Society of Sports Nutrition*, 11(1), p. 20. doi: 10.1186/1550-2783-11-20.
- Hemilä, H. (2011), Zinc lozenges may shorten the duration of colds: a systematic review. *The open respiratory medicine journal*, 5(1).
- Hemilä, H. (2017), 'Zinc lozenges and the common cold: a meta-analysis comparing zinc acetate and zinc gluconate, and the role of zinc dosage', *JRSM open*, 8(5), p. 2054270417694291. doi: 10.1177/2054270417694291.
- Hemilä, H. and Chalker, E. (2013), 'Vitamin C for preventing and treating the common cold', *The Cochrane Database of Systematic Reviews*, (1), p. CD000980. doi: 10.1002/14651858.CD000980.pub4.
- Herman, P. and Polivy, J. (1991), 'Fat is a psychological issue', *New Scientist*, 16 Nov., pp. 41–5.
- Hespel, P. *et al.* (2001), 'Oral creatine supplementation facilitates the rehabilitation of disuse atrophy and alters the expression of muscle myogenic factors in humans', *The Journal of Physiology*, 536(Pt 2), pp. 625–633. doi: 10.1111/j.1469-7793.2001.0625c.xd.
- Hevia-Larraín, V. *et al.* (2021), 'High-Protein Plant-Based Diet Versus a Protein-Matched Omnivorous Diet to Support Resistance Training Adaptations: A Comparison Between Habitual Vegans and Omnivores', *Sports Medicine*. doi: 10.1007/s40279-021-01434-9.
- Hew-Butler, T. *et al.* (2015), 'Statement of the Third International Exercise-Associated Hyponatremia Consensus Development Conference, Carlsbad, California, 2015', *Clinical Journal of Sport Medicine*, 25(4), pp. 303–320. doi: 10.1097/JSM.0000000000000221.
- Hickey, H. S. *et al.* (1994), 'Drinking behaviour and exercise-thermal stress: role of drink carbonation', *Int. J. Sport Nutr.*, vol. 4, pp. 8–12.
- Higgins, S. *et al.* (2015), 'The effects of pre-exercise caffeinated-coffee ingestion on endurance performance: an evidence-based review', *Int. J. Sport Nutr. Exerc. Metab.*, Nov 16 (Epub ahead of print).
- Hill, A. M. *et al.* (2007), 'Combining fish-oil supplements with regular aerobic exercise improves body composition and cardiovascular disease risk factors'. *Am. J. Clin. Nutr.*, vol. 85(5), pp. 1267–74..
- Hingley, L. *et al.* (2017), 'DHA-rich Fish Oil Increases the Omega-3 Index and Lowers the Oxygen Cost of Physiologically Stressful Cycling in Trained Individuals. *International journal of sport nutrition and exercise metabolism*. doi: 10.1123/ijsnem.2016-0150.
- Hitchins, S. *et al.* (1999), 'Glycerol hyperhydration improves cycle time-trial performance in hot humid conditions', *Eur. J. Appl. Physiol. Occup. Physiol.*, vol. 80(5), pp. 494–501.
- Hobson, R. M. *et al.* (2012), 'Effects of -alanine supplementation on exercise performance: a meta-analysis', *Amino Acids*, vol. 43(1), pp. 25–37.
- Hodgson, A. B., Randell, R. K. and Jeukendrup, A. E. (2013) (a), 'The metabolic and performance effects of caffeine compared to coffee during endurance exercise', *PloS One*, 8(4), p. e59561. doi: 10.1371/journal.pone.0059561.
- Hodgson, A. B., Randell, R. K. and Jeukendrup, A. E. (2013) (b), 'The Effect of Green Tea Extract on Fat Oxidation at Rest and during Exercise: Evidence of Efficacy and Proposed Mechanisms', *Advances in Nutrition*, 4(2), pp. 129–140. doi: 10.3945/an.112.003269.
- Hoffman, J. R. *et al.* (2008), 'Short-duration beta-alanine supplementation increases training

- volume and reduces subjective feelings of fatigue in college football players', *Nutr. Res.*, vol. 28, pp. 31–5.
- Hoffman, M. D. *et al.* (2016), 'VIEW: Is Drinking to Thirst Adequate to Appropriately Maintain Hydration Status During Prolonged Endurance Exercise? Yes', *Wilderness & Environmental Medicine*, 27(2), pp. 192–195. doi: 10.1016/j.wem.2016.03.003.
- Holt, S. J. (1992), 'Relationship of satiety to postprandial glycaemic, insulin and cholecystokinin responses'. *Appetite*, vol. 18, pp. 129–41.
- Hoon, M. W. *et al.* (2014), 'Nitrate supplementation and high-intensity performance in competitive cyclists', *Applied Physiology, Nutrition, and Metabolism*, 0, 0, 10.1139/apnm-2013-0574.
- Hoon, MW1, Johnson NA, Chapman PG, Burke LM (2013), The effect of nitrate supplementation on exercise performance in healthy individuals: a systematic review and meta-analysis. *Int J Sport Nutr Exerc Metab.* 2013 Oct; 23(5):522–32. Epub 2013 Apr 9.
- Hooper, L. *et al.* (2012), 'Reduced or modified dietary fat for preventing cardiovascular disease', *Cochrane Database Syst. Rev.*, May 16;5.
- Hooper, L. *et al.* (2015) (a), 'Effects of total fat intake on body weight', *The Cochrane Database of Systematic Reviews*, (8), p. CD011834. doi: 10.1002/14651858.CD011834.
- Hooper, L. *et al.* (2015) (b), 'Reduction in saturated fat intake for cardiovascular disease', *The Cochrane Database of Systematic Reviews*, (6), p. CD011737. doi: 10.1002/14651858.CD011737.
- Hooper, L. *et al.* (2020), 'Reduction in saturated fat intake for cardiovascular disease', *Cochrane Database of Systematic Reviews*, (5). doi: 10.1002/14651858.CD011737.pub2.
- Hord N. G., Tang, Y. and Bryan, N. S. (2009), 'Food sources of nitrates and nitrites: the physiologic context for potential health benefits', *Am. J. Clin. Nutr.*, vol. 90(1), pp. 1–10.
- Houtkooper, L. B. (2000), 'Body composition', in Manore, M. M. and Thompson, J. L., *Sport Nutrition for Health and Performance*, Human Kinetics, pp. 199–219.
- Howarth, K. R. *et al.* (2009), 'Coingestion of protein with carbohydrate during recovery from endurance exercise stimulates skeletal muscle protein synthesis in humans', *J. Appl. Physiol.*, vol. 106, pp. 1394–1402.
- Howatson, G., McHugh, M. P., Hill, J. A., *et al.* (2010), 'Influence of tart cherry juice on indices of recovery following marathon running', *Scand. J. Med. Sci. Sports*, vol. 20(6), pp. 843–52.
- Howe, S. T. *et al.* (2013), 'The effect of beta-alanine supplementation on isokinetic force and cycling performance in highly trained cyclists', *Int. J. Sport Nutr. Exerc. Metab.* Dec, vol. 23(6), pp. 562–70 (Epub 2013 Apr 18).
- Hu, T. *et al.* (2012), 'Effects of low-carbohydrate diets versus low-fat diets on metabolic risk factors: a meta-analysis of randomized controlled clinical trials', *Am. J. Epidemiol.*, vol 176 Suppl 7, pp. S44–54.
- Hudson, J. L. *et al.* (2020), 'Protein Intake Greater than the RDA Differentially Influences Whole-Body Lean Mass Responses to Purposeful Catabolic and Anabolic Stressors: A Systematic Review and Meta-analysis', *Advances in Nutrition (Bethesda, Md.)*, 11(3), pp. 548–558. doi: 10.1093/advances/nmz106.
- Hughes, R. L. (2019), 'A Review of the Role of the Gut Microbiome in Personalized Sports Nutrition', *Frontiers in Nutrition*, 6, p. 191. doi: 10.3389/fnut.2019.00191.
- Hulston C. J. *et al.* (2010), 'Training with low muscle glycogen enhances fat metabolism in well-trained cyclists', *Med. Sci. Sports Exerc.*, vol. 42, pp. 2046–55.
- Hultman, E. *et al.* (1996), 'Muscle creatine loading in man', *J. Appl. Physiol.*, vol. 81, pp. 232–9.
- Igwe, E. O. *et al.* (2019), 'A systematic literature review of the effect of anthocyanins on gut microbiota populations', *Journal of Human Nutrition and Dietetics: The Official Journal of the*

- British Dietetic Association*, 32(1), pp. 53–62. doi: 10.1111/jhn.12582.
- Ihalainen, J. K. *et al.* (2021), ‘Body Composition, Energy Availability, Training, and Menstrual Status in Female Runners’, *International Journal of Sports Physiology and Performance*, pp. 1–6. doi: 10.1123/ijsp.2020-0276.
- Impey, S. G. *et al.* (2016), ‘Fuel for the work required: a practical approach to amalgamating train-low paradigms for endurance athletes’, *Physiological Reports*, 4(10). doi: 10.14814/phy2.12803.
- Impey, S. G. *et al.* (2018), ‘Fuel for the Work Required: A Theoretical Framework for Carbohydrate Periodization and the Glycogen Threshold Hypothesis’, *Sports Medicine (Auckland, N.Z.)*, 48(5), pp. 1031–1048. doi: 10.1007/s40279-018-0867-7.
- International Olympic Committee (IOC) (2005), ‘IOC Position Stand on the female athlete triad’ [http://www.olympic.org/assets/importednews/documents/en\\_report\\_917.pdf](http://www.olympic.org/assets/importednews/documents/en_report_917.pdf) (Accessed March 2016).
- International Olympic Committee (IOC) (2011), Consensus Statement on Sports Nutrition 2010, *Sports Sci* 4,29 Suppl 1: S3–4. <http://www.olympic.org/Documents/Reports/EN/CONSENSUSFINAL-v8-en.pdf>.
- Ivy, J. L. *et al.* (1988), ‘Muscle glycogen synthesis after exercise: effect of time of carbohydrate ingestion’, *J. Appl. Physiol.*, vol. 64, pp. 1480–5.
- Ivy, J. L. *et al.* (2002), ‘Early post-exercise muscle glycogen recovery is enhanced with carbohydrate-protein supplement’, *J. Appl. Physiol.*, vol. 93, pp. 1337–44.
- Ivy, J. L. *et al.* (2003), ‘Effect of a carbohydrate-protein supplement on endurance performance during exercise of varying intensity’. *Int. J. Sport Nutr. Exerc. Metab.*, vol. 13, pp. 388–401.
- Ivy, J. L. *et al.* (2009), ‘Improved cycling time-trial performance after ingestion of a caffeine energy drink’, *Int. J. Sport Nutr. Exerc. Metab.*, vol. 19(1), pp. 61–78.
- Jäger, R. *et al.* (2011), ‘Analysis of the efficacy, safety, and regulatory status of novel forms of creatine’, *Amino Acids*, vol. 40(5), pp. 1369–83.
- Jäger, R. *et al.* (2017), ‘International Society of Sports Nutrition Position Stand: protein and exercise’, *Journal of the International Society of Sports Nutrition*, 14(1), p. 20. doi: 10.1186/s12970-017-0177-8.
- Jäger, R. *et al.* (2019), ‘International Society of Sports Nutrition Position Stand: Probiotics’, *Journal of the International Society of Sports Nutrition*, 16(1), p. 62. doi: 10.1186/s12970-019-0329-0.
- Jakobsen, M. U. *et al.* (2009), ‘Major types of dietary fat and risk of coronary heart disease: a pooled analysis of 11 cohort studies’, *Am. J. Clin. Nutr.*, vol. 89, pp. 1425–32.
- Jakubowski, J. S. *et al.* (2020), ‘ki, J. S. –32. Supplementation with the Leucine Metabolite  $\beta$ -hydroxy- $\beta$ -methylbutyrate (HMB) does not Improve Resistance Exercise-Induced Changes in Body Composition or Strength in Young Subjects: A Systematic Review and Meta-Analysis’, on with the Leucine Metabolite ry fat and risk of coronary heart disease: a pooled analysis of 11 cohort studies’. y’. –4. [http://, abolic Stressors: A Systematic Review and Meta-analysis](http://abolicStressors: A Systematic Review and Meta-analysis)’, S44–54.nalysis th*Nutrients*, 12(5), p. 1523. doi: 10.3390/nu12051523.
- James, L. J. *et al.* (2017), ‘Hypohydration impairs endurance performance: a blinded study’, *Physiological Reports*, 5(12). doi: 10.14814/phy2.13315.
- James, L. J. *et al.* (2019) (a), ‘Cow’s milk as a post-exercise recovery drink: implications for performance and health’, *European Journal of Sport Science*, 19(1), pp. 40–48. doi: 10.1080/17461391.2018.1534989.
- James, L. J. *et al.* (2019) (b), ‘Does Hypohydration Really Impair Endurance Performance? Methodological Considerations for Interpreting Hydration Research’, *Sports Medicine (Auckland, N.Z.)*, 49(Suppl 2), pp. 103–114. doi: 10.1007/s40279-019-01188-5.

- Jamurtas, A. Z. *et al.* (2011), 'The effects of low and high glycemic index foods on exercise performance and beta-endorphin responses', *J. Int. Soc. Sports Nutr.*, vol. 8, p. 15.
- Janelle, K. C. and Barr, S. I. (1995), 'Nutrient intakes and eating behavior scores of vegetarian and non-vegetarian women', *J. Am. Diet. Assoc.*, vol. 95, pp. 180–6, 189.
- Jang, L.-G. *et al.* (2019), 'The combination of sport and sport-specific diet is associated with characteristics of gut microbiota: an observational study', *Journal of the International Society of Sports Nutrition*, 16(1), p. 21. doi: 10.1186/s12970-019-0290-y.
- Janssen, I. *et al.* (2000), 'Skeletal muscle mass and distribution in 468 men and women aged 18–88 yr', *J. Appl. Physiol.*, vol. 89(1), pp. 81–8.
- Jentjens, R. L. *et al.* (2001), 'Addition of protein and amino acids to carbohydrates does not enhance postexercise muscle glycogen synthesis', *Journal of Applied Physiology (Bethesda, Md.: 1985)*, 91(2), pp. 839–846. doi: 10.1152/jap.2001.91.2.839.
- Jentjens, R. L. P. G. *et al.* (2004), 'Oxidation of combined ingestion of glucose and fructose during exercise', *Journal of Applied Physiology*, 96(4), pp. 1277–1284. doi: 10.1152/jap.2004.96.4.1277.
- Jéquier, E. (2002), 'Pathways to obesity', *International Journal of Obesity and Related Metabolic Disorders: Journal of the International Association for the Study of Obesity*, 26 Suppl 2, pp. S12–17. doi: 10.1038/sj.ijo.0802123.
- Jeukendrup, A. E. (2004), 'Carbohydrate intake during exercise and performance', *Nutrition*, 20(7–8), pp. 669–677. doi: 10.1016/j.nut.2004.04.017.
- Jeukendrup, A. E. (2004), 'Carbohydrate intake during exercise and performance', *Nutrition*, vol. 20, pp. 669–77.
- Jeukendrup, A. E. (2008), 'Carbohydrate feeding during exercise', *Eur. J. Sports Sci.*, vol. 8(2), pp. 77–86.
- Jeukendrup, A. E. (2017), 'Training the Gut for Athletes', *Sports Medicine (Auckland, N.Z.)*, 47(Suppl 1), pp. 101–110. doi: 10.1007/s40279-017-0690-6.
- Jeukendrup, A. E. A. (2014), 'A step towards personalized sports nutrition: carbohydrate intake during exercise', *Sports Med.*, vol. 44 (Suppl 1), pp. 25–33.
- Jeukendrup, A. E. A. *et al.* (1997), 'Carbohydrate-electrolyte feedings improve 1-hour time trial cycling performance', *Int. J. Sports Med.*, vol. 18, pp. 125–9.
- Jeukendrup, A. E. A., *et al.* (2000), 'Relationship between gastro-intestinal complaints and endotoxaemia, cytokine release and the acute-phase reaction during and after a long-distance triathlon in highly trained men', *Clin. Sci. (Lond.)*, vol. 98(1), pp. 47–55.
- Jeukendrup, A. E.A. (2010), 'Carbohydrate and exercise performance: the role of multiple transportable carbohydrates', *Curr. Opin. Clin. Nutr. Metab. Care*, vol. 13(4), pp. 452–57.
- Jeukendrup, A. E.A. (2014), 'A step towards personalized sports nutrition: carbohydrate intake during exercise', *Sports Medicine (Auckland, N.Z.)*, 44 Suppl 1, pp. S25–33. doi: 10.1007/s40279-014-0148-z.
- Johnston, B. C. *et al.* (2014), 'Comparison of weight loss among named diet programs in overweight and obese adults: a meta-analysis', *JAMA*, 312(9), pp. 923–933. doi: 10.1001/jama.2014.10397.
- Johnston, C. S. *et al.* (2006), 'Ketogenic low-carbohydrate diets have no metabolic advantage over nonketogenic low-carbohydrate diets', *Am. J. Clin. Nutr.*, vol. 83(5), pp. 1055–61.
- Jonnalagadda, S. S. *et al.* (2004), 'Food preferences, dieting behaviours, and body image perceptions of elite figure skaters', *Int. J. Sports Nutr. Exerc. Metab.*, vol. 14, pp. 594–606.
- Josse, A. R. *et al.* (2010), 'Body composition and strength changes in women with milk and resistance exercise', *Med. Sci. Sports Exerc.*, vol. 42(6), pp. 1122–30.
- Jouris K. *et al.* (2011), 'The effect of omega-3 fatty acid supplementation on the inflammatory



- response to eccentric strength exercise'. *J. Sports Sci. Med.*, vol. 10, pp. 432–38.
- Jowko, E. *et al.* (2001), 'Creatine and HMB additively increase lean body mass and muscle strength during a weight training programme'. *Nutrition*, vol. 17(7), pp. 558–66.
- Joyce, S., *et al.* (2012), 'Acute and chronic loading of sodium bicarbonate in highly trained swimmers', *Eur. J. Appl. Physiol.*, vol. 112(2), pp. 461–9.
- Judkins, C. (2008), 'Investigation into supplementation contamination levels in the UK market', *HFL Sport Science*. [www.informed-sport.com](http://www.informed-sport.com)
- Kamber, M. *et al.* (2001), 'Nutritional supplements as a source for positive doping cases?' *Int. J. Sports Nutr.*, vol. 11, pp. 258–63.
- Kammer L. *et al.* (2009), 'Cereal and non-fat milk support muscle recovery following exercise', *J. Int. Soc. Sports Nutr.*, vol. 6, p. 11.
- Kannel, W. B. *et al.* (1986), 'Overall and coronary heart disease mortality rates in relation to major risk factors in 325,348 men screened for the MRFIT. Multiple Risk Factor Intervention Trial', *American Heart Journal*, 112(4), pp. 825–836. doi: 10.1016/0002-8703(86)90481-3.
- Karlssoon, J. and Saltin, B. (1971), 'Diet, muscle glycogen and endurance performance', *J. Appl. Physiol.*, vol. 31, pp. 201–6.
- Karp, J. R. *et al.* (2006), 'Chocolate milk as a post-exercise recovery aid', *Int. J. Sport Nutr. Exerc. Metab.*, vol. 16, pp. 78–91.
- Karsch–Völk, M., Barrett, B., Kiefer, D., Bauer, R., Ardjomand–Woelkart, K., & Linde, K. (2014), *Echinacea for preventing and treating the common cold.*, The Cochrane Library.
- Kasper, A. M. *et al.* (2015), 'Carbohydrate mouth rinse and caffeine improves high-intensity interval running capacity when carbohydrate restricted', *Eur. J. Sport Sci.*, vol. 2, pp. 1–9 (Epub ahead of print).
- Kasper, A. M. *et al.* (2019), 'Case Study: Extreme Weight Making Causes Relative Energy Deficiency, Dehydration, and Acute Kidney Injury in a Male Mixed Martial Arts Athlete', *International Journal of Sport Nutrition and Exercise Metabolism*, 29(3), pp. 331–338. doi: 10.1123/ijsnem.2018-0029.
- Kasper, A. M. *et al.* (2020), 'High Prevalence of Cannabidiol Use Within Male Professional Rugby Union and League Players: A Quest for Pain Relief and Enhanced Recovery', *International Journal of Sport Nutrition and Exercise Metabolism*, 30(5), pp. 315–322. doi: 10.1123/ijsnem.2020-0151.
- Katsanos, C. S. *et al.* (2006), 'A high proportion of leucine is required for optimal stimulation of the rate of muscle protein synthesis by essential amino acids in the elderly', *Am. J. Physiol. Endocrinol. Metab.*, vol. 291, pp. E381–E387.
- Kawabata, F. *et al.* (2014), 'Supplementation with eicosapentaenoic acid-rich fish oil improves exercise economy and reduces perceived exertion during submaximal steady-state exercise in normal healthy untrained men', *Bioscience, Biotechnology, and Biochemistry*, 78(12), pp. 2081–2088. doi: 10.1080/09168451.2014.946392.
- Kay, C. D. *et al.* (2012), 'Relative impact of flavonoid composition, dose and structure on vascular function: a systematic review of randomised controlled trials of flavonoid-rich food products', *Molecular Nutrition & Food Research*, 56(11), pp. 1605–1616. doi: 10.1002/mnfr.201200363.
- Keane, K. M. *et al.* (2018), 'Effects of montmorency tart cherry (L. Prunus Cerasus) consumption on nitric oxide biomarkers and exercise performance', *Scandinavian Journal of Medicine & Science in Sports*, 28(7), pp. 1746–1756. doi: 10.1111/sms.13088.
- Keay, N. *et al.* (2019), 'Clinical evaluation of education relating to nutrition and skeletal loading in competitive male road cyclists at risk of relative energy deficiency in sports (RED-S): 6-month randomised controlled trial', *BMJ Open Sport — Exercise Medicine*, 5(1). doi:

- 10.1136/bmjsem-2019-000523.
- Keay, N., Fogelman, I. and Blake, G. (1997), 'Bone mineral density in professional female dancers.', *British Journal of Sports Medicine*, 31(2), pp. 143–147. doi: 10.1136/bjsem.31.2.143.
- Keay, N., Francis, G. and Hind, K. (2018), 'Low energy availability assessed by a sport-specific questionnaire and clinical interview indicative of bone health, endocrine profile and cycling performance in competitive male cyclists', *BMJ Open Sport & Exercise Medicine*, 4(1), p. e000424. doi: 10.1136/bmjsem-2018-000424.
- Keay, N., Overseas, A. and Francis, G. (2020), 'Indicators and correlates of low energy availability in male and female dancers', *BMJ Open Sport & Exercise Medicine*, 6(1), p. e000906. doi: 10.1136/bmjsem-2020-000906.
- Keizer, H. A. *et al.* (1986), 'Influence of liquid or solid meals on muscle glycogen resynthesis, plasma fuel hormone response and maximal physical working capacity', *Int. J. Sports Med.*, vol. 8, pp. 99–104.
- Kenefick, R. W. and Cheuvront, S. N. (2012), 'Hydration for recreational sport and physical activity', *Nutrition Reviews*, 70(s2), pp. S137–S142. doi: <https://doi.org/10.1111/j.1753-4887.2012.00523.x>.
- Kennerly, K. *et al.* (2011), 'Influence of banana versus sports beverage ingestion on 75 km cycling performance and exercise-induced inflammation', *Med. Sci. Sports Exerc.*, vol. 43(5), pp. 340–341.
- Kenney, W. L. and Chiu, P. (2001), 'Influence of age on thirst and fluid intake', *Med. Sci. Sports Exerc.*, vol. 33(9), pp. 1524–32.
- Keohane, D. M. *et al.* (2019), 'Four men in a boat: Ultra-endurance exercise alters the gut microbiome', *Journal of Science and Medicine in Sport*, 22(9), pp. 1059–1064. doi: 10.1016/j.jsams.2019.04.004.
- Kerksick, C. M. *et al.* (2018), 'ISSN exercise & sports nutrition review update: research & recommendations', *Journal of the International Society of Sports Nutrition*, 15(1), p. 38. doi: 10.1186/s12970-018-0242-y.
- Key, T. J. A. *et al.* (1996), 'Dietary habits and mortality in a cohort of 11,000 vegetarians and health conscious people: results of a 17-year follow-up', *Br. Med. J.*, vol. 313, pp. 775–9.
- Keys, A.B. (1980), *Seven Countries: A Multivariate Analysis of Death and Coronary Heart Disease* (Harvard University Press, 1980).
- Kiens, B. *et al.* (1990), 'Benefit of simple carbohydrates on the early post-exercise muscle glycogen repletion in male athletes', *Med. Sci. Sports Ex.* (suppl.), S88.
- Killer, S.C. *et al.* (2014), 'No Evidence of Dehydration with Moderate Daily Coffee Intake: A Counterbalanced Cross-Over Study in a Free-Living Population', *PLoS ONE*, vol. 9 (1): e84154.
- Kim, H., Caulfield, L. E. and Rebholz, C. M. (2018), 'Healthy Plant-Based Diets Are Associated with Lower Risk of All-Cause Mortality in US Adults', *The Journal of Nutrition*, 148(4), pp. 624–631. doi: 10.1093/jn/nxy019.
- King, D. S. *et al.* (1999), 'Effects of oral androstenedione on serum testosterone and adaptations to resistance training in young men', *J. Am. Med. Assoc.*, vol. 281 (21), pp. 2020–8.
- King, S. *et al.* (2014), 'Effectiveness of probiotics on the duration of illness in healthy children and adults who develop common acute respiratory infectious conditions: a systematic review and meta-analysis', *The British Journal of Nutrition*, 112(1), pp. 41–54. doi: 10.1017/S0007114514000075.
- Kirkpatrick, C. F. *et al.* (2019), 'Review of current evidence and clinical recommendations on the effects of low-carbohydrate and very-low-carbohydrate (including ketogenic) diets for the

- management of body weight and other cardiometabolic risk factors: A scientific statement from the National Lipid Association Nutrition and Lifestyle Task Force', *Journal of Clinical Lipidology*, 13(5), pp. 689-711. e1. doi: 10.1016/j.jacl.2019.08.003.
- Knez, W. L. and Peake, J. M. (2010), 'The prevalence of vitamin supplementation in ultra-endurance triathletes', *Int. J. Sport Nutr. Exerc. Metab.*, vol. 20(6), pp. 507-14.
- Koopman, R. (2011), 'Dietary protein and exercise training in ageing', *Proc. Nutr. Soc.*, vol. 70, pp. 104-13.
- Koopman, R. *et al.* (2005), 'Combined ingestion of protein and free leucine with carbohydrate increases post-exercise muscle protein synthesis in vivo in male subjects', *Am. J. Physiol. Endocrinol. Metab.*, vol. 288(4), pp. E645-653.
- Kreider, R. (2003) (a), 'Effects of whey protein supplementation with casein or BCAA and glutamine on training adaptations: body composition', *Med. Sci. Sport Exerc.*, vol. 35(5), suppl. 1, p. S395.
- Kreider, R. B. (2003) (b), 'Effects of creatine supplementation on performance and training adaptations', *Mol. Cell Biochem.*, vol. 244(1-2), pp. 89-94.
- Kreider, R. B. *et al.* (1996), 'Effects of ingesting supplements designed to promote lean tissue accretion on body composition during resistance training', *Int. J. Sport Nutr.*, vol. 63, pp. 234-46.
- Kreider, R. B. *et al.* (2000), 'Effects of calcium-HMB supplementation during training on markers of catabolism, body composition, strength and sprint. performance', *J. Exerc. Physiol.*, vol. 3 (4), pp. 48-59.
- Kreider, R. B. *et al.* (2002), 'Effects of conjugated linoleic acid supplementation during resistance training on body composition, bone density, strength, and selected hematological markers', *J. Strength Cond. Res.*, vol. 16(3), pp. 325-34.
- Kreider, R. B. *et al.* (2017), 'International Society of Sports Nutrition position stand: safety and efficacy of creatine supplementation in exercise, sport, and medicine', *Journal of the International Society of Sports Nutrition*, 14(1), p. 18. doi: 10.1186/s12970-017-0173-z.
- Kristensen, K. *et al.* (1999), 'Hormone replacement therapy affects body composition and leptin differently in obese and non-obese postmenopausal women', *The Journal of Endocrinology*, 163(1), pp. 55-62. doi: 10.1677/joe.0.1630055.
- Kristiansen, M. *et al.* (2005), 'Dietary supplement use by varsity athletes at a Canadian University', *Int. J. Sport Nutr. Exerc. Metab.*, vol. 15, pp. 195-210.
- Lambert, E. V. *et al.* (1994), 'Enhanced endurance in trained cyclists during moderate intensity exercise following 2 weeks adaptation to a high fat diet', *Eur. J. Appl. Physiol.*, vol. 69, pp. 287-293.
- Lane, S. C. *et al.* (2013) (a), 'Effect of a carbohydrate mouth rinse on simulated cycling time-trial performance commenced in a fed or fasted state', *Applied Physiology, Nutrition, and Metabolism = Physiologie Appliquee, Nutrition Et Metabolisme*, 38(2), pp. 134-139. doi: 10.1139/apnm-2012-0300.
- Lane, S. C. *et al.* (2013) (b), 'Caffeine ingestion and cycling power output in a low or normal muscle glycogen state', *Med. Sci. Sports Exerc.*, Aug. 45(8), pp. 1577-84.
- Lane, S. C. *et al.* (2013) (c), 'Single and combined effects of beetroot juice and caffeine supplementation on cycling time-trial performance', *Appl. Physiol. Nutr. Metab.*, 10.1139/apnm-2013-0336
- Lane, S. C. *et al.* (2015), 'Effects of sleeping with reduced carbohydrate availability on acute training responses', *J. Appl. Physiol.*, vol. 119(6), pp. 643-55.
- Langan-Evans, C., Close, G. L. and Morton, J. P. (2011), 'Making Weight in Combat Sports', *Strength & Conditioning Journal*, 33(6), pp. 25-39. doi: 10.1519/SSC.0b013e318231bb64.

- Lansley, K. I. *et al.* (2011) (a), 'Dietary nitrate supplementation reduces the O<sub>2</sub> cost of walking and running: a placebo-controlled study', *J. Appl. Physiol.*, vol. 110, pp. 591–600.
- Lansley, K. I., *et al.* (2011) (b), 'Acute dietary nitrate supplementation improves cycling time-trial performance', *Med. Sci. Sports Exerc.*, vol. 43, pp. 1125–31.
- Lantzouni, E. *et al.* (2002), 'Reversibility of growth stunting in early onset anorexia nervosa: a prospective study', *The Journal of Adolescent Health: Official Publication of the Society for Adolescent Medicine*, 31(2), pp. 162–165. doi: 10.1016/s1054-139x(02)00342-7.
- Lanza, I. R. *et al.* (2008), 'Endurance exercise as a countermeasure for aging', *Diabetes*, vol. 57(11), pp. 2933–42.
- Larson-Meyer, D. E. and Willis, K. S. (2010), 'Vitamin D and athletes', *Curr. Sports Med. Rep.*, vol. 9(4), pp. 220–6.
- Lassale Camille, *et al.* (2015), "Abstract 16: A Pro-Vegetarian Food Pattern and Cardiovascular Mortality in the Epic Study." *Circulation*, vol. 131, no. suppl\_1, Mar. 2015, pp. A16–A16. ahajournals.org (Atypon), doi:10.1161/circ.131.suppl\_1.16.
- Laursen, P. B. *et al.* (2006), 'Core temperature and hydration status during an Ironman triathlon', *Br. J. Sports Med.* 2006 Apr; 40(4), pp. 320–5; discussion 325.
- Layman, D. K. *et al.* (2005), 'Dietary protein and exercise have additive effects on body composition during weight loss in adult women', *J. Nutr.* vol. 35(8), pp. 1903–10.
- Le, L. T. and Sabaté, J. (2014), 'Beyond meatless, the health effects of vegan diets: findings from the Adventist cohorts', *Nutrients*, 6(6), pp. 2131–2147. doi: 10.3390/nu6062131.
- Lean, M. E. J. *et al.* (1995), 'Waist circumference as a measure for indicating need for weight management', *BMJ*, vol. 311, pp. 158–61.
- Leckey, J. J. *et al.* (2016), 'Altering fatty acid availability does not impair prolonged, continuous running to fatigue: Evidence for carbohydrate dependence'. *J. Appl. Physiol.*, vol. 120(2), pp. 107–13.
- Leckey, J. J. *et al.* (2017), 'Ketone Diester Ingestion Impairs Time-Trial Performance in Professional Cyclists', *Frontiers in Physiology*, 8, p. 806. doi: 10.3389/fphys.2017.00806.
- Leddy, J. *et al.* (1997), 'Effect of a high or a low fat diet on cardiovascular risk factors in male and female runners', *Medicine & Science in Sports & Exercise*, 29(1), pp. 17–25.
- Leeds, A., Brand Miller, J., Foster-Powell, K. and Colagiuri, S. (2000), *The Glucose Revolution* (London: Hodder and Stoughton), p. 29.
- Leenders, M. *et al.* (2013), 'Elderly men and women benefit equally from prolonged resistance-type exercise training', *J. Gerontol. A. Biol. Sci. Med. Sci.*, vol. 68(7), pp. 769–79.
- Lehnen, T. E. *et al.* (2015), 'A review on effects of conjugated linoleic fatty acid (CLA) upon body composition and energetic metabolism', *J. Int. Soc. Sports Nutr.*, vol. 12, p. 36.
- Lemon, P. W. R. (1992), 'Protein requirements and muscle mass/strength changes during intensive training in novice bodybuilders', *J. Appl. Physiol.*, vol. 73, pp. 767–75.
- Lemon, P. W. R. (1995), 'Do athletes need more dietary protein and amino acids?', *Int. J. Sport Nutr.*, vol. 5 pp. s39–61.
- Lemon, P. W. R. (1998), 'Effects of exercise on dietary protein requirements', *Int. J. Sport Nutr.*, vol. 8, pp. 426–47.
- Lenn, J. *et al.* (2002), 'The effects of fish oil and isoflavones on delayed onset muscle soreness'. *Med. Sci. Sports Exerc.*, vol. 34(10), pp. 1605–13.
- Levine, S. A., Gordon, B. and Derick, C. L. (1924), 'Some changes in the chemical constituents of the blood following a marathon race: with special reference to the development of hypoglycemia', *Journal of the American Medical Association*, 82(22), pp. 1778–1779. doi: 10.1001/jama.1924.02650480034015.
- Levitsky, D. A. and Pacanowski, C. R. (2013), 'Effect of skipping breakfast on subsequent energy

- intake', *Physiol Behav.* vol. 119, pp. 9–16.
- Lewis, N. A. *et al.* (2020), 'Are There Benefits from the Use of Fish Oil Supplements in Athletes? A Systematic Review', *Advances in Nutrition (Bethesda, Md.)*, 11(5), pp. 1300–1314. doi: 10.1093/advances/nmaa050.
- LGC (2015), Clean Sport [http://www.informed-sport.com/sites/default/files/LGC\\_Clean\\_Sport\\_trifold\\_0315\\_EN\\_4336\\_digital2\\_0.pdf](http://www.informed-sport.com/sites/default/files/LGC_Clean_Sport_trifold_0315_EN_4336_digital2_0.pdf). Accessed March 2016.
- Li, Y., Hruby, A., Bernstein, A. M., *et al.* (2015), 'Saturated fats compared with unsaturated fats and sources of carbohydrates in relation to risk of coronary heart disease: a prospective cohort study', *J. Am. Coll. Cardiol.*, vol. 66(14), pp. 1538–48.
- Lis, D. *et al.* (2016), 'Food avoidance in athletes: FODMAP foods on the list', *Applied Physiology, Nutrition, and Metabolism = Physiologie Appliquee, Nutrition Et Metabolisme*, 41(9), pp. 1002–1004. doi: 10.1139/apnm-2015-0428.
- Lis, D. M. and Baar, K. (2019), 'Effects of Different Vitamin C-Enriched Collagen Derivatives on Collagen Synthesis', *International Journal of Sport Nutrition and Exercise Metabolism*, 29(5), pp. 526–531. doi: 10.1123/ijsnem.2018-0385.
- Lis, D. M. *et al.* (2018), 'Low FODMAP: A Preliminary Strategy to Reduce Gastrointestinal Distress in Athletes', *Medicine & Science in Sports & Exercise*, 50(1), pp. 116–123. doi: 10.1249/MSS.0000000000001419.
- Lissner, L. *et al.* (1987), 'Dietary fat and the regulation of energy intake in human subjects', *The American Journal of Clinical Nutrition*, 46(6), pp. 886–892. doi: 10.1093/ajcn/46.6.886.
- Liu, T.-H. *et al.* (2009), 'No effect of short-term arginine supplementation on nitric oxide production, metabolism and performance in intermittent exercise in athletes', *The Journal of Nutritional Biochemistry*, 20(6), pp. 462–468. doi: 10.1016/j.jnutbio.2008.05.005.
- Lloyd, T. *et al.* (1986), 'Women athletes with menstrual irregularity have increased musculoskeletal injuries', *Med. Sci. Sports Ex.*, vol. 18, pp. 3427–9.
- Logan-Sprenger, H. M. *et al.* (2015), 'The effect of dehydration on muscle metabolism and time trial performance during prolonged cycling in males', *Physiological Reports*, 3(8). doi: 10.14814/phy2.12483.
- Logue, D. *et al.* (2018), 'Low Energy Availability in Athletes: A Review of Prevalence, Dietary Patterns, Physiological Health, and Sports Performance', *Sports Medicine (Auckland, N.Z.)*, 48(1), pp. 73–96. doi: 10.1007/s40279-017-0790-3.
- Logue, D. M. *et al.* (2020), 'Low Energy Availability in Athletes 2020: An Updated Narrative Review of Prevalence, Risk, Within-Day Energy Balance, Knowledge, and Impact on Sports Performance', *Nutrients*, 12(3), p. 835. doi: 10.3390/nu12030835.
- Lohman, T. G. (1992), 'Basic concepts in body composition assessment', In *Advances in Body Composition Assessment*, Human Kinetics, pp. 109–118.
- Longland, T. M. *et al.* (2016), 'Higher compared with lower dietary protein during an energy deficit combined with intense exercise promotes greater lean mass gain and fat mass loss: a randomized trial', *Am. J. Clin. Nutr.* Jan 27 (Epub ahead of print).
- Loucks, A. B. (2003), 'Energy availability, not body fatness, regulates reproductive function in women', *Exerc. Sport Sci. Rev.*, vol. 31, pp. 144–148.
- Loucks, A. B. and Thuma, J. R. (2003), 'Luteinizing hormone pulsatility is disrupted at a threshold of energy availability in regularly menstruating women', *The Journal of Clinical Endocrinology and Metabolism*, 88(1), pp. 297–311. doi: 10.1210/jc.2002-020369.
- Loucks, A. B. *et al.* (1989), 'Alterations in the hypothalamic-pituitary-ovarian and the hypothalamic-pituitary axes in athletic women', *J. Clinical Endocrinol. Metab.*, vol. 68, pp. 402–22.

- Loucks, A. B., Kiens, B. and Wright, H. H. (2011), 'Energy availability in athletes', *Journal of Sports Sciences*, 29 Suppl 1, pp. S7-15. doi: 10.1080/02640414.2011.588958.
- Lovell, G. (2008), 'Vitamin D status of females in an elite gymnastic programme', *Clin. J. Sports Med.*, vol. 18, pp. 159–61.
- Luden, N. D. *et al.* (2007), 'Post-exercise carbohydrate-protein-antioxidant ingestion increase CK and muscle soreness in cross-country runners', *Int. J. Sports Exerc. Metab.*, vol. 17, pp. 109–122.
- Lukaski, H.C. (2004), 'Vitamin and mineral status: effects on physical performance'. *Nutrition*, vol. 20, 632–44.
- Lun, V. *et al.* (2012), 'Dietary supplementation practices in Canadian high-performance athletes', *Int. J. Sport Nutr. Exerc. Metab.*, vol. 22(1), pp. 31–7.
- Lynch, Heidi M., *et al.* (2016), "Cardiorespiratory Fitness and Peak Torque Differences between Vegetarian and Omnivore Endurance Athletes: A Cross-Sectional Study.", *Nutrients*, vol. 8, no. 11, Nov. 2016, p. 726. [www.mdpi.com](http://www.mdpi.com), doi:10.3390/nu8110726.
- MacConnie, S. E. *et al.* (1986), 'Decreased hypothalamic gonadotropin-releasing hormone secretion in male marathon runners', *The New England Journal of Medicine*, 315(7), pp. 411–417. doi: 10.1056/NEJM198608143150702.
- Mach, N. and Fuster-Botella, D. (2017), 'Endurance exercise and gut microbiota: A review', *Journal of Sport and Health Science*, 6(2), pp. 179–197. doi: 10.1016/j.jshs.2016.05.001.
- Macintosh, B. R. *et al.* (1995), 'Caffeine ingestion and performance of a 1500-metre swim'. *Can. J. Appl. Physiol.*, vol. 20 (2): pp. 168–77.
- Mackie, G. M., Samocha-Bonet, D. and Tam, C. S. (2017), 'Does weight cycling promote obesity and metabolic risk factors?', *Obesity Research & Clinical Practice*, 11(2), pp. 131–139. doi: 10.1016/j.orcp.2016.10.284.
- Macnaughton, L. S. *et al.* (2016), 'The response of muscle protein synthesis following whole-body resistance exercise is greater following 40 g than 20 g of ingested whey protein', *Physiol. Rep.* vol. 4 (15). pii: e12893.
- Mäestu, J. *et al.* (2010), 'Anabolic and catabolic hormones and energy balance of the male bodybuilders during the preparation for the competition', *Journal of Strength and Conditioning Research*, 24(4), pp. 1074–1081. doi: 10.1519/JSC.0b013e3181cb6fd3.
- Mäestu, J. *et al.* (2010), 'Anabolic and catabolic hormones and energy balance of the male bodybuilders during the preparation for the competition', *Journal of Strength and Conditioning Research*, 24(4), pp. 1074–1081. doi: 10.1519/JSC.0b013e3181cb6fd3.
- Maffucci, D. M. and McMurray, R. G. (2000), 'Towards optimising the timing of the pre-exercise meal', *Int. J. Sport Nutr.*, vol. 10, pp. 103–13.
- Maggini, S., Pierre, A. and Calder, P. C. (2018), 'Immune Function and Micronutrient Requirements Change over the Life Course', *Nutrients*, 10(10). doi: 10.3390/nu10101531.
- Mamerow, M. M. *et al.* (2014), 'Dietary Protein Distribution Positively Influences 24-h Muscle Protein Synthesis in Healthy Adults', *J. Nutrition*, vol. 144, pp. 876–80.
- Margolis, L. M. *et al.* (2021), 'Congestion of Carbohydrate and Protein on Muscle Glycogen Synthesis after Exercise: A Meta-analysis', *Medicine and Science in Sports and Exercise*, 53(2), pp. 384–393. doi: 10.1249/MSS.0000000000002476.
- Mariotti, FrançoisF., and Christopher D. Gardner (2019), "Dietary Protein and Amino Acids in Vegetarian Diets—A Review.", *Nutrients*, vol. 11, no. 11, Nov. 2019, p. 2661. [www.mdpi.com](http://www.mdpi.com), doi:10.3390/nu11112661.
- Marquet, L. A. *et al.* (2016) (a), 'Enhanced endurance performance by periodization of CHO intake: 'sleep low' strategy', *Med. Sci. Sports Exerc.*, vol. 48(4), pp. 663–72.
- Marquet, L.-A. *et al.* (2016) (b), 'Periodization of Carbohydrate Intake: Short-Term Effect on

- Performance', *Nutrients*, 8(12), p. 755. doi: 10.3390/nu8120755.
- Martin, W. F., Armstrong, L. E. and Rodriguez, N. R. (2005), 'Dietary protein intake and renal function', *Nutrition & Metabolism*, 2(1), p. 25. doi: 10.1186/1743-7075-2-25.
- Martinez, L. R. and Haymes, E. M. (1992), 'Substrate utilisation during treadmill running in prepubescent girls and women', *Med. Sci. Sports Exerc.*, vol. 24, pp. 975–83.
- Martinsen, M. *et al.* (2010), 'Dieting to win or to be thin? A study of dieting and disordered eating among adolescent elite athletes and non-athlete controls', *British Journal of Sports Medicine*, 44(1), pp. 70–76. doi: 10.1136/bjism.2009.068668.
- Mason, W. L. *et al.* (1993), 'Carbohydrate ingestion during exercise: liquid vs solid feedings', *Med. Sci. Sports Ex.*, vol. 25, pp. 966–9.
- Maughan, R. J. (1995), 'Creatine supplementation and exercise performance', *Int. J. Sport Nutr.*, vol. 5, pp. 94–101.
- Maughan, R. J. and Shirreffs, S. M. (2012), 'Nutrition for sports performance: issues and opportunities', *Proc. Nutr. Soc.*; vol. 71 (1), pp. 112–19.
- Maughan, R. J. *et al.* (1996), 'Rehydration and recovery after exercise', *Sports Sci. Ex.*, vol. 9(62), pp. 1–5.
- Maughan, R. J. *et al.* (2016), 'A randomized trial to assess the potential of different beverages to affect hydration status: development of a beverage hydration index', *Am J. Clin. Nutr.*, vol. 103, pp. 717–23.
- Maughan, R. J. *et al.* (2018), 'IOC consensus statement: dietary supplements and the high-performance athlete', *British Journal of Sports Medicine*, 52(7), pp. 439–455. doi: 10.1136/bjsports-2018-099027.
- Mayhew, D. L. *et al.* (2002), 'Effects of long term creatine supplementation on liver and kidney function in American Football players', *Int. J. Sport Nutr.*, vol. 12, pp. 453–60.
- McConnell, G. K. *et al.* (1997), 'Influence of ingested fluid volume on physiological responses during prolonged exercise', *Acta. Phys. Scand.*, vol. 160, pp. 149–56.
- McCubbin, A. J. *et al.* (2020), 'Hydrogel Carbohydrate-Electrolyte Beverage Does Not Improve Glucose Availability, Substrate Oxidation, Gastrointestinal Symptoms or Exercise Performance, Compared With a Concentration and Nutrient-Matched Placebo', *International Journal of Sport Nutrition and Exercise Metabolism*, 30(1), pp. 25–33. doi: 10.1123/ijsnem.2019-0090.
- McDonald, D. *et al.* (2018), 'American Gut: an Open Platform for Citizen Science Microbiome Research', *mSystems*, 3(3). doi: 10.1128/mSystems.00031-18.
- McGlory, C. *et al.* (2016), 'Fish oil supplementation suppresses resistance exercise and feeding-induced increases in anabolic signaling without affecting myofibrillar protein synthesis in young men', *Physiological Reports*, 4(6). doi: 10.14814/phy2.12715.
- McLeay, Y. *et al.* (2012), 'Effect of New Zealand blueberry consumption on recovery from eccentric exercise-induced muscle damage', *Journal of the International Society of Sports Nutrition*, 9(1), p. 19. doi: 10.1186/1550-2783-9-19.
- Mears, S. A. *et al.* (2020), 'Addition of sodium alginate and pectin to a carbohydrate-electrolyte solution does not influence substrate oxidation, gastrointestinal comfort, or cycling performance', *Applied Physiology, Nutrition, and Metabolism = Physiologie Appliquee, Nutrition Et Metabolisme*, 45(6), pp. 675–678. doi: 10.1139/apnm-2019-0802.
- Meier, C. *et al.* (2004), 'Supplementation with oral vitamin D and calcium during winter prevents seasonal bone loss: a randomised controlled open-label prospective trial', *J. Bone Mineral Res.*, vol. 19, pp. 1221–30.
- Melby, C. *et al.* (1993), 'Effect of acute resistance exercise on resting metabolic rate', *J. Appl. Physiol.*, vol. 75, pp. 1847–53.
- Melin, A. *et al.* (2015), 'Energy availability and the female athlete triad in elite endurance

- athletes', *Scandinavian Journal of Medicine & Science in Sports*, 25(5), pp. 610–622. doi: <https://doi.org/10.1111/sms.12261>.
- Melina, V., Craig, W. and Levin, S. (2016), 'Position of the Academy of Nutrition and Dietetics: Vegetarian Diets', *Journal of the Academy of Nutrition and Dietetics*, 116(12), pp. 1970–1980. doi: 10.1016/j.jand.2016.09.025.
- Mensink, R.P. *et al.* (2003), 'Effects of dietary fatty acids and carbohydrate on the ratio of serum total to HDL cholesterol and on serum lipids and apolipoproteins: a meta-analysis of 60 controlled trials', *American Journal of Clinical Nutrition*, 77 (5), pp. 1146–1155.
- Menzies, C. *et al.* (2020), 'Frequent Carbohydrate Ingestion Reduces Muscle Glycogen Depletion and Postpones Fatigue Relative to a Single Bolus', *International Journal of Sport Nutrition and Exercise Metabolism*, pp. 1–7. doi: 10.1123/ijsnem.2019-0291.
- Mercuro, G. *et al.* (2006), 'Impairment of physical exercise capacity in healthy postmenopausal women', *American Heart Journal*, 151(4), pp. 923–927. doi: 10.1016/j.ahj.2005.06.027.
- Mero, A. A. *et al.* (2010), 'Moderate energy restriction with high protein diet results in healthier outcome in women', *Journal of the International Society of Sports Nutrition*, 7(1), p. 4. doi: 10.1186/1550-2783-7-4.
- Messina, Mark, *et al.* (2018), "No Difference Between the Effects of Supplementing With Soya Protein Versus Animal Protein on Gains in Muscle Mass and Strength in Response to Resistance Exercise.", *International Journal of Sport Nutrition and Exercise Metabolism*, vol. 28, no. 6, Nov. 2018, pp. 674–85. [journals.humankinetics.com](http://journals.humankinetics.com), doi:10.1123/ijsnem.2018-0071.
- MHRA (2012), <http://www.mhra.gov.uk/home/groups/comms-po/documents/news/con174847.pdf>
- Middleton, N., Jelen, P. and Bell, G. (2004), 'Whole blood and mononuclear cell glutathione response to dietary whey protein supplementation in sedentary and trained male human subjects', *International journal of food sciences and nutrition*. doi: 10.1080/096374080410001666504.
- Miescher, E. and Fortney, S. M. (1989), 'Responses to dehydration and rehydration during heat exposure in young and older men', *Am. J. Physiol.*, vol. 257, pp. R1050–R1056.
- Mifflin, M. D. *et al.* (1990), 'A new predictive equation for resting energy expenditure in healthy individuals', *J. Am. Diet. Assoc.*, vol. 51, pp. 241–7.
- Mihic, S. *et al.* (2000), 'Acute creatine loading increases fat free mass but does not affect blood pressure, plasma creatinine or CK activity in men and women', *Med. Sci. Sport. Exerc.*, vol. 32, pp. 291–6.
- Miller, S. L. *et al.* (2002), 'Metabolic responses to provision of mixed protein-carbohydrate supplementation during endurance exercise attenuate the increases in cortisol', *Int. J. Sport Nutr.*, vol. 12, pp. 384–97.
- Minehan, M. R. *et al.* (2002), 'Effect of flavour and awareness of kilojoule content of drinks on preference and fluid balance in team sports', *Int. J. Sport Nutr.*, vol. 12, pp. 81–92.
- Minetto, M. A. *et al.* (2013), 'Origin and development of muscle cramps', *Exercise and Sport Sciences Reviews*, 41(1), pp. 3–10. doi: 10.1097/JES.0b013e3182724817.
- Misra, M. *et al.* (2004), 'Effects of anorexia nervosa on clinical, hematologic, biochemical, and bone density parameters in community-dwelling adolescent girls', *Pediatrics*, 114(6), pp. 1574–1583. doi: 10.1542/peds.2004-0540.
- Mitchell, C. M. *et al.* (2019), 'Does Exercise Alter Gut Microbial Composition? A Systematic Review', *Medicine and Science in Sports and Exercise*, 51(1), pp. 160–167. doi: 10.1249/MSS.0000000000001760.
- Mitchell, W. K. *et al.* (2012), 'Sarcopenia, dynapenia, and the impact of advancing age on human



- skeletal muscle size and strength; a quantitative review', *Front Physiol.* vol. 3, p. 260.
- Mizumura, K. and Taguchi, T. (2016) (b), 'Delayed onset muscle soreness: Involvement of neurotrophic factors', *The journal of physiological sciences: JPS*, 66(1), pp. 43–52. doi: 10.1007/s12576-015-0397-0.
- Mohr, A. E. *et al.* (2020), 'The athletic gut microbiota', *Journal of the International Society of Sports Nutrition*, 17(1), p. 24. doi: 10.1186/s12970-020-00353-w.
- Montain, S. J. and Coyle, E. F. (1992), 'The influence of graded dehydration on hyperthermia and cardiovascular drift during exercise', *J. Appl. Physiol.*, vol. 73, pp. 1340–50.
- Moore, D. R. (2019), 'Maximizing Post-exercise Anabolism: The Case for Relative Protein Intakes', *Frontiers in Nutrition*, 6. doi: 10.3389/fnut.2019.00147.
- Moore, D. R. *et al.* (2009), 'Ingested protein dose response of muscle and albumin protein synthesis after existence exercise in young men', *Am. J. Clin. Nutr.*, vol. 89, pp. 161–8.
- Moore, D. R. *et al.* (2014), 'Beyond muscle hypertrophy: why dietary protein is important for endurance athletes', *Appl. Physiol. Nutr. Metab.*, vol 39(9), pp. 987–97
- Moore, D. R. *et al.* (2015), 'Protein ingestion to stimulate myofibrillar protein synthesis requires greater relative protein intakes in healthy older versus younger men', *J. Gerontol. A. Biol. Sci. Med. Sci.* vol. 70(1), pp. 57–62.
- Morehen, J. C. *et al.* (2020), 'Montmorency tart cherry juice does not reduce markers of muscle soreness, function and inflammation following professional male rugby League match-play', *European Journal of Sport Science*, pp. 1–10. doi: 10.1080/17461391.2020.1797181.
- Morrison, L. J. *et al.* (2004), 'Prevalent use of dietary supplements among people who exercise at a commercial gym'. *Int. J. Sport Nutr. Exerc. Metab.*, vol. 14, pp. 481–92.
- Morton, J. P. *et al.* (2009), 'Reduced carbohydrate availability does not modulate training-induced heat shock protein adaptations but does upregulate oxidative enzyme activity in human skeletal muscle', *Journal of Applied Physiology (Bethesda, Md.: 1985)*, 106(5), pp. 1513–1521. doi: 10.1152/jappphysiol.00003.2009.
- Morton, R. W. *et al.* (2018), 'A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults', *British Journal of Sports Medicine*, 52(6), pp. 376–384. doi: 10.1136/bjsports-2017-097608.
- Mountjoy, M. *et al.* (2014), 'The IOC consensus statement: beyond the Female Athlete Triad--Relative Energy Deficiency in Sport (RED-S)', *British Journal of Sports Medicine*, 48(7), pp. 491–497. doi: 10.1136/bjsports-2014-093502.
- Mountjoy, M. *et al.* (2015), 'Authors' 2015 additions to the IOC consensus statement: Relative Energy Deficiency in Sport (RED-S)', *British Journal of Sports Medicine*, 49(7), pp. 417–420. doi: 10.1136/bjsports-2014-094371.
- Mountjoy, M. *et al.* (2018), 'IOC consensus statement on relative energy deficiency in sport (RED-S): 2018 update', *British Journal of Sports Medicine*, 52(11), pp. 687–697. doi: 10.1136/bjsports-2018-099193.
- Mozaffarian, D. *et al.* (2006), 'Trans fatty acids and cardiovascular disease', *New Eng. J. Med.*, vol. 354, pp. 1601–13.
- Mozaffarian, D. *et al.* (2011), muls 'Components of a cardioprotective diet: new insights', *Circulation*, vol. 123, pp. 2870–91.
- Mullins, V. A. *et al.* (2001), 'Nutritional status of US elite female heptathletes during training', *Int. J. Sport Nutr.*, vol. 11, pp. 299–314.
- Muls, E. *et al.* (1995), 'Is weight cycling detrimental to health? A review of the literature in humans', *International Journal of Obesity and Related Metabolic Disorders: Journal of the International Association for the Study of Obesity*, 19 Suppl 3, pp. S46-50.

- Muoio, D. M. *et al.* (1994), 'Effect of dietary fat on metabolic adjustments to maximal VO<sub>2</sub> and endurance in runners', *Med. Sci. Sports Exerc.*, vol. 26(1), pp. 81–8.
- Murphy, C. H. *et al.* (2015), 'Considerations for protein intake in managing weight loss in athletes', *Eur. J. Sport Sci.*, vol 15(1), pp. 21–8.
- Murphy, M. *et al.* (2012), 'Whole beetroot consumption acutely improves running performance', *J. Acad. Nutr. Diet.*, vol. 112, pp. 548–52.
- Murray, R. *et al.* (1999), 'A comparison of the gastric emptying characteristics of selected sports drinks', *Int. J. Sports Nutr.*, vol. 9, pp. 263–74.
- Nattiv, A. *et al.* (2007), 'American College of Sports Medicine position stand. The female athlete triad', *Med. Sci. Sports Exerc.*, vol. 39, pp. 1867–82.
- Nebl, Josefine, *et al.* (2019), 'Exercise Capacity of Vegan, Lacto-Ovo-Vegetarian and Omnivorous Recreational Runners.', *Journal of the International Society of Sports Nutrition*, vol. 16, no. 1, May 2019, p. 23. BioMed Central, doi:10.1186/s12970-019-0289-4.
- Needleman I *et al.* (2014), 'Oral health and impact on performance of athletes participating in the London 2012 Olympic Games: a cross-sectional study', *Br J Sports Med.*, vol. 47(16), pp. 1054–8.
- Needleman, I. *et al.* (2015), 'Oral health and elite sport performance', *Br. J. Sports Med.*, vol. 49, pp. 3–6.
- Nelson, A. *et al.* (1997), 'Creatine supplementation raises anaerobic threshold', *FASEB J.*, vol. 11, A586 (abstract).
- Nelson, M. E. *et al.* (1986), 'Diet and bone status in amenorrheic runners', *Am. J. Clin. Nutr.*, vol. 43, pp. 910–16.
- Neufer, P. D. *et al.* (1987), 'Improvements in exercise performance: effects of carbohydrate feedings and diet', *J. Appl. Physiol.*, vol. 62, pp. 983–8.
- Neychev, V. K. and Mitev VI. (2005), 'The aphrodisiac herb *Tribulus terrestris* does not influence the androgen production in young men', *J. Ethnopharmacol.*, vol. 101(1–3), pp. 319–20.
- Nichols, J. F. *et al.* (2007), 'Disordered eating and menstrual irregularity in high school athletes in lean-build and non lean-build sports', *Int. J. Sport Nutr. Exerc. Metab.*, vol. 17, pp. 364–77.
- Nielsen, F. H. and Lukaski, H. C. (2006), 'Update on the relationship between magnesium and exercise', *Magnesium Research*, 19(3), pp. 180–189.
- Nieman, D. C. (1999), 'Physical fitness and vegetarian diets: is there a relation?', *The American Journal of Clinical Nutrition*, 70(3), pp. 570s–575s. doi: 10.1093/ajcn/70.3.570s.
- Nieman, D. C. (1999), 'Physical fitness and vegetarian diets: is there a relation?' *Am. J. Clin. Nutr.*, (Sep) vol. 70 (3 suppl), pp. 570S–5S.
- Nieman, D. C. *et al.* (1989), 'Hematological, anthropometric, and metabolic comparisons between vegetarian and nonvegetarian elderly women'. *Int. J. Sports Med.*, vol. 10, pp. 243–50.
- Nieman, D. C. *et al.* (2007), 'Quercetin reduces illness but not immune perturbations after intense exercise', *Med. Sci. Sports Exerc.*, vol. 39, pp. 1561–69.
- Nieman, D. C., Henson, D. A., McAnulty, S. R., *et al.* (2004), 'Vitamin E and immunity after the Kona Triathlon World Championship', *Med. Sci. Sports Exerc.*, Aug 36(8), pp. 1328–35.
- Noakes, T. D. (1993), 'Fluid replacement during exercise', *Exerc. Sport Sci. Rev.*, vol. 21, pp. 297–330.
- Noakes, T. D. (2000), 'Hyponatremia in distance athletes: pulling the IV on the dehydration myth', *Phys. Sportsmed.*, vol. 26 (Sept), pp. 71–6.
- Noakes, T. D. (2007), 'Drinking guidelines for exercise: what evidence is there that athletes should drink as much as possible to replace the weight lost during exercise or ad libitum?', *J. Sports Sci.*, vol. 25 (7), pp. 781–96.

- Noakes, T. D. (2010), 'Is drinking to thirst optimum?' *Ann. Nutr. Metab.*, vol, 57, Suppl 2, pp. S9–17.
- Noakes, T. D. (2012), *Waterlogged: The Serious Problem of Overhydration in Endurance Sports*, (Champaign, IL:/Human Kinetics).
- Noreen, E. E. *et al.* (2010), 'Effects of supplemental fish oil on resting metabolic rate, body composition, and salivary cortisol in healthy adults'. *J. Int. Soc. Sports Nutr.*, vol. 7, p. 31.
- Norris, M. L. *et al.* (2016), 'Gastrointestinal complications associated with anorexia nervosa: A systematic review', *International Journal of Eating Disorders*, 49(3), pp. 216–237. doi: <https://doi.org/10.1002/eat.22462>.
- Norton, L. E. and Layman, D. K. (2006), 'Leucine Regulates Translation Initiation of Protein Synthesis in Skeletal Muscle after Exercise', *The Journal of Nutrition*, 136(2), pp. 533S-537S. doi: 10.1093/jn/136.2.533S.
- Nosaka, K. *et al.* (2006), 'Effects of amino acid supplementation on muscle soreness and damage'. *Int. J. Sports Nutr. Exerc. Metab.*, vol. 16, pp. 620–35.
- Nyakayiru, J. *et al.* (2017), 'Beetroot Juice Supplementation Improves High-Intensity Intermittent Type Exercise Performance in Trained Soccer Players', *Nutrients*, 9(3). doi: 10.3390/nu9030314.
- O'Donovan, C. M. *et al.* (2020), 'Distinct microbiome composition and metabolome exists across subgroups of elite Irish athletes', *Journal of Science and Medicine in Sport*, 23(1), pp. 63–68. doi: 10.1016/j.jsams.2019.08.290.
- Oliver, J. M., Anzalone, A. J. and Turner, S. M. (2018), 'Protection Before Impact: the Potential Neuroprotective Role of Nutritional Supplementation in Sports-Related Head Trauma', *Sports Medicine (Auckland, N.Z.)*, 48(Suppl 1), pp. 39–52. doi: 10.1007/s40279-017-0847-3.
- Onywere, V. O. *et al.* (2004), 'Food and macronutrient intake of elite kenyan distance runners'. *Int. J. Sport Nutr. Exerc. Metab.*, vol. 14(6), pp. 709–19.
- Ostendorf, D. M. *et al.* (2018), 'No consistent evidence of a disproportionately low resting energy expenditure in long-term successful weight-loss maintainers', *The American Journal of Clinical Nutrition*, 108(4), pp. 658–666. doi: 10.1093/ajcn/nqy179.
- Ostojic, S. (2004), 'Creatine supplementation in young soccer players'. *Int. J. Sport Nutr. Exerc. Metab.*, vol. 14, pp. 95–103.
- Otis, C. L. *et al.* (1997), 'American College of *Sports Medicine* position stand. The Female Athlete Triad'. *Med. Sci. Sport Exerc.*, vol. 29, pp. i–ix.
- Otte, J. A. *et al.* (2001), 'Exercise induces gastric ischemia in healthy volunteers: a tonometry study', *Journal of Applied Physiology (Bethesda, Md.: 1985)*, 91(2), pp. 866–871. doi: 10.1152/jap.2001.91.2.866.
- Owens, B. M. (2007), 'The potential effects of pH and buffering capacity on dental erosion'. *Gen. Dent.*, vol. 55(6), pp. 527–31.
- Owens, D. J., Allison, R. and Close, G. L. (2018), 'Vitamin D and the Athlete: Current Perspectives and New Challenges', *Sports Medicine*, 48(1), pp. 3–16. doi: 10.1007/s40279-017-0841-9.
- Owens, D. J., Fraser, W. D. and Close, G. L. (2015), 'Vitamin D and the athlete: emerging insights', *European Journal of Sport Science*, 15(1), pp. 73–84. doi: 10.1080/17461391.2014.944223.
- Paddon-Jones, D. *et al.* (2001), 'Short term HMB supplementation does not reduce symptoms of eccentric muscle damage'. *Int. J. Sport Nutr.*, vol. 11, pp. 442–50.
- Pagoto, S. L. and Appelhans, B. M. (2013), 'A call for an end to the diet debates'. *JAMA*, vol 310(7), pp. 687–688.
- Palmer-Green, D. *et al.* (2013), 'The Injury/Illness Performance Project (IIPP): A Novel

- Epidemiological Approach for Recording the Consequences of Sports Injuries and Illnesses', *Journal of Sports Medicine*, 2013, pp. 1–9. doi: 10.1155/2013/523974.
- Pannemans, D. L. *et al.* (1997), 'Calcium excretion, apparent calcium absorption and calcium balance in young and elderly subjects: influence of protein intake'. *Brit. J. Nutr.*, vol. 77(5), pp. 721–9.
- Papageorgiou, M. *et al.* (2018), 'Reduced energy availability: implications for bone health in physically active populations', *European Journal of Nutrition*, 57(3), pp. 847–859. doi: 10.1007/s00394-017-1498-8.
- Parr, E. B. *et al.* (2014), 'Alcohol ingestion impairs maximal post-exercise rates of myofibrillar protein synthesis following a single bout of concurrent training', *PLoS One*, 9(2), p. e88384. doi: 10.1371/journal.pone.0088384.
- Parry-Billings, M. *et al.* (1992), 'Plasma amino acid concentrations in over-training syndrome: possible effects on the immune system'. *Med. Sci. Sport Ex.*, vol. 24, pp. 1353–8.
- Pasiakos S. M. *et al.* (2013), 'Effects of high-protein diets on fat-free mass and muscle protein synthesis following weight loss: a randomized controlled trial'. *FASEB J.*, vol. 27(9), pp.3837–47.
- Pasiakos, S. M, and McClung, J. P. (2011), 'Supplemental dietary leucine and the skeletal muscle anabolic response to essential amino acids'. *Nutr. Rev.*, vol. 69(9), pp. 550–57.
- Pasiakos, S. M. *et al.* (2010), 'Acute energy deprivation affects skeletal muscle protein synthesis and associated intracellular signaling proteins in physically active adults', *The Journal of Nutrition*, 140(4), pp. 745–751. doi: 10.3945/jn.109.118372.
- Pasiakos, S. M., McClung, H. L., McClung, J. P. (2011), 'Leucine-enriched essential amino acid supplementation during moderate steady state exercise enhances post-exercise muscle protein synthesis'. *Am. J. Clin. Nutr.*, vol. 94(3), pp. 809–18.
- Passricha, S. R. *et al.* (2014), 'Iron Supplementation Benefits Physical Performance in Women of Reproductive Age: A Systematic Review and Meta-Analysis'. *J. Nutr.* 2014 jn.113.189589; first published online April 9, 2014.
- Passé, D. H. *et al.* (2004), 'Palatability and voluntary intake of sports beverages, diluted orange juice and water during exercise'. *Int. J. Sport Nutr. Exerc. Metab.*, vol. 14, pp. 272–84.
- Patterson, S. D. and Gray, S. C. (2007), 'Carbohydrate-gel supplementation and endurance performance during intermittent high-intensity shuttle running'. *Int. J. Sports Nutr. Exerc. Metab.*, vol. 17, pp. 445–55.
- Paulsen, G., Cumming, K. T. *et al.* (2013), 'Vitamin C and E supplementation hampers cellular adaptation to endurance training in humans: a double-blind randomized controlled trial'. *J. Physiol.* 2013.267419
- Peeling, P. *et al.* (2008), 'Athletic induced iron deficiency: new insights into the role of inflammation, cytokines and hormones', *European Journal of Applied Physiology*, 103(4), pp. 381–391. doi: 10.1007/s00421-008-0726-6.
- Pennings, B. *et al.* (2011), 'Exercising before protein intake allows for greater use of dietary protein-derived amino acids for de novo muscle protein synthesis in both young and elderly men'. *Am. J. Clin. Nutr.*, vol. 93, pp. 322–31.
- Pennings, B. *et al.* (2012), 'Amino acid absorption and subsequent muscle protein accretion following graded intakes of whey protein in elderly men'. *Am. J. Physiol. Endocrinol. Metab.*, vol. 302, pp. E992–9.
- Peternejl, T. T. and Coombes, J. S. (2011), 'Antioxidant supplementation during exercise training: beneficial or detrimental?' *Sports Med.*, vol. 41(12), pp. 1043–69.
- Peters, E. M. *et al.* (1993), 'Vitamin C supplementation reduces the incidence of post-race symptoms of upper-respiratory-tract infection in ultra-marathon runners'. *Am. J. Clin. Nutr.*,

- vol. 57, pp. 170–4.
- Peters, E. M. *et al.* (2001), ‘Vitamin C supplementation attenuates the increases in circulating cortisol, adrenaline and anti-inflammatory polypeptides following ultra-marathon running’. *Int. J. Sports Med.*, vol. 22(7), pp. 537–43.
- Petersen, L. M. *et al.* (2017), ‘Community characteristics of the gut microbiomes of competitive cyclists’, *Microbiome*, 5. doi: 10.1186/s40168-017-0320-4.
- Petersen, L. M. *et al.* (2017), ‘Community characteristics of the gut microbiomes of competitive cyclists’, *Microbiome*, 5, p. 98. doi: 10.1186/s40168-017-0320-4.
- Peterson, M. *et al.* (2011), ‘Influence of resistance exercise on lean body mass in aging adults: a metaanalysis’. *Med. Sci. Sports Exerc.*, vol. 43, pp. 249–58.
- Petkus, D. L., Murray-Kolb, L. E. and De Souza, M. J. (2017), ‘The Unexplored Crossroads of the Female Athlete Triad and Iron Deficiency: A Narrative Review’, *Sports Medicine*, 47(9), pp. 1721–1737. doi: 10.1007/s40279-017-0706-2.
- Pfeiffer, B. *et al.* (2010a) (a), ‘Oxidation of solid versus liquid CHO sources during exercise’. *Med. Sci. Sports Exerc.*, vol. 42(11), pp. 2030–7.
- Pfeiffer, B. *et al.* (2010b) (b), ‘CHO oxidation from a CHO gel compared with a drink during exercise’. *Med. Sci. Sports Exerc.*, vol. 42(11), pp. 2038–45.
- Phillips, P. A. *et al.* (1984), ‘Reduced thirst after water deprivation in healthy elderly men’. *N. Engl. J. Med.*, vol. 311, pp. 753–59.
- Phillips, S. M. (2012), ‘Dietary protein requirements and adaptive advantages in athletes’. *Br J. Nutr.*, vol. 108, Suppl. 2, pp. S158–67.
- Phillips, S. M. and Van Loon, L. J. C. (2011) (a), ‘Dietary protein for athletes: from requirements to optimum adaptation’, *Journal of Sports Sciences*, 29 Suppl 1, pp. S29–38. doi: 10.1080/02640414.2011.619204.
- Phillips, S. M. and Van Loon, L. J. C. (2011) (b), ‘Dietary protein for athletes: from requirements to optimum adaptation’, *Journal of Sports Sciences*, 29 Suppl 1, pp. S29–38. doi: 10.1080/02640414.2011.619204.
- Phillips, S. M. *et al.* (1997), ‘Mixed muscle protein synthesis and breakdown after resistance training in humans’. *Am. J. Physiol.*, vol. 273(1), pp. E99–E107.
- Phillips, S. M. *et al.* (1999), ‘Resistance training reduces acute exercise-induced increase in muscle protein turnover’. *Am. J. Physiol.*, vol. 276(1), pp. E118–24.
- Phillips, S. M. *et al.* (2005), ‘Dietary protein to support anabolism with resistance exercise in young men’. *J. Am. Coll. Nutr.*, vol. 24(2), pp. 134S–9S.
- Phillips, S. M. *et al.* (2007), ‘A critical examination of dietary protein requirements, benefits and excesses in athletes’. *Int. J. Sports Nutr. Exerc. Metab.*, vol. 17, pp. 58–78.
- Phillips, S. M., Chevalier, S. and Leidy, H. J. (2016), ‘Protein “requirements” beyond the RDA: implications for optimizing health’, *Applied Physiology, Nutrition, and Metabolism = Physiologie Appliquee, Nutrition Et Metabolisme*, 41(5), pp. 565–572. doi: 10.1139/apnm-2015-0550.
- Phillips, S. M., *et al.* (2011)(c), ‘Nutrition for weight and resistance training’. In: Lanham-New, S. A. *et al.* (eds), *Sport and Exercise Nutrition* (Oxford: Wiley-Blackwell).
- Phillips, T. *et al.* (2003), ‘A dietary supplement attenuates IL-6 and CRP after eccentric exercise in untrained males’. *Med. Sci. Sports Exerc.*, vol. 35(12), pp. 2032–7.
- Philpott, J. D., Witard, O. C. and Galloway, S. D. R. (2019), ‘Applications of omega-3 polyunsaturated fatty acid supplementation for sport performance’, *Research in Sports Medicine (Print)*, 27(2), pp. 219–237. doi: 10.1080/15438627.2018.1550401.
- Phinney, S. D. *et al.* (1983), ‘The human metabolic response to chronic ketosis without caloric restriction: preservation of submaximal exercise capability with reduced carbohydrate oxidation’. *Metabolism*, vol. 32(8), pp. 69–76.

- Pilegaard, H. *et al.* (2005), 'Substrate availability and transcriptional regulation of metabolic genes in human skeletal muscle during recovery from exercise'. *Metabolism*, vol. 54(8), pp. 1048–55.
- Pillay, S. *et al.* (2021), 'Longer race distance predicts gastrointestinal illness-related medical encounters in 153 208 endurance runner race starters – SAFER XVI', *The Journal of Sports Medicine and Physical Fitness*. doi: 10.23736/S0022-4707.21.12072-9.
- Pinto, C. L. *et al.* (2016), 'Impact of creatine supplementation in combination with resistance training on lean mass in the elderly', *Journal of Cachexia, Sarcopenia and Muscle*, 7(4), pp. 413–421. doi: <https://doi.org/10.1002/jcsm.12094>.
- Plotkin, D. L. *et al.* (2021), 'Isolated Leucine and Branched-Chain Amino Acid Supplementation for Enhancing Muscular Strength and Hypertrophy: A Narrative Review', *International Journal of Sport Nutrition and Exercise Metabolism*, pp. 1–10. doi: 10.1123/ijsnem.2020-0356.
- Pöschmüller, M. *et al.* (2016), 'A systematic review and meta-analysis of carbohydrate benefits associated with randomized controlled competition-based performance trials'. *J. Int. Soc. Sports Nutr.* vVol. 13, pp 27.
- Poffé, C. *et al.* (2020), 'Exogenous ketosis impacts neither performance nor muscle glycogen breakdown in prolonged endurance exercise', *Journal of Applied Physiology (Bethesda, Md.: 1985)*, 128(6), pp. 1643–1653. doi: 10.1152/jappphysiol.00092.2020.
- Polidori, D. *et al.* (2016), 'How Strongly Does Appetite Counter Weight Loss? Quantification of the Feedback Control of Human Energy Intake', *Obesity (Silver Spring, Md.)*, 24(11), pp. 2289–2295. doi: 10.1002/oby.21653.
- Pollock, M. L. and Jackson, A. S. (1984), 'Research progress invalidation of clinical methods of assessing body composition'. *Med. Sci. Sport Ex.*, vol. 16, pp. 606–13.
- Poortmans, J. R. and Francaux, M. (1999), 'Longterm oral creatine supplementation does not impair renal function in healthy athletes'. *Med. Sci. Sports Ex.*, vol. 31(8), pp. 1103–10.
- Pottier, A. *et al.* (2010), 'Mouth rinse but not ingestion of a carbohydrate solution improves 1-h cycle time trialtime-trial performance', *Scand. J. Med. Sci. Sports*, vol. 20(1), pp. 105–11.
- Powers, M. E. (2002), 'The safety and efficacy of anabolic steroid precursors: what is the scientific evidence?'. *J. Athol. Training*, vol. 37(3), pp. 300–5.
- Powers, S. *et al.* (2011), 'Antioxidant and vitamin D supplements for athletes: sense or nonsense?' *J. Sports Sci.*, vol. 29, Suppl. 1, S47–55.
- Public Health England (2011), 'Government Dietary Recommendation's.
- Public Health England (2016), 'The Eatwell Guide'.. Available at: <https://www.gov.uk/government/publications/the-eatwell-guide> (Accessed: 30 April 2021).
- Pugh, J. N. *et al.* (2018), 'Prevalence, Severity and Potential Nutritional Causes of Gastrointestinal Symptoms during a Marathon in Recreational Runners', *Nutrients*, 10(7). doi: 10.3390/nu10070811.
- Pugh, J. N. *et al.* (2019), 'Four weeks of probiotic supplementation reduces GI symptoms during a marathon race', *European Journal of Applied Physiology*, 119(7), pp. 1491–1501. doi: 10.1007/s00421-019-04136-3.
- Pyne, D. B. *et al.* (2015), 'Probiotics supplementation for athletes – clinical and physiological effects', *European Journal of Sport Science*, 15(1), pp. 63–72. doi: 10.1080/17461391.2014.971879.
- Quesnele, J. J. *et al.* (2014), 'The effects of Beta-alanine supplementation on performance: a systematic review of the literature'. *Int. J. Sport Nutr. Exerc. Metab.*, vol. 24(1), pp. 14–27.
- Ravussin, E. *et al.* (2019), 'Early Time-Restricted Feeding Reduces Appetite and Increases Fat Oxidation But Does Not Affect Energy Expenditure in Humans', *Obesity (Silver Spring, Md.)*,

- 27(8), pp. 1244–1254. doi: 10.1002/oby.22518.
- Rawson, E. S. and Volek, J. S. (2003), 'Effects of creatine supplementation and resistance training on muscle strength and weightlifting performance'. *J. Strength Cond. Res.*, vol. 17, pp. 822–31.
- Ready, S. L. *et al.* (1999), 'The effect of two sports drink formulations on muscle stress and performance'. *Med. Sci. Sports Exerc.*, vol. 31(5), p. S119.
- Reidy, P. T. *et al.* (2013), 'Protein blend ingestion following resistance exercise promotes human muscle protein synthesis'. *J. Nutr.*, vol. 143(4), pp. 410–16.
- Rennie, M. J. (2009), 'Anabolic resistance: the effects of aging, sexual dimorphism, and immobilization on human muscle protein turnover'. *Appl. Physiol. Nutr. Metab.*, vol. 34, pp. 377–81.
- Rennie, M. J. and Tipton, K. D. (2000), 'Protein and amino acid metabolism during and after exercise and the effects of nutrition', *Annual Review of Nutrition*, 20, pp. 457–483. doi: 10.1146/annurev.nutr.20.1.457.
- Res, P. T. *et al.* (2012), 'Protein ingestion prior to sleep improves post-exercise overnight recovery'. *Med. Sci. Sports Exerc.*, Feb 9 (Epub ahead of print).
- Reubinoff, B. E. *et al.* (1995), 'Effects of hormone replacement therapy on weight, body composition, fat distribution, and food intake in early post-menopausal women: a prospective study', *Fertility and Sterility*, 64(5), pp. 963–968. doi: 10.1016/s0015-0282(16)57910-2.
- Ribeiro, A. S. *et al.* (2016), 'Effect of conjugated linoleic acid associated with aerobic exercise on body fat and lipid profile in obese women: a randomized, double-blinded, and placebo-controlled trial'. *Int. J. Sport Nutr. Exerc. Metab.*, vol. 26(2), pp. 135–44.
- Richter, E. A. *et al.* (1991), 'Immune parameters in male athletes after a lacto-ovo-vegetarian diet and a mixed Western diet'. *Med. Sci. Sports Exerc.*, vol. 23, pp. 517–21.
- Rickenlund, A. *et al.* (2005), 'Amenorrhea in Female Athletes Is Associated with Endothelial Dysfunction and Unfavorable Lipid Profile', *The Journal of Clinical Endocrinology & Metabolism*, 90(3), pp. 1354–1359. doi: 10.1210/jc.2004-1286.
- Ritz, P and Rockwell, M (2021), *Promoting Optimal Omega-3 Fatty Acid Status in Athletes Gatorade Sports Science Institute*. Available at: <http://www.gssi-web.org:80/en/sports-science-exchange/Article/promoting-optimal-omega-3-fatty-acid-status-in-athletes> (Accessed: 13 July 2021).
- Ritz, P. P. *et al.* (2020), 'Dietary and Biological Assessment of the Omega-3 Status of Collegiate Athletes: A Cross-Sectional Analysis', *PloS One*, 15(4), p. e0228834. doi: 10.1371/journal.pone.0228834.
- Rivera-Brown, A. M. *et al.* (1999), 'Drink composition, voluntary drinking and fluid balance in exercising trained heat-acclimatized boys'. *J. Appl. Physiol.*, vol. 86, pp. 78–84.
- Rizkalla, S. *et al.* (2004), 'Improved plasma glucose control, whole body glucose utilisation and lipid profile on a low glycaemic index diet in type 2 diabetic men: a randomised controlled trial'. *Diab. Care*, vol. 27, pp. 1866–72.
- Roberts, J. D. *et al.* (2016), 'An Exploratory Investigation of Endotoxin Levels in Novice Long Distance Triathletes, and the Effects of a Multi-Strain Probiotic/Prebiotic, Antioxidant Intervention', *Nutrients*, 8(11). doi: 10.3390/nu8110733.
- Robinson. T. M. *et al.* (2000), 'Dietary creatine supplementation does not affect some haematological indices, or indices of muscle damage and hepatic and renal function'. *Brit. J. Sports Med.*, vol. 34, pp. 284–8.
- Rodriguez, N. R., Di Marco, N. M., Langley, S., *et al.* (2009), 'American College of Sports Medicine position stand: Nutrition and athletic performance', *Med. Sci. Sports Exerc.*, vol. 41(3), pp. 709–31.

- Rogers, J. *et al.* (2005), 'Gastric emptying and intestinal absorption of a low carbohydrate sport drink during exercise'. *Int. J. Sport Nutr. Exerc. Metab.*, vol. 15, pp. 220–35.
- Rogers, M. A. *et al.* (2021), 'Prevalence of impaired physiological function consistent with Relative Energy Deficiency in Sport (RED-S): an Australian elite and pre-elite cohort', *British Journal of Sports Medicine*, 55(1), pp. 38–45. doi: 10.1136/bjsports-2019-101517.
- Rogerson, S. *et al.* (2007), 'The effect of five weeks of Tribulus terrestris supplementation on muscle strength and body composition during preseason training in elite rugby league players'. *J. Strength Cond. Res.*, vol. 21(2), pp. 348–53.
- Rokitzki, L. *et al.* (1994), 'a-tocopherol supplementation in racing cyclists during extreme endurance training'. *Int. J. Sports Nutr.*, vol. 4, pp. 235–64.
- Rollo, I. *et al.* (2008), 'The influence of carbohydrate mouth rinse on self-selected speeds during a 30-min treadmill run', *Int. J. Sport Nutr. Exerc. Metab.*, vol. 18(6), pp. 585–600.
- Rolls, B. J. and Shide, D. J. (1992), 'The influence of fat on food intake and body weight'. *Nutr. Revs.*, vol. 50(10), pp. 283–90.
- Romano-Ely, B. C. *et al.* (2006), 'Effects of an isocaloric carbohydrate-protein-antioxidant drink on cycling performance'. *Med. Sci. Sports Exerc.*, vol. 38, pp. 1608–16.
- Rosen, L. W. *et al.* (1986), 'Pathogenic weightcontrol behavior in female athletes'. *Phys. Sports Med.*, vol. 14, pp. 79–86.
- Rosenbaum, M. and Leibel, R. L. (2010), 'Adaptive thermogenesis in humans', *International Journal of Obesity (2005)*, 34 Suppl 1, pp. S47-55. doi: 10.1038/ijo.2010.184.
- Rossow, L. M. *et al.* (2013), 'Natural bodybuilding competition preparation and recovery: a 12-month case study', *International Journal of Sports Physiology and Performance*, 8(5), pp. 582–592. doi: 10.1123/ijsp.8.5.582.
- Rowbottom, D. G. *et al.* (1996), 'The energizing role of glutamine as an indicator of exercise stress and overtraining'. *Sports Med.*, vol. 21(2), pp. 80–97.
- Ruddick-Collins, L. C., Morgan, P. J. and Johnstone, A. M. (2020), 'Mealtime: A circadian disruptor and determinant of energy balance?', *Journal of Neuroendocrinology*, 32(7), p. e12886. doi: <https://doi.org/10.1111/jne.12886>.
- Rustad, P.I. *et al.* (2016), 'Intake of Protein Plus Carbohydrate during the First Two Hours after Exhaustive Cycling Improves Performance the following Day'. *PLoS One*, vol. 11(4) e0153229.
- Rutherford, Shane M., *et al.* (2015), "Protein Digestibility-Corrected Amino Acid Scores and Digestible Indispensable Amino Acid Scores Differentially Describe Protein Quality in Growing Male Rats.", *The Journal of Nutrition*, vol. 145, no. 2, Feb. 2015, pp. 372–79. PubMed, doi:10.3945/jn.114.195438.
- Rynders, C. A. *et al.* (2019), 'Effectiveness of Intermittent Fasting and Time-Restricted Feeding Compared to Continuous Energy Restriction for Weight Loss', *Nutrients*, 11(10), p. 2442. doi: 10.3390/nu11102442.
- Sacks, F. M. *et al.* (2017), 'Dietary Fats and Cardiovascular Disease: A Presidential Advisory From the American Heart Association', *Circulation*, 136(3), pp. e1–e23. doi: 10.1161/CIR.0000000000000510.
- SACN (2015), 'Carbohydrates and Health', [https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment\\_data/file/474242/sacn-carbohydrates-and-health-report-2015.pdf](https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/474242/sacn-carbohydrates-and-health-report-2015.pdf). Accessed May 2021.
- SACN (2019) *Saturated fats and health: SACN report* (no date) GOV.UK. Available at: <https://www.gov.uk/government/publications/saturated-fats-and-health-sacn-report> (Accessed: 30 April 2021).
- Santos, V. C. *et al.* (2013), 'Effects of DHA-rich fish oil supplementation on lymphocyte



- function before and after a marathon race'. *Int. J. Sport Nutr. Exerc. Metab.*, vol. 23(2), pp. 161–9.
- Santosa, S. and Jensen, M. D. (2013), 'Adipocyte fatty acid storage factors enhance subcutaneous fat storage in postmenopausal women', *Diabetes*, 62(3), pp. 775–782. doi: 10.2337/db12-0912.
- Saunders, B. *et al.* (2017), 'β-alanine supplementation to improve exercise capacity and performance: a systematic review and meta-analysis', *British Journal of Sports Medicine*, 51(8), pp. 658–669. doi: 10.1136/bjsports-2016-096396.
- Saunders, M. J. (2007), 'Coingestion of carbohydrate-protein during endurance exercise: influence on performance and recovery'. *Int. J. Sports Nutr. Exerc. Metab.*, vol. 17, S87–S103.
- Saunders, M. J. *et al.* (2004), 'Effects of a carbohydrate-protein-beverage on cycling endurance and muscle damage'. *Med. Sci. Sports Exerc.*, vol. 36, pp. 1233–8.
- Sawka, M. N. (1992), 'Physiological consequences of hypohydration: exercise performance and thermoregulation'. *Med. Sci. Sports Exerc.*, vol. 24, pp. 657–70.
- Sawka, M. N. *et al.* (2007), 'American College of Sports Medicine Position stand. Exercise and fluid replacement'. *Med. Sci. Sports Exerc.*, vol. 39, pp. 377–90.
- Sawka, M. N., Cheuvront, S. N. and Kenefick, R. W. (2015), 'Hypohydration and Human Performance: Impact of Environment and Physiological Mechanisms', *Sports Medicine (Auckland, N.z.)*, 45(Suppl 1), pp. 51–60. doi: 10.1007/s40279-015-0395-7.
- Sawka, M. N., Coyle, E.F. (1999), 'Influence of body water and blood volume on thermoregulation and exercise performance in the heat', *Exerc. Sport. Sci. Rev.*, 1999;27:167-218. PMID: 10791017.
- Schoenfeld, B. J. and Aragon, A. A. (2018), 'How much protein can the body use in a single meal for muscle-building? Implications for daily protein distribution', *Journal of the International Society of Sports Nutrition*, 15(1), p. 10. doi: 10.1186/s12970-018-0215-1.
- Schoenfeld, B. J. *et al.* (2013), 'The effect of protein timing on muscle strength and hypertrophy: a meta-analysis'. *J. Int. Soc. Sports Nutr.*, vol. 10(1), pp. 53.
- Schoenfeld, B. J. *et al.* (2017) (a), 'Strength and Hypertrophy Adaptations Between Low- vs. High-Load Resistance Training: A Systematic Review and Meta-analysis', *Journal of Strength and Conditioning Research*, 31(12), pp. 3508–3523. doi: 10.1519/JSC.0000000000002200.
- Schoenfeld, B. J. *et al.* (2017) (b), 'Pre- versus post-exercise protein intake has similar effects on muscular adaptations'. *PeerJ.*, vol. 5, e2825.
- Schokman, C. P. *et al.* (1999), 'Pre- and post game macronutrient intake of a group of elite Australian Football Players'. *Int. J. Sport Nutr.*, vol. 9, pp. 60–9.
- Seebohar, B. (2014), 'Metabolic efficiency training: teaching the body to burn more fat'. (2nd edn), [www.enrgperformance.com](http://www.enrgperformance.com).
- Seiler, D. *et al.* (1989), 'Effects of long-distance running on iron metabolism and hematological parameters'. *Int. J. Sports Med.*, vol. 10, pp. 357–62.
- Seip, R. L. and Semenkovich, C. F. (1998), 'Skeletal muscle lipoprotein lipase: molecular regulation and physiological effects in relation to exercise'. *Exerc. Sport Sci. Rev.*, vol. 26, pp. 191–218.
- Sender, R., Fuchs, S. and Milo, R. (2016), 'Revised Estimates for the Number of Human and Bacteria Cells in the Body', *PLoS Biology*, 14(8). doi: 10.1371/journal.pbio.1002533.
- Senefeld, J. W. *et al.* (2020), 'Ergogenic Effect of Nitrate Supplementation: A Systematic Review and Meta-analysis', *Medicine and Science in Sports and Exercise*, 52(10), pp. 2250–2261. doi: 10.1249/MSS.0000000000002363.
- Shahidi, F. and Ambigaipalan, P. (2018), 'Omega-3 Polyunsaturated Fatty Acids and Their Health Benefits', *Annual Review of Food Science and Technology*, 9, pp. 345–381. doi:

- 10.1146/annurev-food-111317-095850.
- Shaw, G. *et al.* (2017), 'Vitamin C-enriched gelatin supplementation before intermittent activity augments collagen synthesis<sup>12</sup>', *The American Journal of Clinical Nutrition*, 105(1), pp. 136–143. doi: 10.3945/ajcn.116.138594.
- Sherman, W. M. *et al.* (1981), 'Effect of exercise-diet manipulation on muscle glycogen and its subsequent utilisation during performance'. *Int. J. Sports Med.*, vol. 2, pp. 114–18.
- Sherman, W. M. *et al.* (1991), 'Carbohydrate feedings 1 hour before exercise improve cycling performance'. *Am. J. Clin. Nutr.*, vol. 54, pp. 866–70.
- Sherman, W. M. *et al.* (1993), 'Dietary carbohydrate, muscle glycogen, and exercise performance during 7 d of training', *The American Journal of Clinical Nutrition*, 57(1), pp. 27–31. doi: 10.1093/ajcn/57.1.27.
- Shi, X. *et al.* (2004), 'Gastrointestinal discomfort during intermittent high-intensity exercise: effect of carbohydrate-electrolyte beverage', *International Journal of Sport Nutrition and Exercise Metabolism*, 14(6), pp. 673–683. doi: 10.1123/ijsnem.14.6.673.
- Shimizu, K. *et al.* (2011), 'Influences of weight loss on monocytes and T-cell subpopulations in male judo athletes', *Journal of Strength and Conditioning Research*, 25(7), pp. 1943–1950. doi: 10.1519/JSC.0b013e3181e4f9c6.
- Shimomura, Y. *et al.* (2006), 'Nutraceutical effects of branched chain amino acids on skeletal muscle'. *J. Nutr.*, vol. 136, pp. 529–32.
- Shing, C. M., Hunter, D. C. and, Stevenson, L. M. (2009), 'Bovine colostrum supplementation and exercise performance: potential mechanisms'. *Sports Med.*, vol. 39(12), pp. 1033–54.
- Shing, C.M. *et al.* (2007), 'Effects of bovine colostrum supplementation on immune variables in highly trained cyclists'. *J. Appl. Physiol.*, vol. 102, pp. 1113–22.
- Shirreffs S. M. *et al.* (2004), 'Fluid and electrolyte needs for preparation and recovery from training and competition'. *J. Sports Sci.*, vol. 22(1), pp. 57–63.
- Shirreffs, S. M. and Sawka, M. N. (2011), 'Fluid and electrolyte needs for training, competition, and recovery'. *J. Sports Sci.*, vol. 29, suppl 1, pp. S39–46.
- Shirreffs, S. M. *et al.* (2007), 'Milk as an effective post-exercise rehydration drink'. *Br J. Nutr.*, vol. 98, pp. 173–180.
- Shirreffs, S. M., *et al.* (1996), 'Post-exercise rehydration in man: effects of volume consumed. and drink sodium content'. *Med. Sci. Sports Ex.*, vol. 28, pp. 1260–71.
- Short, S. H. and Short, W. R. (1983), 'Four-year study of university athletes' dietary intake'. *J. Am. Diet. Assoc.*, vol. 82, p. 632.
- Siervo, M. *et al.* (2013), 'Inorganic nitrate and beetroot juice supplementation reduces blood pressure in adults: a systematic review and meta-analysis', *The Journal of Nutrition*, 143(6), pp. 818–826. doi: 10.3945/jn.112.170233.
- Silva, M. R. G. and Paiva, T. (2015), 'Low energy availability and low body fat of female gymnasts before an international competition', *European Journal of Sport Science*, 15(7), pp. 591–599. doi: 10.1080/17461391.2014.969323.
- Silva-Cavalcante, M. D. *et al.* (2013), 'Caffeine increases anaerobic work and restores cycling performance following a protocol designed to lower endogenous carbohydrate availability'. *PLoS One*, vol. 8(8), e72025.
- Sim, M. *et al.* (2019), 'Iron considerations for the athlete: a narrative review', *European Journal of Applied Physiology*, 119(7), pp. 1463–1478. doi: 10.1007/s00421-019-04157-y.
- Simopoulos, A. P. and Robinson, J. (1998), *The Omega Plan* (New York, HarperCollins).
- Simpson, R. J. *et al.* (2020), 'Can exercise affect immune function to increase susceptibility to infection?', *Exercise Immunology Review*, 26, pp. 8–22.
- Singh, Pramil N., *et al.* (2003), 'Does Low Meat Consumption Increase Life Expectancy in

- Humans?”, *The American Journal of Clinical Nutrition*, vol. 78, no. 3, Sept. 2003, pp. 526S-532S. academic.oup. com, doi:10.1093/ajcn/78.3.526S.
- Sinning, W. E. (1998), ‘Body composition in athletes’, In Roche, A. F. *et al.* (eds.) *Human Body Composition*, (Champaign, IL: Human Kinetics), pp. 257–73.
- Skaug, A., Sveen, O., and Raastad, T. (2014), ‘An antioxidant and multivitamin supplement reduced improvements in VO<sub>2</sub>max’. *J. Sports Med. Phys. Fitness.*, vol. 54(1), pp. 63–9.
- Slater, G. *et al.* (2001), ‘HMB supplementation does not affect changes in strength or body composition during resistance training in trained men’. *Int. J. Sport Nutr.*, vol. 11, pp. 383–96.
- Sloth, B. *et al.* (2004), ‘No difference in body weight decrease between a low GI and high GI diet but reduced LDL cholesterol after 10 wk ad libitum intake of the low GI diet’. *Am. J. Clin Nutr.*, vol. 80, pp. 337–47.
- Smith, G. I. *et al.* (2011) (a), ‘Dietary omega-3 fatty acid supplementation increases the rate of muscle protein synthesis in older adults: a randomized controlled trial’, *The American Journal of Clinical Nutrition*, 93(2), pp. 402–412. doi: 10.3945/ajcn.110.005611.
- Smith, G. I. *et al.* (2011) (b), ‘Omega-3 polyunsaturated fatty acids augment the muscle protein anabolic response to hyperinsulinaemia-hyperaminoacidaemia in healthy young and middle-aged men and women’, *Clinical Science (London, England: 1979)*, 121(6), pp. 267–278. doi: 10.1042/CS20100597.
- Smith, J. W. *et al.* (2010), ‘Fuel selection and cycling endurance performance with ingestion of [13C] glucose: evidence for a carbohydrate dose response’, *Journal of Applied Physiology (Bethesda, Md.: 1985)*, 108(6), pp. 1520–1529. doi: 10.1152/jappphysiol.91394.2008.
- Snijders, T. *et al.* (2015), ‘Protein ingestion before sleep increases muscle mass and strength gains during prolonged resistance-type exercise training in healthy young men’. *J. Nutr.* In press, Apr 29, 2015.
- Snyder, A. C. *et al.* (1989), ‘Influence of dietary iron source on measures of iron status among female runners’. *Med. Sci. Sports Exerc.*; vol. 21, pp. 7–10.
- Sobiecki, Jakub G., *et al.* (2016), ‘High Compliance with Dietary Recommendations in a Cohort of Meat Eaters, Fish Eaters, Vegetarians, and Vegans: Results from the European Prospective Investigation into Cancer and Nutrition-Oxford Study.’, *Nutrition Research (New York, N.Y.)*, vol. 36, no. 5, May 2016, pp. 464–77. PubMed, doi:10.1016/j.nutres.2015.12.016.
- Soenen, S. *et al.* (2012), ‘Relatively high-protein or “low-carb” energy-restricted diets for body weight loss and body weight maintenance?’, *Physiology & Behavior*, 107(3), pp. 374–380. doi: 10.1016/j.physbeh.2012.08.004.
- Somerville, V. S., Braakhuis, A. J. and Hopkins, W. G. (2016), ‘Effect of Flavonoids on Upper Respiratory Tract Infections and Immune Function: A Systematic Review and Meta-Analysis’, *Advances in Nutrition*, 7(3), pp. 488–497. doi: 10.3945/an.115.010538.
- Sorrenti, V. *et al.* (2020), ‘Deciphering the Role of Polyphenols in Sports Performance: From Nutritional Genomics to the Gut Microbiota toward Phytonutritional Epigenomics’, *Nutrients*, 12(5), p. 1265. doi: 10.3390/nu12051265.
- Southward, K., Rutherford-Markwick, K. J. and Ali, A. (2018), ‘The Effect of Acute Caffeine Ingestion on Endurance Performance: A Systematic Review and Meta-Analysis’, *Sports Medicine (Auckland, N.Z.)*, 48(8), pp. 1913–1928. doi: 10.1007/s40279-018-0939-8.
- Souza, D. B. *et al.* (2017), ‘Acute effects of caffeine-containing energy drinks on physical performance: a systematic review and meta-analysis’, *European Journal of Nutrition*, 56(1), pp. 13–27. doi: 10.1007/s00394-016-1331-9.

- Spector, T. (2015), *The Diet Myth: The real science behind what we eat* (WBH).
- Speedy, D. B. *et al.* (1999), 'Hyponatremia in ultra-distance triathletes'. *Med. Sci. Sports Exerc.*, vol. 31, pp. 809–15.
- Spence, L. *et al.* (2007), 'Incidence, etiology, and symptomatology of upper respiratory illness in elite athletes', *Medicine and Science in Sports and Exercise*, 39(4), pp. 577–586. doi: 10.1249/mss.0b013e31802e851a.
- Spencer, E. A. *et al.* (2003), 'Weight gain over 5 years in 21,966 meat-eating, fish-eating, vegetarian, and vegan men and women in EPIC-Oxford'. *Int. J. Obes. Relat. Metab. Disord.* Jun; 27(6), pp. 728–34.
- Spriet, L. (1995), 'Caffeine and performance'. *Int. J. Sport Nutr.*, vol. 5, pp. S84–S99.
- Spriet, L. L. (2014(a)), 'Exercise and Sport Performance with Low Doses of Caffeine', *Sports Medicine*, 44(2), pp. 175–184. doi: 10.1007/s40279-014-0257-8.
- Spriet, L. L. (2014(b)), 'New insights into the interaction of carbohydrate and fat metabolism during exercise', *Sports Medicine (Auckland, N.Z.)*, 44 Suppl 1, pp. S87-96. doi: 10.1007/s40279-014-0154-1.
- Steen, S. N. and McKinney, S. (1986), 'Nutrition assessment of college wrestlers'. *Phys. Sports Med.*, vol. 14, pp. 100–6.
- Steenge, G. R. *et al.* (1998), 'The stimulatory effect of insulin on creatine accumulation in human skeletal muscle'. *Am. J. Physiol.*, vol. 275, pp. E974–9.
- Stegen, S. *et al.* (2014), 'The beta-alanine dose for maintaining moderately elevated muscle carnosine levels'. *Med. Sci. Sports Exerc.*, 1 (Epub ahead of print).
- Stellingwerff, T. and Cox, G. R. (2014), 'Systematic review: Carbohydrate supplementation on exercise performance or capacity of varying durations', *Applied Physiology, Nutrition, and Metabolism = Physiologie Appliquee, Nutrition Et Metabolisme*, 39(9), pp. 998–1011. doi: 10.1139/apnm-2014-0027.
- Stellingwerff, T. and Cox, G. R. (2014), 'Systematic review: Carbohydrate supplementation on exercise performance or capacity of varying durations', *Applied Physiology, Nutrition, and Metabolism = Physiologie Appliquee, Nutrition Et Metabolisme*, 39(9), pp. 998–1011. doi: 10.1139/apnm-2014-0027.
- Stellingwerff, T. *et al.* (2006), 'Decreased PDH activation and glycogenolysis during exercise following fat adaptation with carbohydrate restoration'. *Am. J. Physiol. Endocrinol. Metab.*, vol. 290(2), pp. 380–8.
- Stevenson, E. *et al.* (2005), 'Improved recovery from prolonged exercise following the consumption of low glycaemic index carbohydrate meals'. *Int. J. Sport Nutr. Exerc. Metab.*, vol. 15, pp. 333–49.
- Stoffel, N. U. *et al.* (2020), 'Iron absorption from supplements is greater with alternate day than with consecutive day dosing in iron-deficient anemic women', *Haematologica*, 105(5), pp. 1232–1239. doi: 10.3324/haematol.2019.220830.
- Strauss, R. H., Lanese, R. R. and Malarkey, W. B. (1985), 'Weight loss in amateur wrestlers and its effect on serum testosterone levels', *JAMA*, 254(23), pp. 3337–3338.
- Stroehlein, J. K. *et al.* (2021), 'Vitamin D supplementation for the treatment of COVID-19: a living systematic review', *The Cochrane Database of Systematic Reviews*, 5, p. CD015043. doi: 10.1002/14651858. CD015043.
- Sun, G. *et al.* (2005), 'Comparison of multifrequency bioelectrical impedance analysis with dual-energy X-ray absorptiometry for assessment of percentage body fat in a large, healthy population'. *Am. J. Clin Nutr.*, vol. 81, pp. 74–8.
- Sundgot-Borgen, J. (1994a)(a), 'Eating disorders in female athletes'. *Sports Med.*, vol. 17(3), pp. 176–88.

- Sundgot-Borgen, J. (1994b)(b), 'Risk and trigger factors for the development of eating disorders in female elite athletes'. *Med. Sci. Sports Ex.*, vol. 26, pp. 414–19.
- Sundgot-Borgen, J. and Garthe, I. (2011), 'Elite athletes in aesthetic and Olympic weight-class sports and the challenge of body weight and body compositions', *Journal of Sports Sciences*, 29 Suppl 1, pp. S101-114. doi: 10.1080/02640414.2011.565783.
- Sundgot-Borgen, J. and Larsen, S. (1993), 'Nutrient intake and eating behaviour in elite female athletes suffering from anorexia nervosa, anorexia athletica and bulimia nervosa'. *Int. J. Sport Nutr.*, vol. 3, pp. 431–42.
- Sundgot-Borgen, J. and Torstveit M. K. (2004), 'Prevalence of eating disorders in elite athletes is higher than in the general population'. *Clin. J. Sport Nutr.*, vol. 14, pp. 25–32.
- Sundgot-Borgen, J. and Torstveit, M. K. (2010), 'Aspects of disordered eating continuum in elite high-intensity sports', *Scandinavian Journal of Medicine & Science in Sports*, 20 Suppl 2, pp. 112–121. doi: 10.1111/j.1600-0838.2010.01190.x.
- Sundgot-Borgen, J. and Torstveit, M. K. (2010), 'Aspects of disordered eating continuum in elite high-intensity sports'. *Scand. J. Med. Sci. Sports.*, vol. 20 (Suppl 2), pp. 112–21.
- Sundgot-Borgen, J. *et al.* (2013), 'How to minimise the health risks to athletes who compete in weight-sensitive sports review and position statement on behalf of the Ad Hoc Research Working Group on Body Composition, Health and Performance, under the auspices of the IOC Medical Commission', *British Journal of Sports Medicine*, 47(16), pp. 1012–1022. doi: 10.1136/bjsports-2013-092966.
- Sutehall, S. *et al.* (2020), 'Addition of an Alginate Hydrogel to a Carbohydrate Beverage Enhances Gastric Emptying', *Medicine and Science in Sports and Exercise*, 52(8), pp. 1785–1792. doi: 10.1249/MSS.0000000000002301.
- Sutton, E. F. *et al.* (2018), 'Early Time-Restricted Feeding Improves Insulin Sensitivity, Blood Pressure, and Oxidative Stress Even without Weight Loss in Men with Prediabetes', *Cell Metabolism*, 27(6), pp. 1212-1221.e3. doi: 10.1016/j.cmet.2018.04.010.
- Swaminathan, R. *et al.* (1985), 'Thermic effect of feeding carbohydrate, fat, protein and mixed meal in lean and obese subjects'. *Am. J. Clin. Nutr.*, vol. 42, pp. 177–81.
- Talanian, J. L. and Spriet, L. L. (2016), 'Low and moderate doses of caffeine late in exercise improve performance in trained cyclists', *Applied Physiology, Nutrition, and Metabolism = Physiologie Appliquee, Nutrition Et Metabolisme*, 41(8), pp. 850–855. doi: 10.1139/apnm-2016-0053.
- Tanaka, H. and Seals, D. R. (2008), 'Endurance exercise performance in Masters athletes: age-associated changes and underlying physiological mechanisms.' *J Physiol.*, Volvol. 586(1), p.55–63.
- Tang, J. E. *et al.* (2009), 'Ingestion of whey hydrolysate, casein, or soya protein isolate: effects on mixed muscle protein synthesis at rest and following resistance exercise in young men'. *J. Appl. Physiol.*, vol. 107(3), pp. 987–92.
- Tarnopolsky, M. and MacLennan, D. P. (1988), 'Influence of protein intake and training status in nitrogen balance and lean body mass'. *J. Appl. Physiol.*, vol. 64, pp. 187–93.
- Tarnopolsky, M. and MacLennan, D. P. (1992), 'Evaluation of protein requirements for trained strength athletes'. *J. Appl. Physiol.*, vol. 73, pp. 1986–95.
- Tarnopolsky, M. and MacLennan, D. P. (1997), 'Post exercise protein-carbohydrate and carbohydrate supplements increase muscle glycogen in males and females'. *J. Appl. Physiol.*, Abstracts, vol. 4, p. 332A.
- Tarnopolsky, M. and MacLennan, D. P. (2000), 'Creatine monohydrate supplementation enhances high-intensity exercise performance in males and females'. *Int. J. Sport Nutr.*, vol. 10, pp. 452–63.

- Terada, T. *et al.* (2019), 'Overnight fasting compromises exercise intensity and volume during sprint interval training but improves high-intensity aerobic endurance', *The Journal of Sports Medicine and Physical Fitness*, 59(3), pp. 357–365. doi: 10.23736/S0022-4707.18.08281-6.
- Theodorou, A. A. *et al.* (2011), 'No effect of antioxidant supplementation on muscle performance and blood redox status adaptations to eccentric training'. *Am. J. Clin. Nutr.*, vol. 93(6), pp. 1373–83.
- Thomas, D. E. *et al.* (1991), 'Carbohydrate feeding before exercise: effect of glycaemic index'. *Int. J. Sports Med.*, vol. 12, pp. 180–6.
- Thomas, D. E. *et al.* (1994), 'Plasma glucose levels after prolonged strenuous exercise correlate inversely with glycaemic response to food consumed before exercise'. *Int. J. Sports Nutr.*, vol. 4, pp. 261–73.
- Thomas, D. T., Erdman, K. A. and Burke, L. M. (2016), 'American College of Sports Medicine Joint Position Statement. Nutrition and Athletic Performance', *Medicine and Science in Sports and Exercise*, 48(3), pp. 543–568. doi: 10.1249/MSS.0000000000000852.
- Thomas, M. H. and Burns, S. P. (2016), 'Increasing Lean Mass and Strength: A Comparison of High Frequency Strength Training to Lower Frequency Strength Training', *International Journal of Exercise Science*, 9(2), pp. 159–167.
- Thompson, C. *et al.* (2016), 'Dietary nitrate supplementation improves sprint and high-intensity intermittent running performance'. *Nitric Oxide*, vol. 61, pp. 55–61.
- Tipton K. D. *et al.* (2001), 'Timing of amino acid-carbohydrate ingestion alters anabolic response of muscle to resistance exercise'. *Am. J. Physiol.*, vol. 281(2), pp. E197–206.
- Tipton, C. M. (1987), 'Commentary: physicians should advise wrestlers about weight loss'. *Phys. Sports Med.*, vol. 15, pp. 160–5.
- Tipton, K. D. (2015), 'Nutritional Support for Exercise-Induced Injuries', *Sports Medicine (Auckland, N.Z.)*, 45 Suppl 1, pp. S93-104. doi: 10.1007/s40279-015-0398-4.
- Tipton, K. D. and Witard, O. C. (2007), 'Protein requirements and recommendations for athletes: relevance of ivory tower arguments for practical recommendations'. *Clin. Sports Med.*, vol. 26(1), pp. 17–36.
- Tipton, K. D. and Wolfe, R. (2007), 'Protein needs and amino acids for athletes'. *J. Sports Sci.*, vol 22(1), pp. 65–79.
- Tipton, K. D. *et al.* (2004), 'Ingestion of casein and whey proteins result in muscle anabolism after resistance exercise'. *Med. Sci. Sports Exerc.*, vol. 36 (12), pp. 2073–81.
- Tipton, K. D. *et al.* (2007), 'Stimulation of net protein synthesis by whey protein ingestion before and after exercise'. *Am. J. Physiol. Endocrinol. Metab.*, vol. 292(1), pp. E71–6.
- Tobias, Deirdre K. *et al.* (2015), 'Effect of low-fat diet interventions versus other diet interventions on long-term weight change in adults: a systematic review and meta-analysis'. *The Lancet Diabetes & Endocrinology*, vol. 3(12), pp. 968–79.
- Tornberg, Å. B. *et al.* (2017), 'Reduced Neuromuscular Performance in Amenorrhoeic Elite Endurance Athletes', *Medicine & Science in Sports & Exercise*, 49(12), pp. 2478–2485. doi: 10.1249/MSS.0000000000001383.
- Torstveit, M. K. and J. Sundgot-Borgen (2005), 'Participation in leanness sports but not training volume is associated with menstrual dysfunction: a national survey of 1276 athletes and controls'. *Br. J. Sports Med.*, vol. 39, pp. 141–7.
- Toth, M. J. *et al.* (2000), 'Menopause-related changes in body fat distribution', *Annals of the New York Academy of Sciences*, 904, pp. 502–506. doi: 10.1111/j.1749-6632.2000.tb06506.x.
- Toth, P. P. (2005), 'The "good cholesterol": High-density lipoprotein', *Circulation*, vol. 111(5), pp. 89–91.

- Trapp, D., Knez, W. and Sinclair, W. (2010), 'Could a vegetarian diet reduce exercise-induced oxidative stress? A review of the literature', *Journal of Sports Sciences*, 28(12), pp. 1261–1268. doi: 10.1080/02640414.2010.507676.
- Trombold, J. R. *et al.* (2011), 'The Effect of Pomegranate Juice Supplementation on Strength and Soreness after Eccentric Exercise', *Journal of strength and conditioning research*. doi: 10.1519/JSC.0b013e318220d992.
- Trommelen, J. *et al.* (2017), 'Fructose and Sucrose Intake Increase Exogenous Carbohydrate Oxidation during Exercise', *Nutrients*, 9(2). doi: 10.3390/nu9020167.
- Truby, H. *et al.* (2008), 'Commercial weight loss diets meet nutrient requirements in free living adults over 8 weeks: a randomised controlled weight loss trial'. *Nutrition Journal* 2008, vol. 7, p. 25.
- Tsintzas, O. K. *et al.* (1995), 'Influence of carbohydrate electrolyte drinks on marathon running performance'. *Eur. J. Appl. Physiol.*, vol. 70, pp. 154–60.
- U.S. Department of Health and Human Services (2018). Physical Activity Guidelines for Americans, 2nd edition. Washington, DC.
- Uauy, R. *et al.* (2009), 'WHO Scientific Update on trans fatty acids: summary and conclusions'. *Eur. J. Clin. Nutr.*, vol. 63, pp. S68–S75.
- Unnithan, V. B. *et al.* (2001), 'Is there a physiologic basis for creatine use in children and adolescents?' *J. Strength Cond. Res.*, vol. 15(4), pp. 524–8.
- Vahedi, K. (2000), 'Ischaemic stroke in a sportsman who consumed mahuang extract and creatine monohydrate for bodybuilding'. *J. Neur., Neurosurgery and Psych.*, vol. 68, pp. 112–13.
- Valdes, A. M. *et al.* (2018), 'Role of the gut microbiota in nutrition and health', *BMJ*, 361, p. k2179. doi: 10.1136/bmj.k2179.
- Van Elswyk, M. E., Weatherford, C. A. and McNeill, S. H. (2018), 'A Systematic Review of Renal Health in Healthy Individuals Associated with Protein Intake above the US Recommended Daily Allowance in Randomized Controlled Trials and Observational Studies', *Advances in Nutrition (Bethesda, Md.)*, 9(4), pp. 404–418. doi: 10.1093/advances/nmy026.
- Van Essen M. and Gibala, M. J. (2006), 'Failure of protein to improve time-trial performance when added to a sports drink'. *Med. Sci. Sports Exerc.*, vol. 38(8), pp. 1476–83.
- Van Loon, L. J. C. (2007), 'Application of protein or protein hydrolysates to improve postexercise recovery'. *Int. J. Sports Nutr. Exerc. Metab.*, vol. 17, S104–17.
- Van Loon, L. J. C. (2014), 'Is there a need for protein ingestion during exercise?' *Sports Med.*, vol. 44 (Suppl. 1), pp. 105–11.
- Van Proeyen, K. *et al.* (2011), 'Beneficial metabolic adaptations due to endurance exercise training in the fasted state'. *J. Appl. Physiol.*, vol. 110(1), pp. 236–45.
- Van Someren, K. A. *et al.* (2005), 'Supplementation with HMB and KIC reduces signs and symptoms of exercise-induced muscle damage in man'. *Int. J. Sport Nutr. Exerc. Metab.*, vol. 15, pp. 413–24.
- Van Thienen, R. *et al.* (2009), 'Beta-alanine improves sprint performance in endurance cycling'. *Med. Sci. Sports Exerc.*, vol. 41, pp. 898–903.
- van Vliet, S. *et al.* (2017), 'Consumption of whole eggs promotes greater stimulation of postexercise muscle protein synthesis than consumption of isonitrogenous amounts of egg whites in young men', *The American Journal of Clinical Nutrition*, 106(6), pp. 1401–1412. doi: 10.3945/ajcn.117.159855.
- van Vliet, S., Burd, N. A. and van Loon, L. J. C. (2015), 'The Skeletal Muscle Anabolic Response to Plant- versus Animal-Based Protein Consumption', *The Journal of Nutrition*, 145(9), pp. 1981–1991. doi: 10.3945/jn.114.204305.

- Vandenbogaerde, T. J. and Hopkins, W. G. (2011), 'Effects of acute carbohydrate supplementation on endurance performance: a meta-analysis'. *Sports Med.*, vol. 41(9), pp. 773–92.
- Venables, M. *et al.* (2005), 'Erosive effect of a new sports drink on dental enamel during exercise'. *Med. Sci. Sports Exerc.*, vol. 37(1), pp. 39–44.
- Volek, J. S. (1997), 'Response of testosterone and cortisol concentrations to high-intensity resistance training following creatine supplementation'. *J. Strength Cond. Res.*, vol. 11, pp. 182–7.
- Volek, J. S. *et al.* (1999), 'Performance and muscle fibre adaptations to creatine supplementation and heavy resistance training'. *Med. Sci. Sports Ex.*, vol. 31(8), pp. 1147–56.
- Volek, J. S. *et al.* (2013), 'Whey protein supplementation during resistance training augments lean body mass'. *J. Am. Coll. Nutr.*, vol. 32(2), pp. 122–35.
- Volpe, S. L. (2010), 'Alcohol and Athletic Performance', *ACSM's Health & Fitness Journal*, 14(3), pp. 28–30. doi: 10.1249/FIT.0b013e3181daa567.
- Wackerhage, H. *et al.* (2019), 'Stimuli and sensors that initiate skeletal muscle hypertrophy following resistance exercise', *Journal of Applied Physiology (Bethesda, Md.: 1985)*, 126(1), pp. 30–43. doi: 10.1152/jappphysiol.00685.2018.
- Waitrose Food and Drink Report 2018–2019  
<https://www.waitrose.com/content/dam/waitrose/Inspiration/Waitrose%20&%20Partners%20>
- Wall, B. T. and van Loon, L. J. C. (2013) (a), 'Nutritional strategies to attenuate muscle disuse atrophy', *Nutrition Reviews*, 71(4), pp. 195–208. doi: 10.1111/nure.12019.
- Wall, B. T. *et al.* (2013) (b), 'Disuse impairs the muscle protein synthetic response to protein ingestion in healthy men', *The Journal of Clinical Endocrinology and Metabolism*, 98(12), pp. 4872–4881. doi: 10.1210/jc.2013-2098.
- Wall, B. T. *et al.* (2014), 'Substantial skeletal muscle loss occurs during only 5 days of disuse', *Acta Physiologica (Oxford, England)*, 210(3), pp. 600–611. doi: 10.1111/apha.12190.
- Wallis, G. A. *et al.* (2005), 'Oxidation of combined ingestion of maltodextrins and fructose during exercise', *Medicine and Science in Sports and Exercise*, 37(3), pp. 426–432. doi: 10.1249/01.mss.0000155399.23358.82.
- Walser, B., Giordano, R. M. and Stebbins, C. L. (2006), 'Supplementation with omega-3 polyunsaturated fatty acids augments brachial artery dilation and blood flow during forearm contraction'. *Eur. J. Appl. Physiol.* Jun; 97(3), pp. 347–54.
- Walsh, N. P. (2018), 'Recommendations to maintain immune health in athletes', *European Journal of Sport Science*, 18(6), pp. 820–831. doi: 10.1080/17461391.2018.1449895.
- Walsh, N. P. (2019), 'Nutrition and athlete immune health: a new perspective.' Available at: <http://www.gssiweb.org:80/sports-science-exchange/article/nutrition-and-athlete-immune-health-a-new-perspective> (Accessed: 6 May 2021).
- Wang, Y. *et al.* (2005), 'Comparison of abdominal adiposity and overall obesity in predicting
- Warburton, D. E., Nicol, C. W. and Bredin, S. S. (2006), 'Health benefits of physical activity: the evidence'. *CMAJ*, vol. 174(6), pp. 801–9.
- Warren, J. *et al.* (2003), 'Low glycaemic index breakfasts and reduced food intake in preadolescent children'. *Paediatrics*, vol. 112, pp. 414–19.
- Wasserfurth, P. *et al.* (2020), 'Reasons for and Consequences of Low Energy Availability in Female and Male Athletes: Social Environment, Adaptations, and Prevention', *Sports Medicine – Open*. doi: 10.1186/s40798-020-00275-6.
- Watt, K. K. O. *et al.* (2004), 'Skeletal muscle total creatine content and creatine transporter gene expression in vegetarians prior to and following creatine supplementation', *Int. J. Sport Nutr. Exerc. Metab.*, vol. 14, pp. 517–31.



- Watt, M. J. *et al.* (2002), 'Intramuscular triacylglycerol, glycogen and acetyl group metabolism during 4 h of moderate exercise in man', *The Journal of Physiology*, 541(Pt 3), pp. 969–978. doi: 10.1113/jphysiol.2002.018820.
- WCRF/AICR (2007), 'Food, Nutrition, Physical Activity, and the Prevention of Cancer: a Global Perspective'.
- Wee, S. L. *et al.* (1999), 'Influence of high and low glycemic index meals on endurance running capacity', *Medicine and Science in Sports and Exercise*, 31(3), pp. 393–399. doi: 10.1097/00005768-199903000-00007.
- Weisgarber, K. D., Candow, D. G. and Vogt, E. S. M. (2012), 'Whey protein before and during resistance exercise has no effect on muscle mass and strength in untrained young adults'. *Int. J. Sport Nutr. Exerc. Metab.*, vol. 22(6), pp. 463–9.
- Wellington, B. M., Leveritt, M. D. and Kelly, V. G. (2017), 'The Effect of Caffeine on Repeat-High-Intensity-Effort Performance in Rugby League Players', *International Journal of Sports Physiology and Performance*, 12(2), pp. 206–210. doi: 10.1123/ijsspp.2015-0689.
- Wells, K. R. *et al.* (2020), 'The Australian Institute of Sport (AIS) and National Eating Disorders Collaboration (NEDC) position statement on disordered eating in high performance sport', *British Journal of Sports Medicine*, 54(21), pp. 1247–1258. doi: 10.1136/bjsports-2019-101813.
- Wemple, R. D. *et al.* (1997), 'Caffeine vs. caffeine-free sports drinks: effects on urine production at rest and during prolonged exercise'. *Int. J. Sports Med.*, vol. 18(1), pp. 40–6.
- West, N. P. *et al.* (2011), 'Supplementation with *Lactobacillus fermentum* VRI (PCC) reduces lower respiratory illness in athletes and moderate exercise-induced immune perturbations'. *Nutrition Journal*, vol. 10, p. 30.
- Westcott, W. L. (2012), 'Resistance training is medicine: effects of strength training on health', *Current Sports Medicine Reports*, 11(4), pp. 209–216. doi: 10.1249/JSR.0b013e31825dabb8.
- Westerterp-Plantenga, M. S. *et al.* (2012), 'Dietary protein – its role in satiety, energetics, weight loss and health'. *Br. J. Nutr.*, vol. 108, Suppl 2, pp. S105–12.
- Westman, E. C. *et al.* (2007), 'Low-carbohydrate nutrition and metabolism', *The American Journal of Clinical Nutrition*, 86(2), pp. 276–284. doi: 10.1093/ajcn/86.2.276.
- Wilborn, C. D. *et al.* (2004), 'Effects of zinc magnesium aspartate (ZMA) supplementation on training adaptations and markers of anabolism and catabolism'. *J. Int. Soc. Sports Nutr.*, vol. 1(2), pp. 12–20.
- Wilk, B. and Bar-Or, O. (1996), 'Effect of drink flavour and NaCl on voluntary drinking and rehydration in boys exercising in the heat'. *J. Appl. Physiol.*, vol. 80, pp. 1112–17.
- Wilkinson, S. B. *et al.* (2007), 'Consumption of fluid skim milk promotes greater protein accretion after resistance exercise than does consumption of an isonitrogenous and isoenergetic soysoya-protein beverage'. *Am. J. Clin. Nutr.*, vol. 85(4), pp. 1031–40.
- Wilkinson, S. B. *et al.* (2008), 'Differential effects of resistance and endurance exercise in the fed state on signalling molecule phosphorylation and protein synthesis in human muscle', *The Journal of Physiology*, 586(Pt 15), pp. 3701–3717. doi: 10.1113/jphysiol.2008.153916.
- Williams, C. and Devlin, J. T. (eds) (1992), *Foods, Nutrition and Performance: An International Scientific Consensus* (London: Chapman and Hall).
- Williams, M. (2005), 'Dietary supplements and sports performance: amino acids', *Journal of the International Society of Sports Nutrition*, 2, pp. 63–67. doi: 10.1186/1550-2783-2-2-63.
- Williams, M. H. (1985), *Nutritional Aspects of Human Physical and Athletic Performance*, (Springfield, IL: Charles C Thomas Publisher).
- Williams, M. H. (1992), *Nutrition for Fitness and Sport* (Dubuque, IO: WilliAm. C. Brown).
- Williams, M. H. (1998), *The Ergogenics Edge* (Champaign, IL: Human Kinetics).

- Williams, M. H. (1999), *Nutrition for Health, Fitness and Sport*, 5th ed. (New York: McGraw-Hill).
- Williams, M. H. *et al.* (1999), *Creatine: The Power Supplement* (Champaign, IL: Human Kinetics).
- Williams, T. D. *et al.* (2020), 'Effect of Acute Beetroot Juice Supplementation on Bench Press Power, Velocity, and Repetition Volume', *The Journal of Strength & Conditioning Research*, 34(4), pp. 924–928. doi: 10.1519/JSC.0000000000003509.
- Williamson, D. A. *et al.* (1995), 'Structured equation modeling of risk factors for the development of eating disorder symptoms in female athletes'. *Int. J. Eating Disorders*, vol. 17(4), 387–93.
- Willis, L. H. *et al.* (2012), 'Effects of aerobic and/or resistance training on body mass and fat mass in overweight or obese adults'. *J. Appl. Physiol.* Dec, 113(12), pp. 1831–7. <http://www.ncbi.nlm.nih.gov/pubmed/23019316>
- Willoughby, D. S. *et al.* (2011), 'Effects of 7 Days of Arginine-Alpha-Ketoglutarate Supplementation on Blood Flow, Plasma L-Arginine, Nitric Oxide Metabolites, and Asymmetric Dimethyl Arginine After Resistance Exercise', *International Journal of Sport Nutrition and Exercise Metabolism*, 21(4), pp. 291–299. doi: 10.1123/ijsnem.21.4.291.
- Willoughby, D. S., Stout, J. R. and Wilborn, C. D. (2007), 'Effects of resistance training and protein plus amino acid supplementation on muscle anabolism, mass, and strength', *Amino Acids*, 32(4), pp. 467–477. doi: 10.1007/s00726-006-0398-7.
- Wilmore, J. H. (1983), 'Body composition in sport and exercise'. *Med. Sci. Sports Ex.*, vol. 15, pp. 21–31.
- Wilson, J. M. *et al.* (2013), 'International Society of Sports Nutrition Position Stand: beta-hydroxy-beta-methylbutyrate (HMB)', *Journal of the International Society of Sports Nutrition*, 10(1), p. 6. doi: 10.1186/1550-2783-10-6.
- Wilson, P. B. (2017), 'Frequency of Chronic Gastrointestinal Distress in Runners: Validity and Reliability of a Retrospective Questionnaire', *International Journal of Sport Nutrition and Exercise Metabolism*, 27(4), pp. 370–376. doi: 10.1123/ijsnem.2016-0305.
- Witard O. C., *et al.* (2016), 'Protein Considerations for Optimising Skeletal Muscle Mass in Healthy Young and Older Adults.' *Nutrients*. Vol 8 (4), p. 181.
- Witard, O. C. *et al.* (2011), 'Effect of increased dietary protein on tolerance to intensified training'. *Med. Sci. Sports Exerc.*, vol. 43(4), pp. 598–607.
- Witard, O. C. *et al.* (2014) (a), 'High dietary protein restores overreaching induced impairments in leukocyte trafficking and reduces the incidence of upper respiratory tract infection in elite cyclists'. *Brain Behav. Immun.*, vol. 39, pp. 211–9.
- Witard, O. C. *et al.* (2014) (b), 'Myofibrillar muscle protein synthesis rates subsequent to a meal in response to increasing doses of whey protein at rest and after resistance exercise'. *Am. J. Clin. Nutr.*, vol. 99(1), pp. 86–95.
- Witard, O. C., Garthe, I. and Phillips, S. M. (2019), 'Dietary Protein for Training Adaptation and Body Composition Manipulation in Track and Field Athletes', *International Journal of Sport Nutrition and Exercise Metabolism*, 29(2), pp. 165–174. doi: 10.1123/ijsnem.2018-0267.
- Wojtaszewski, J. F. P. *et al.* (2003), 'Regulation of 5'AMP-activated protein kinase activity and substrate utilization in exercising human skeletal muscle', *American Journal of Physiology. Endocrinology and Metabolism*, 284(4), pp. E813–822. doi: 10.1152/ajpendo.00436.2002.
- Wolfe, R. R. (2017), 'Branched-chain amino acids and muscle protein synthesis in humans: myth or reality?', *Journal of the International Society of Sports Nutrition*, 14, p. 30. doi: 10.1186/s12970-017-0184-9.

- Woods, A. L., Garvican-Lewis, L. A., Lundy, B., *et al.* (2017) (a), 'New approaches to determine fatigue in elite athletes during intensified training: Resting metabolic rate and pacing profile', *PLoS One*, 12(3), p. e0173807. doi: 10.1371/journal.pone.0173807.
- Woods, A. L., Garvican-Lewis, L. A., Rice, A., *et al.* (2017) (b), '12 days of altitude exposure at 1800 m does not increase resting metabolic rate in elite rowers', *Applied Physiology, Nutrition, and Metabolism = Physiologie Appliquee, Nutrition Et Metabolisme*, 42(6), pp. 672–676. doi: 10.1139/apnm-2016-0693.
- World Cancer Research Fund (WCRF) (2018), 'Diet, Nutrition, Physical Activity and Cancer: A Global Perspective The Third Expert Report'.
- World Health Organization (2003), 'Expert consultation on diet, nutrition and the prevention of chronic diseases'. WHO Technical Report Series No. 916, Geneva, Switzerland.
- World Health Organization WHO (2016 no date), WHO *Effects of saturated fatty acids on serum lipids and lipoproteins: a systematic review and regression analysis* (no date) WHO. World Health Organization. Available at: [http://www.who.int/nutrition/publications/nutrientrequirements/sfa\\_systematic\\_review/en/](http://www.who.int/nutrition/publications/nutrientrequirements/sfa_systematic_review/en/) (Accessed: 30 April 2021).
- Wright, D. W. *et al.* (1991), 'Carbohydrate feedings before, during or in combination improve cycling endurance performance'. *J. Appl. Physiol.*, vol. 71, pp. 1082–88.
- Wu, C. L. and Williams, C. (2006), 'A low glycaemic index meal before exercise improves endurance running capacity in men'. *Int. J. Sports Nutr. Exerc. Metab.*, vol 16, pp. 510–27.
- Wu, C. L. *et al.* (2003), 'The influence of high carbohydrate meals with different glycaemic indices on substrate utilisation during subsequent exercise'. *Br. J. Nutr.*, vol. 90(6), pp. 1049–56.
- Wu, G. D. *et al.* (2011), 'Linking long-term dietary patterns with gut microbial enterotypes', *Science (New York, N.Y.)*, 334(6052), pp. 105–108. doi: 10.1126/science.1208344.
- Wylie, L. J. *et al.* (2013), 'Beetroot juice and exercise: pharmacodynamic and dose-response relationships'. *J. Appl. Physiol.*, vol 115(3), pp. 325–36.
- Yang, Y. *et al.* (2012) (a), 'Myofibrillar protein synthesis following ingestion of soy protein isolate at rest and after resistance exercise in elderly men', *Nutrition & Metabolism*, 9(1), p. 57. doi: 10.1186/1743-7075-9-57.
- Yang, Y. *et al.* (2012) (b), 'Resistance exercise enhances myofibrillar protein synthesis with graded intakes of whey protein in older men', *The British Journal of Nutrition*, 108(10), pp. 1780–1788. doi: 10.1017/S0007114511007422.
- Yaspelkis, B. B., *et al.* (1993), 'Carbohydrate supplementation spares muscle glycogen during variable-intensity exercise'. *J. Appl. Physiol.* Oct, 75(4), pp. 1477–85.
- Yeo, W. K. *et al.* (2008), 'Skeletal muscle adaptation and performance responses to once a day versus twice-every-second-day endurance training regimens'. *J. Appl. Physiol.*, vol. 105, pp. 1462–70.
- Yeo, W. K. *et al.* (2010), 'Acute signalling responses to intense endurance training commenced with low or normal muscle glycogen', *Experimental Physiology*, 95(2), pp. 351–358. doi: 10.1113/expphysiol.2009.049353.
- Yfanti, C. *et al.* (2010), 'Antioxidant supplementation does not alter endurance training adaptation'. *Med. Sci. Sports Exerc.*, vol. 42(7), pp. 1388–95.
- Young, V. R., and Pellett, P. L. (1994), Pellett. "Plant Proteins in Relation to Human Protein and Amino Acid Nutrition.", *The American Journal of Clinical Nutrition*, vol. 59, no. 5, May 1994, pp. 1203S-1212S. [academic.oup.com](http://academic.oup.com), doi:10.1093/ajcn/59.5.1203S.
- Yüksel, H. *et al.* (2006) 'Effects of oral continuous 17 $\beta$ -estradiol plus norethisterone acetate replacement therapy on abdominal subcutaneous fat, serum leptin levels and body

- composition', *Gynecological endocrinology: the official journal of the International Society of Gynecological Endocrinology*, 22, pp. 381–7. doi: 10.1080/09513590600842281.
- Zajac, A. *et al.* (2014), 'The effects of a ketogenic diet on exercise metabolism and physical performance in off-road cyclists'. *Nutrients*, vol. 6(7), pp. 2493–508.
- Zawadzki, K. M. *et al.* (1992), 'Carbohydrate-protein complex increases the rate of muscle glycogen storage after exercise'. *J. Appl. Physiol.*, vol. 72, pp. 1854–9.
- Ziberna, L. *et al.* (2013), 'The endothelial plasma membrane transporter bilitranslocase mediates rat aortic vasodilation induced by anthocyanins'. *Nutr. Metab. Cardiovasc. Dis.*, vol. 23(1), pp. 68–74.
- Ziegenfuss, T. *et al.* (1997), 'Acute creatine ingestion: effects on muscle volume, anaerobic power, fluid volumes and protein turnover'. *Med. Sci. Sports Ex.*, vol. 29, supp. 127.
- Ziegler, P. J. *et al.* (1999), 'Nutritional and physiological status of US National Figure Skaters'. *Int. J. Sport Nutr.*, vol. 9, pp. 345–60.
- Zouhal, H. *et al.* (2011) 'Inverse relationship between percentage body weight change and finishing time in 643 forty-two-kilometre marathon runners', *British Journal of Sports Medicine*, 45(14), pp. 1101–1105. doi: 10.1136/bjism.2010.074641.
- Zouhal, H. *et al.* (2020), '<p>Exercise Training and Fasting: Current Insights</p>', *Open Access Journal of Sports Medicine*, 11, pp. 1–28. doi: 10.2147/OAJSM.S224919.
- Zuhl, M. *et al.* (2014), 'Exercise regulation of intestinal tight junction proteins', *British Journal of Sports Medicine*, 48(12), pp. 980–986. doi: 10.1136/bjsports-2012-091585.