

References

Chapter 1

1. Messier SP, Gutekunst DJ, Davis C, Devita P. Weight loss reduces knee-joint loads in overweight and obese older adults with knee osteoarthritis. *Arthritis Rheum.* 2005;52(7):2026-2032.
2. Migliaccio S, Greco EA, Fornari R, Donini LM, Di Luigi L, Lenzi A. Skeletal alterations in women affected by obesity. *Aging Clin Exp Res.* 2013;25 Suppl 1:S35-S37.
3. Ackerman IN, Osborne RH. Obesity and increased burden of hip and knee joint disease in Australia: results from a national survey. *BMC Musculoskelet Disord.* 2012;13:254.
4. Sundquist K, Winkleby M, Li X, Ji J, Hemminki K, Sundquist J. Familial transmission of coronary heart disease: a cohort study of 80,214 Swedish adoptees linked to their biological and adoptive parents. *Am Heart J.* 2011;162(2):317-323.
5. Craft LL, Perna FM. The benefits of exercise for the clinically depressed. *Prim Care Companion J Clin Psychiatry.* 2004;6(3):104-111.
6. Miller MC. *Understanding Depression.* Harvard Medical School; 2011.
7. Schoenfeld TJ, Rada P, Pieruzzini PR, Hsueh B, Gould E. Physical exercise prevents stress-induced activation of granule neurons and enhances local inhibitory mechanisms in the dentate gyrus. *J Neurosci.* 2013;33(18):7770-7777.
8. Driver HS, Taylor SR. Exercise and sleep. *Sleep Med Rev.* 2000;4(4):387-402.
9. Peterlin BL, Rosso AL, Rapoport AM, Scher AI. Obesity and migraine: the effect of age, gender and adipose tissue distribution. *Headache.* 2010;50(1):52-62.
10. Gelaye B, Sacco S, Brown WJ, et al. Body composition status and the risk of migraine: a meta-analysis. *Neurology.* 2017;88(19):1795-1804.
11. Viardot A, Lord RV, Samaras K. The effects of weight loss and gastric banding on the innate and adaptive immune system in type 2 diabetes and prediabetes. *J Clin Endocrinol Metab.* 2010;doi:10.1210/jc.2009-2371.
12. Iglay HB, Thyfault JP, Apolzan JW, Campbell WW. Resistance training and dietary protein: effects on glucose tolerance and contents of skeletal muscle insulin signaling proteins in older persons. *Am J Clin Nutr.* 2007;85(4):1005-1013.
13. Dolezal BA, Potteiger JA. Concurrent resistance and endurance training influence basal metabolic rate in nondieting individuals. *J Appl Physiol (1985).* 1998;85(2):695-700.
14. Josse AR, Tang JE, Tarnopolsky MA, Phillips SM. Body composition and strength changes in women with milk and resistance exercise. *Med Sci Sports Exerc.* 2010;42(6):1122-1130.
15. Abboud G, Greer BK, Campbell SC, Panton LB. Effects of load-volume on EPOC after acute bouts of resistance training in resistance-trained men. *J Strength Cond Res.* 2013;27(7):1936-1941.
16. Vella CA, Kravitz L. Exercise after-burn: a research update. *IDEA Fitness Journal.* 2004;November.
17. Clark JE. Diet, exercise or diet with exercise: comparing the effectiveness of treatment options for weight-loss and changes in fitness for adults (18-65 years old) who are overweight, or obese; systematic review and meta-analysis. *J Diabetes Metab Disord.* 2015;14:31.
18. Willis LH, Slentz CA, Bateman LA, et al. Effects of aerobic and/or resistance training on body mass and fat mass in overweight or obese adults. *J Appl Physiol (1985).* 2012;113(12):1831-1837.
19. Heden T, Lox C, Rose P, Reid S, Kirk EP. One-set resistance training elevates energy expenditure for 72 h similar to three sets. *Eur J Appl Physiol.* 2011;111(3):477-484.
20. Bahr R, Sejersted OM. Effect of intensity of exercise on excess postexercise oxygen consumption. *Metabolism.* 1991;40(8):836-841.
21. Phelain JF, Reinke E, Harris MA, Melby CL. Postexercise energy expenditure and substrate oxidation in young women resulting from exercise bouts of different intensity. *J Am Coll Nutr.* 1997;16(2):140-146.
22. Yingling VR, Yack HJ, White SC. The effect of rearfoot motion on attenuation of the impulse wave at impact during running. *J Appl Biomech.* 1996;12:313-325.

Chapter 2

1. Wang Z, Heshka S, Zhang K, Boozer CN, Heymsfield, SB. Resting energy expenditure: systematic organization and critique of prediction methods. *Obes Res.* 2001;9:331-336.
2. Wang Z, Ying Z, Bosy-Westphal A, et al. Evaluation of specific metabolic rates of major organs and tissues: comparison between men and women. *Am J Hum Biol.* 2011;23(3):333-338.
3. Wolfe RR. The underappreciated role of muscle in health and disease. *Am J Clin Nutr.* 2006;84:475-482.
4. Segal KR, Gutin B, Nyman AM, Pi-Sunyer FX. Thermic effect of food at rest, during exercise, and after exercise in lean and obese men of similar body weight. *J Clin Invest.* 1985;76(3):1107-1112.
5. Smith J, McNaughton L. The effects of intensity of exercise on excess post-exercise oxygen consumption and energy expenditure in moderately trained men and women. *Eur J Appl Physiol.* 1993;67:420-425.
6. Wallace MB, Mills BD, Browning CL. Effects of cross-training on markers of insulin resistance/hyperinsulinemia. *Med Sci Sports Exerc.* 1997;29(9):1170-1175.

Chapter 3

1. Sacks FM, Bray GA, Carey VJ, et al. Comparison of weight-loss diets with different compositions of fat, protein, and carbohydrates. *N Engl J Med.* 2009;360(9):859-873.
2. Zalesin KC, Franklin BA, Lillystone MA, et al. Differential loss of fat and lean mass in the morbidly obese after bariatric surgery. *Metab Syndr Relat Disord.* 2010;8(1):15-20.
3. Santarpia L, Contaldo F, Pasanisi F. Body composition changes after weight-loss interventions for overweight and obesity. *Clin Nutr.* 2013;32(2):157-161.
4. Chaston TB, Dixon JB, O'Brien PE. Changes in fat-free mass during significant weight loss: a systematic review. *Int J Obes (Lond).* 2007;31(5):743-750.
5. Redman LM, Heilbronn LK, Martin CK, et al. Metabolic and behavioral compensations in response to caloric restriction: implications for the maintenance of weight loss. *PLoS One.* 2009;4(2):e4377.
6. Garthe I, Raastad T, Refsnes PE, Koivisto A, Sundgot-Borgen J. Effect of two different weight-loss rates on body composition

- and strength and power-related performance in elite athletes. *Int J Sport Nutr Exerc Metab.* 2011;21(2):97-104.
7. Mero AA, Huovinen H, Matintupa O, et al. Moderate energy restriction with high protein diet results in healthier outcome in women. *J Int Soc Sports Nutr.* 2010;7(1):4.
 8. Martin CK, Das SK, Lindblad L, et al. Effect of calorie restriction on the free-living physical activity levels of nonobese humans: results of three randomized trials. *J Appl Physiol.* 2011;110(4):956-963.
 9. Mozaffarian D, Katan M, Ascherio A, Stampfer M, Willett W. Trans fatty acids and cardiovascular disease. *N Engl J Med.* 2006;354:1601-1613.
 10. Holmstrup, ME, Owns CM, Fairchild TJ, Kanaley JA. Effect of meal frequency on glucose and insulin excursions over the course of a day. *E Spen Eur E J Clin Nutr Metab.* 2010;5(6):e277-e280.
 11. Halton T, Hu F. The effects of high protein diets on thermogenesis, satiety, and weight loss: a critical review. *J Am Coll Nutr.* 2004;23:373-385.
 12. Kirpich AR, Maryniuk, MD. The 3 R's of glycemic index: recommendations, research, and the real world. *Clin Diabetes* 2011;29(4):155-159.
 13. Hall KD. What is the required energy deficit per unit weight loss? *Int J Obes (Lond).* 2008;32(3):573-576.
 14. van der Ploeg GE, Brooks AG, Withers RT, Dollman J, Leane F, Chatterton BE. Body composition changes in female bodybuilders during preparation for competition. *Eur J Clin Nutr.* 2001;55(4):268-277.
 15. Withers RT, Noell CJ, Whittingham NO, Chatterton BE, Schultz CG, Keeves JP. Body composition changes in elite male bodybuilders during preparation for competition. *Aust J Sci Med Sport.* 1997;29(1):11-16.
 16. Alcock J. *Animal behavior: an evolutionary approach.* Sinaur Press; 1975.
 17. Staddon J. A note on the evolutionary significance of "supernormal" stimuli. *Am Nat.* 1975;109(969): 541-545.
 18. Tinbergen N. Social releasers and the experimental method required for their study. *Wilson Bull.* 1948;60:6-51.
 19. Guthrie H. Selection and quantification of typical food portions by young adults. *J Am Diet Assoc.* 1984;84(12):1440-1444.
 20. Schwartz J, Byrd-Bredbenner C. Portion distortion: Typical portion sizes selected by young adults. *J Am Diet Assoc.* 2006;106(9):1412-1418.
 21. Mettler S, Mitchell N, Tipton KD. Increased protein intake reduces lean body mass loss during weight loss in athletes. *Med Sci Sports Exerc.* 2010;42(2):326-337.
 22. Frestedt JL, Zenk JL, Kuskowski MA, Ward LS, Bastian ED. A whey-protein supplement increases fat loss and spares lean muscle in obese subjects: a randomized human clinical study. *Nutr Metab* 2008;5:8.
 23. Westerterp-Plantenga MS. The significance of protein in food intake and body weight regulation. *Curr Opin Clin Nutr Metab Care.* 2003;6(6):635-638.
 24. Brinkworth G, Noakes M, Parker B, Foster P, Clifton PM. Long-term effects of advice to consume a high-protein, low-fat diet, rather than a conventional weight-loss diet, in obese adults with type 2 diabetes: one-year follow-up of a randomized trial. *Diabetologia.* 2004;47(10):1677-1686.
 25. Westerterp-Plantenga MS, Lejeune MP, Nijs I, van Ooijen M, Kovacs EM. High protein intake sustains weight maintenance after body weight loss in humans. *Int J Obes Relat Metab Disord.* 2004;28(1):57-64.
 26. Lejeune MP, Kovacs EM, Westerterp-Plantenga MS. Additional protein intake limits weight regain after weight loss in humans. *Br J Nutr.* 2005;93(2):281-289.
 27. Westerterp-Plantenga MS, Lejeune MP, Smeets AJ, Luscombe-Marsh ND. Sex differences in energy homeostasis following a diet relatively high in protein exchanged with carbohydrate, assessed in a respiration chamber in humans. *Physiol Behav.* 2009;97(3-4):414-419.
 28. Poortmans JR, Dellalieux O. Do regular high-protein diets have potential health risks on kidney function in athletes? *Int J Sport Nutr Exerc Metab.* 2000;10(1):28-38.
 29. Martin WF, Armstrong LE, Rodriguez NR. Dietary protein intake and renal function. *Nutr Metab (Lond).* 2005;2:25.
 30. Heaney RP. Effects of caffeine on bone and the calcium economy. *Food Chem Toxicol.* 2002;40(9):1263-1270.
 31. Kerstetter JE, O'Brien KO, Insogna KL. Low protein intake: the impact on calcium and bone homeostasis in humans. *J Nutr.* 2003;133(3):855S-861S.
 32. Porter KH, Johnson MA. Dietary protein supplementation and recovery from femoral fracture. *Nutr Rev.* 1998;56(11):337-340.
 33. Goldstein ER, Ziegenfuss T, Kalman D, et al. International society of sports nutrition position stand: caffeine and performance. *J Int Soc Sports Nutr.* 2010;7:5.
- ### Chapter 9
1. Cheung K, Hume P, Maxwell L. Delayed onset muscle soreness: treatment strategies and performance factors. *Sports Med.* 2003;33(2):145-164.
- ### Chapter 10
1. Medeiros HS Jr, Mello RS, Amorim MZ, Koch AJ, Machado M. Planned intensity reduction to maintain repetitions within recommended hypertrophy range. *Int J Sports Physiol Perform.* 2013;8(4):384-390.
- ### Chapter 11
1. Medeiros HS Jr, Mello RS, Amorim MZ, Koch AJ, Machado M. Planned intensity reduction to maintain repetitions within recommended hypertrophy range. *Int J Sports Physiol Perform.* 2013;8(4):384-390.
- ### Chapter 13
1. von Loeffelholz C, Birkenfeld AL. The role of non-exercise activity thermogenesis in human obesity. [In: Endotext [Internet].] South Dartmouth (MA): MDText.com, Inc.; 2000. 2018 Apr 9.
 2. Nedeltcheva AV, Kilgus JM, Imperial J, Schoeller DA, Penev PD. Insufficient sleep undermines dietary efforts to reduce adiposity. *Ann Intern Med.* 2010 Oct 5;153(7):435-441.
 3. Tasali E, Wroblewski K, Kahn E, Kilgus J, Schoeller DA. Effect of sleep extension on objectively assessed energy intake among adults with overweight in real-life settings: a randomized clinical trial. *JAMA Intern Med.* 2022;182(4):365-374.
 4. Sale DG. Neural adaptation to resistance training. *Med Sci Sports Exerc.* 1988;20(5 Suppl):S135-S145.
 5. Selye H. Stress and the general adaptation syndrome. *Br Med J.* 1950;1:1383.
 6. DeFreitas JM, Beck TW, Stock MS, Dillon MA, Kasishke PR 2nd. An examination of the time course of training-induced skeletal muscle hypertrophy. *Eur J Appl Physiol.* 2011;111(11):2785-2790.
 7. Seynnes OR, de Boer M, Narici MV. Early skeletal muscle hypertrophy and architectural changes in response to high-intensity resistance training. *J Appl Physiol* 2007;102(1):368-373.
 8. Folland JP, Williams AG. The adaptations to strength training: morphological and neurological contributions to increased strength. *Sports Med.* 2007;37(2):145-168.