

- World Health Organization . The SuRF Report 2: Surveillance of Chronic Disease Risk Factors: Country-Level Data and Comparable Estimates. Geneva: WHO; 2005.
- Hales CM, Fryar CD, Carroll MD, Freedman DS, Aoki Y, Ogden CL. Differences in obesity prevalence by demographic characteristics and urbanization level among adults in the United States, 2013–2016. *JAMA*. 2018;319(23):2419. doi: 10.1001/jama.2018.7270. - DOI - PMC - PubMed
- Flegal KM, Kit BK, Orpana H, Graubard BI. Association of all-cause mortality with overweight and obesity using standard body mass index categories: a systematic review and meta-analysis. *JAMA*. 2013;309(1):71–82. doi: 10.1001/jama.2012.113905. - DOI - PMC - PubMed
- Kitahara CM, et al. Association between class III obesity (BMI of 40–59 kg/m<sup>2</sup>) and mortality: a pooled analysis of 20 prospective studies. *PLoS Med*. 2014;11(7):e1001673. doi: 10.1371/journal.pmed.1001673. - DOI - PMC - PubMed
- Branca F, Nikogosian H, Lobstein T, editors. Die Herausforderung Adipositas und Strategien zu ihrer Bekämpfung in der Europäischen Region der WHO: Zusammenfassung. Kopenhagen: WHO Regionalbüro für Europa; 2007.
- Vogt LJ, et al. magnetic resonance imaging of changes in abdominal compartments in obese diabetics during a low-calorie weight-loss program. *PLoS ONE*. 2016;11(4):e0153595. doi: 10.1371/journal.pone.0153595. - DOI - PMC - PubMed
- Klopfenstein BJ, Kim MS, Krisky CM, Szumowski J, Rooney WD, Purnell JQ. Comparison of 3 T MRI and CT for the measurement of visceral and subcutaneous adipose tissue in humans. *Br. J. Radiol*. 2012;85(1018):e826–e830. doi: 10.1259/bjr/57987644. - DOI - PMC - PubMed
- Thörmer G, et al. Software for automated MRI-based quantification of abdominal fat and preliminary evaluation in morbidly obese patients. *J. Magn. Reson. Imaging*. 2013;37(5):1144–1150. doi: 10.1002/jmri.23890. - DOI - PubMed
- Schaudinn A, et al. Predictive accuracy of single- and multi-slice MRI for the estimation of total visceral adipose tissue in overweight to severely obese patients: MRI prediction of visceral fat volumes. *NMR Biomed*. 2015;28(5):583–590. doi: 10.1002/nbm.3286. - DOI - PubMed
- Linder N, et al. Age and gender specific estimation of visceral adipose tissue amounts from radiological images in morbidly obese patients. *Sci. Rep*. 2016 doi: 10.1038/srep22261. - DOI - PMC - PubMed
- Stange R, Linder N, Schaudinn A, Kahn T, Busse H. Dicomflex: a novel framework for efficient deployment of image analysis tools in radiological research. *PLoS ONE*. 2018;13(9):e0202974. doi: 10.1371/journal.pone.0202974. - DOI - PMC - PubMed
- Pausch T, et al. Cachexia but not obesity worsens the postoperative outcome after pancreatoduodenectomy in pancreatic cancer. *Surgery*. 2012;152(3):S81–S88. doi: 10.1016/j.surg.2012.05.028. - DOI - PubMed
- Sadanathan SA, et al. Automated segmentation of visceral and subcutaneous (deep and superficial) adipose tissues in normal and overweight men: automated segmentation of adipose tissue. *J. Magn. Reson. Imaging*. 2015;41(4):924–934. doi: 10.1002/jmri.24655. - DOI - PubMed
- Shen W, Chen J, Gantz M, Velasquez G, Punyanitya M, Heymsfield SB. A single MRI slice does not accurately predict visceral and subcutaneous adipose tissue changes during weight loss. *Obesity*. 2012;20(12):2458–2463. doi: 10.1038/oby.2012.168. - DOI - PMC - PubMed
- Maurovich-Horvat P, Massaro J, Fox CS, Moselewski F, O'Donnell CJ, Hoffmann U. Comparison of anthropometric, area- and volume-based assessment of abdominal subcutaneous and visceral adipose tissue volumes using multi-detector computed tomography. *Int. J. Obes*. 2007;31(3):500–506. doi: 10.1038/sj.ijo.0803454. - DOI - PubMed
- Kanaley JA, Giannopoulou I, Ploutz-Snyder LL. Regional differences in abdominal fat loss. *Int. J. Obes*. 2007;31(1):147–152. doi: 10.1038/sj.ijo.0803359. - DOI - PubMed
- Blüher M. Mechanisms in endocrinology: are metabolically healthy obese individuals really healthy? *Eur. J. Endocrinol*. 2014;171(6):R209–R219. doi: 10.1530/EJE-14-0540. - DOI - PubMed
- Thomas EL, Bell JD. Influence of undersampling on magnetic resonance imaging measurements of intra-abdominal adipose tissue. *Int. J. Obes. Relat. Metab. Disord*. 2003;27(2):211–218. doi: 10.1038/sj.ijo.802229. - DOI - PubMed
- House MG, et al. Preoperative predictors for complications after pancreaticoduodenectomy: impact of BMI and body fat distribution. *J. Gastrointest. Surg*. 2008;12(2):270–278. doi: 10.1007/s11605-007-0421-7. - DOI - PubMed
- Li X-T, Tang L, Chen Y, Li Y-L, Zhang X-P, Sun Y-S. Visceral and subcutaneous fat as new independent predictive factors of

- survival in locally advanced gastric carcinoma patients treated with neo-adjuvant chemotherapy. *J. Cancer Res. Clin. Oncol.* 2015;141(7):1237–1247. doi: 10.1007/s00432-014-1893-y. - DOI - PubMed
- Raman JD, Reynolds C, Hannon M. An increasing proportion of perinephric to subcutaneous fat is associated with adverse perioperative outcomes of robotic partial nephrectomy. *J. Robot. Surg.* 2016;10(3):255–259. doi: 10.1007/s11701-016-0593-9. - DOI - PubMed
- Estimation of abdominal subcutaneous fat volume of obese adults from single-slice MRI data - Regression coefficients and agreement. Linder N, Michel S, Eggebrecht T, Schaudinn A, Blüher M, Dietrich A, Denecke T, Busse H. *Eur J Radiol.* 2020 Sep;130:109184. doi: 10.1016/j.ejrad.2020.109184. Epub 2020 Jul 15. PMID: 32712498
- Garvey WT, et al. American Association of Clinical Endocrinologists and American College of Endocrinology comprehensive clinical practice guidelines for medical care of patients with obesity. *Endocr Pract.* 2016;22 Suppl 3:1. - PubMed - DOI
- Ford ES. The epidemiology of obesity and asthma. *J Allergy Clin Immunol.* 2005;115(5):897. - PubMed - DOI
- Peters-Golden M, et al. Influence of body mass index on the response to asthma controller agents. *Eur Respir J.* 2006;27(3):495–503. - PubMed - DOI
- Xie L, et al. Discontinuation and reduction of asthma medications after metabolic and bariatric surgery: a systematic review and meta-analysis. *Obes Rev.* 2023;24(2):e13527. - PubMed - DOI
- Dias-Junior SA, et al. Effects of weight loss on asthma control in obese patients with severe asthma. *Eur Respir J.* 2014;43(5):1368–77. - PubMed - DOI
- Ross R, et al. Does the relationship between waist circumference, morbidity and mortality depend on measurement protocol for waist circumference? *Obes Rev.* 2008;9(4):312–25. - PubMed - DOI
- Beuther DA, Sutherland ER. Overweight, obesity, and incident asthma: a meta-analysis of prospective epidemiologic studies. *Am J Respir Crit Care Med.* 2007;175(7):661–6. - PubMed - PMC - DOI
- Deng X, et al. Association between overweight or obesity and the risk for childhood asthma and wheeze: an updated meta-analysis on 18 articles and 73 252 children. *Pediatr Obes.* 2019;14(9):e12532. - PubMed - DOI
- Jiang D, et al. Association between abdominal obesity and asthma: a meta-analysis. *Allergy Asthma Clin Immunol.* 2019;15:16. - PubMed - PMC - DOI
- Ma J, Xiao L, Knowles SB. Obesity, insulin resistance and the prevalence of atopy and asthma in US adults. *Allergy.* 2010;65(11):1455–63. - PubMed - DOI
- Musaad SMA, et al. Comparison of anthropometric measures of obesity in childhood allergic asthma: central obesity is most relevant. *J Allergy Clin Immunol.* 2009;123(6):1321. - PubMed - PMC - DOI
- Wang Y, et al. Comparison of abdominal adiposity and overall obesity in predicting risk of type 2 diabetes among men. *Am J Clin Nutr.* 2005;81(3):555–63. - PubMed - DOI
- Xu S, Gilliland FD, Conti DV. Elucidation of causal direction between asthma and obesity: a bi-directional Mendelian randomization study. *Int J Epidemiol.* 2019;48(3):899–907. - PubMed - PMC - DOI
- World Health Organization: Cancer. 2018. [Last accessed on 2021 May 10]. Available from: <https://www.who.int/news-room/factsheets/detail/cancer> .
- Ferlay JE, Lam F, Colombet M, Mery L, Piñeros M. Global Cancer Observatory: Cancer Today. Lyon: International Agency for Research on Cancer. 2020. [Last accessed on 2021 May 09]. Available from: <https://gco.iarc.fr/today/home> .
- International Agency for Research on Cancer. Latest Global Cancer Data: Cancer Burden Rises to 18.1 Million New Cases and 9.6 Million Cancer Deaths in 2018. [Last accessed on 2021 May 10]. Available from: [https://www.iarc.fr/wp-content/uploads/2018/09/pr263\\_E.pdf](https://www.iarc.fr/wp-content/uploads/2018/09/pr263_E.pdf) .
- Keramatinia A, Mohseny M, Akbari ME, Mosavi-Jarrahi A, Monfared ED, Amanpour F, et al. Determinants of survival of common childhood cancers in Iran. *J Res Med Sci.* 2018;23:101. - PMC - PubMed
- A 4 YEARS STUDY IN DIABETIC CHRONIC KIDNEY DISEASE PATIENTS ON KETOAMINOACIDS TREATMENT IN BU-CHAREST. RESULTS AFTER 1-YEAR FOLLOW-UP

- Smith K.B., Smith M.S. Obesity statistics. *Prim. Care Clin. Off. Pract.* 2016;43:121–135. doi: 10.1016/j.pop.2015.10.001. - DOI - PubMed
- Fleming T., Robinson M., Thomson B., Graetz N., Margono C., Mullany E.C., Biryukov S., Abbafati C., Abera S.F., Abraham J.P. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: A systematic analysis for the Global Burden of Disease Study. *Lancet.* 2013;384:766–781. - PMC - PubMed
- Noreik M., Maurmann M., Meier V., Becker I., Röhrig G., Polidori M.C., Schulz R.-J. Resting energy expenditure (REE) in an old-old population: Implications for metabolic stress. *Exp. Gerontol.* 2014;59:47–50. doi: 10.1016/j.exger.2014.06.009. - DOI - PubMed
- Matarese L.E. Indirect Calorimetry: Technical Aspects. *J. Am. Diet. Assoc.* 1997;97:S154–S160. doi: 10.1016/S0002-8223(97)00754-2. - DOI - PubMed
- Blasco Redondo R. Resting energy expenditure; assessment methods and applications. *Nutr. Hosp.* 2015;31:245–254. - PubMed
- Dixon JB, Dixon ME, Anderson ML, Schachter L, O'Brien PE. Daytime sleepiness in the obese: not as simple as obstructive sleep apnea. *Obesity.* 2007;15(10):2504–11. doi: 10.1038/oby.2007.297. - DOI - PubMed
- Dixon JB, Schachter LM, O'Brien PE. Sleep disturbance and obesity: changes following surgically induced weight loss. *Arch Intern Med.* 2001;161(1):102–6. doi: 10.1001/archinte.161.1.102. - DOI - PubMed
- Alosco ML, Gunstad J. The negative effects of obesity and poor glycemic control on cognitive function: a proposed model for possible mechanisms. *Curr Diab Rep.* 2014;14(6):495. doi: 10.1007/s11892-014-0495-z. - DOI - PMC - PubMed
- Whitmer RA, Gustafson DR, Barrett-Connor E, Haan MN, Gunderson EP, Yaffe K. Central obesity and increased risk of dementia more than three decades later. *Neurology.* 2008;71(14):1057–64. doi: 10.1212/01.wnl.0000306313.89165.ef. - DOI - PubMed
- Kopelman P. Health risks associated with overweight and obesity. *Obes Rev.* 2007;8(Suppl 1):13–7. doi: 10.1111/j.1467-789X.2007.00311.x. - DOI - PubMed
- Michalsen A. Prolonged fasting as a method of mood enhancement in chronic pain syndromes: a review of clinical evidence and mechanisms. *Current Pain and Headache Reports.* 2010;14(2):80–87. doi: 10.1007/s11916-010-0104-z. - DOI - PubMed
- Gill S., Panda S. A Smartphone App Reveals Erratic Diurnal Eating Patterns in Humans that Can Be Modulated for Health Benefits. *Cell Metabolism.* 2015;22(5):789–798. doi: 10.1016/j.cmet.2015.09.005. - DOI - PMC - PubMed
- Trepanowski J. F., Kroeger C. M., Barnosky A., et al. Effect of alternate-day fasting on weight loss, weight maintenance, and cardioprotection among metabolically healthy obese Adults. *JAMA Internal Medicine.* 2017;177(7):930–938. doi: 10.1001/jamainternmed.2017.0936. - DOI - PMC - PubMed
- Malinowski B., Zalewska K., Węsierska A., et al. Intermittent fasting in cardiovascular disorders-an overview. *Nutrients.* 2019;11(3):673–690. doi: 10.3390/nu11030673. - DOI - PMC - PubMed
- Zubrzycki A., Cierpka-Kmiec K., Kmiec Z., Wronska A. The role of low-calorie diets and intermittent fasting in the treatment of obesity and type-2 diabetes. *Journal of Physiology and Pharmacology.* 2018;69(5) - PubMed
- Efficacy and tolerability of the modified Atkins diet in children with drug-resistant genetic generalized epilepsy. *Kacker S, Nordli DR Jr, Phitsanu Wong C. Epileptic Disord.* 2022 Apr 1;24(2):295-301. doi: 10.1684/epd.2021.1389.PMID: 34825888
- Ketogenic dietary therapy utilization in Kenya: A qualitative exploration of dietitian's perceptions. *Samia P, Naanyu V, Helen Cross J, Idro R, Boon P, Wilmshurst J, Luchters S. Epilepsy Behav Rep.* 2024 Mar 20;26:100661. doi: 10.1016/j.ebr.2024.100661. eCollection 2024.PMID: 38560597
- Effect of Weekly Subcutaneous Semaglutide vs Daily Liraglutide on Body Weight in Adults With Overweight or Obesity Without Diabetes: The STEP 8 Randomized Clinical Trial. *Rubino DM, Greenway FL, Khalid U, O'Neil PM, Rosenstock J, Sørrig R, Wadden TA, Wizert A, Garvey WT; STEP 8 Investigators. JAMA.* 2022 Jan 11;327(2):138-150. doi: 10.1001/jama.2021.23619.PMID: 35015037
- Effect of Subcutaneous Semaglutide vs Placebo as an Adjunct to Intensive Behavioral Therapy on Body Weight in Adults With Overweight or Obesity: The STEP 3 Randomized Clinical Trial. *Wadden TA, Bailey TS, Billings LK, Davies M, Frias JP, Koroleva A, Lingvay I, O'Neil PM, Rubino DM, Skovgaard D, Walensteen SOR, Garvey WT; STEP 3 Investigators. JAMA.* 2021 Apr 13;325(14):1403-1413. doi: 10.1001/jama.2021.1831. PMID: 33625476
- Comparison of the Atkins, Ornish, Weight Watchers, and Zone diets for weight loss and heart disease risk reduction: a randomized trial.

Dansinger ML, Gleason JA, Griffith JL, Selker HP, Schaefer EJ. *JAMA*. 2005 Jan 5;293(1):43-53. doi: 10.1001/jama.293.1.43. PMID: 15632335

Assessment of food intakes for women adopting the high protein Dukan diet.

Wyka J, Malczyk E, Misiarz M, Zoloteńka-Synowiec M, Całyńiuk B, Baczyńska S. *Rocz Państw Zakł Hig*. 2015;66(2):137-42. PMID: 26024402

Fad diets and their effect on urinary stone formation.

Nouvenne A, Ticinesi A, Morelli I, Guida L, Borghi L, Meschi T. *Transl Androl Urol*. 2014 Sep;3(3):303-12. doi: 10.3978/j.issn.2223-4683.2014.06.01. PMID: 26816783

The Relationship between Modern Fad Diets and Kidney Stone Disease: A Systematic Review of Literature.

Barghouthy Y, Corrales M, Somani B. *Nutrients*. 2021 Nov 26;13(12):4270. doi: 10.3390/nu13124270. PMID: 34959822

Romero V., Akpınar H., Assimos D.G. Kidney Stones: A Global Picture of Prevalence, Incidence, and Associated Risk Factors. *Rev. Urol*. 2010;12:86–96. - PMC - PubMed

Sorokin I., Mamoulakis C., Miyazawa K., Rodgers A., Talati J., Lotan Y. Epidemiology of stone disease across the world. *World J. Urol*. 2017;35:1301–1320. doi: 10.1007/s00345-017-2008-6. - DOI - PubMed

Taylor E.N., Stampfer M.J., Curhan G.C. Obesity, Weight Gain, and the Risk of Kidney Stones. *JAMA*. 2005;293:455–462. doi: 10.1001/jama.293.4.455. - DOI - PubMed

Heilberg I.P., Goldfarb D. Optimum Nutrition for Kidney Stone Disease. *Adv. Chronic Kidney Dis*. 2013;20:165–174. doi: 10.1053/j.ackd.2012.12.001. - DOI - PubMed

Ziemba J.B., Matlaga B.R. Epidemiology and economics of nephrolithiasis. *Investig. Clin. Urol*. 2017;58:299–306. doi: 10.4111/icu.2017.58.5.299. - DOI - PMC - PubMed

Oh R., Gilani B., Uppaluri K.R. *StatPearls*. StatPearls Publishing; Treasure Island, FL, USA: 2021. Low Carbohydrate Diet. - PubMed

Khan S.R., Pearle M.S., Robertson W.G., Gambaro G., Canales B.K., Doizi S., Traxer O., Tiselius H.-G. Kidney stones. *Nat. Rev. Dis. Prim*. 2016;2:16008. doi: 10.1038/nrdp.2016.8. - DOI - PMC - PubMed

Gul Z., Monga M. Medical and Dietary Therapy for Kidney Stone Prevention. *Korean J. Urol*. 2014;55:775–779. doi: 10.4111/kju.2014.55.12.775. - DOI - PMC - PubMed

Institute of Medicine . *Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids*. National Academies Press; Washington, DC, USA: 2005. - DOI - PubMed

Freire R. Scientific evidence of diets for weight loss: Different macronutrient composition, intermittent fasting, and popular diets. *Nutrition*. 2019;69:110549. doi: 10.1016/j.nut.2019.07.001. - DOI - PubMed

Passos J.A., Vasconcelos-Silva P.R., Santos L.A.D.S. Ciclos de atenção a dietas da moda e tendências de busca na internet pelo Google trends. *Ciência Saúde Coletiva*. 2020;25:2615–2631. doi: 10.1590/1413-81232020257.23892018. - DOI - PubMed

Hite A.H., Berkowitz V.G., Berkowitz K. Low-Carbohydrate Diet Review. *Nutr. Clin. Pract*. 2011;26:300–308. doi: 10.1177/0884533611405791. - DOI - PubMed

Ludwig D.S., Ebbeling C.B. The Carbohydrate-Insulin Model of Obesity. *JAMA Intern. Med*. 2018;178:1098–1103. doi: 10.1001/jamainternmed.2018.2933. - DOI - PMC - PubMed

Johnston B.C., Kanters S., Bandayrel K., Wu P., Naji F., Siemieniuk R.A., Ball G.D.C., Busse J., Thorlund K., Guyatt G., et al. Comparison of Weight Loss among Named Diet Programs in Overweight and Obese Adults. *JAMA*. 2014;312:923–933. doi: 10.1001/jama.2014.10397. - DOI - PubMed

Ge L., Sadeghirad B., Ball G.D.C., Da Costa B.R., Hitchcock C.L., Svendrovski A., Kiflen R., Quadri K., Kwon H.Y., Karamouzian M., et al. Comparison of dietary macronutrient patterns of 14 popular named dietary programmes for weight and cardiovascular risk factor reduction in adults: Systematic review and network meta-analysis of randomised trials. *BMJ*. 2020;369:m696. doi: 10.1136/bmj.m696. - DOI - PMC - PubMed

Gardner C.D., Kiazand A., Alhassan S., Kim S., Stafford R.S., Balise R.R., Kraemer H.C., King A.C. Comparison of the Atkins, Zone, Ornish, and LEARN diets for change in weight and related risk factors among overweight premenopausal women: The A TO Z Weight Loss Study: A randomized trial. *JAMA*. 2007;297:969–977. doi: 10.1001/jama.297.9.969. - DOI - PubMed